

## Chapter 3. Differentiation

### 3.7. Derivatives of Inverse Functions and Logarithms

**Note.** Recall that the graph of a one-to-one function  $f$  and its inverse  $f^{-1}$  are mirror images of each other about the line  $y = x$ .




Figure 3.44 page 213

**Theorem 5.** If  $f$  has an interval  $I$  as domain and  $f'(x)$  exists and is never zero on  $I$ , then  $f^{-1}$  is differentiable at every point in its domain. The value of  $(f^{-1})'$  at a point  $b$  in the domain of  $f^{-1}$  is the reciprocal of the value of  $f'$  at the point  $a = f^{-1}(b)$ :

$$\left. \frac{df^{-1}}{dx} \right|_{x=b} = \frac{1}{\left. \frac{df}{dx} \right|_{x=f^{-1}(b)}}.$$

**Proof.** By definition of inverse function,  $f^{-1}(f(x)) = x$  for all  $x \in I$ . Differentiating this equation, we have by the Chain Rule:

$$\begin{aligned} \frac{d}{dx} [f^{-1}(f(x))] &= \frac{d}{dx}[x] \\ f^{-1}'(f(x))f'(x) &= 1 \\ f^{-1}'(f(x)) &= \frac{1}{f'(x)}. \end{aligned}$$

Plugging in  $x = f^{-1}(b)$ , we get the theorem.

*Q.E.D.*

**Example.** Page 221 number 8.

**Theorem.** For  $x > 0$  we have

$$\frac{d}{dx} [\ln x] = \frac{1}{x}.$$

If  $u = u(x)$  is a differentiable function of  $x$ , then for all  $x$  such that  $u(x) > 0$  we have

$$\frac{d}{dx} [\ln u] = \frac{d}{dx} [\ln u(x)] = \frac{1}{u} \left[ \frac{du}{dx} \right] = \frac{1}{u(x)} [u'(x)].$$

**Proof.** We know that  $f(x) = e^x$  is differentiable for all  $x$ , so we can apply Theorem 5 to find the derivative of  $f^{-1}(x) = \ln x$ :

$$\begin{aligned} \frac{d}{dx} [\ln x] &= (f^{-1})'(x) = \frac{1}{f'(f^{-1}(x))} \\ &= \frac{1}{e^{f^{-1}(x)}} \\ &= \frac{1}{e^{\ln x}} \\ &= \frac{1}{x}. \end{aligned}$$

By the Chain Rule

$$\frac{d}{dx} [\ln u(x)] = \frac{d}{du} [\ln u] \frac{du}{dx} = \frac{1}{u} \frac{du}{dx}.$$

*Q.E.D.*

**Note.** We can apply the previous theorem to show that  $\frac{d}{dx} [\ln |x|] = \frac{1}{x}$ .

**Recall.** For any numbers  $a > 0$  and for any real  $x$ ,  $a^x = e^{x \ln a}$ .

**Theorem.** If  $a > 0$  and  $u$  is a differentiable function of  $x$ , then  $a^u$  is a differentiable function of  $x$  and

$$\frac{d}{dx} [a^u] = a^u \ln a \left[ \frac{du}{dx} \right].$$

**Proof.** First

$$\begin{aligned} \frac{d}{dx} [a^x] &= \frac{d}{dx} [e^{x \ln a}] \\ &= e^{x \ln a} \left[ \frac{d}{dx} [x \ln a] \right] \\ &= a^x \ln a. \end{aligned}$$

Combining this result with the Chain Rule yields the theorem. *Q.E.D.*

**Note.** Notice that the previous theorem implies that  $\frac{d}{dx} [a^x] = a^x \ln a$ .

With  $a = e$ , we have the special case  $\frac{d}{dx} [e^x] = e^x(1) = e^x$ . This is

what is *natural* about  $e$ . When you first meet the natural exponential and logarithmic functions in algebra, it is hard to understand what is NATURAL about them. That is because the “natural-ness” is a calculus property (namely this differentiation property).

**Note.** We saw in section 3.2 that  $\frac{d}{dx}[a^x] = a^x \left( \lim_{h \rightarrow 0} \frac{a^h - 1}{h} \right)$ . We said then that the limit exists. We now see that the limit is  $\lim_{h \rightarrow 0} \frac{a^h - 1}{h} = \ln a$ . In particular, for  $a = e$ ,  $\lim_{h \rightarrow 0} \frac{e^h - 1}{h} = \ln e = 1$ .

**Example.** Page 222 number 70.

**Definition.** For any  $a > 0$ ,  $a \neq 1$ , define  $\log_a x = \frac{\ln x}{\ln a}$ . (This is called the *change of base* formula. See page 53.)

**Theorem.** Differentiating a logarithm base  $a$  gives:

$$\frac{d}{dx} [\log_a u] = \frac{1}{\ln a} \frac{1}{u} \frac{du}{dx}.$$

**Proof.** This follows easily:

$$\frac{d}{dx} [\log_a x] = \frac{d}{dx} \left[ \frac{\ln x}{\ln a} \right] = \frac{1}{\ln a} \frac{d}{dx} [\ln x] = \frac{1}{\ln a} \frac{1}{x}.$$

Combining this result with the Chain Rule gives the theorem. *Q.E.D.*

**Examples.** Page 222 numbers 74, and 80.

**Note.** We can, in fact, take the logarithm of a complicated function before differentiating it and then implicitly differentiate the result. This process is called *logarithmic differentiation*. It allows us to use the laws of logarithms instead of some of the complicated rules of differentiation.

**Example.** Page 222 number 90.

**Theorem. Power Rule (General Form).** If  $u$  is a positive differentiable function of  $x$  and  $n$  is any real number, then  $u^n$  is a differentiable function of  $x$  and

$$\frac{d}{dx} [u^n] = nu^{n-1} \frac{du}{dx}.$$

**Proof.** First,

$$\begin{aligned} \frac{d}{dx} [x^n] &= \frac{d}{dx} [e^{n \ln x}] \\ &= e^{n \ln x} \frac{d}{dx} [n \ln x] \text{ by the Chain Rule} \\ &= x^n \frac{n}{x} \\ &= nx^{n-1}. \end{aligned}$$

Combining this with the Chain Rule gives the result.

*Q.E.D.*

**Example.** Page 222 Example 72.

**Theorem 6.** We can find  $e$  as a limit:

$$e = \lim_{x \rightarrow 0} (1 + x)^{1/x}.$$

**Proof.** Let  $f(x) = \ln x$ . Then  $f'(x) = 1/x$  and  $f'(1) = 1$ . Now by the definition of derivative:

$$\begin{aligned} f'(1) &= \lim_{h \rightarrow 0} \frac{f(1+h) - f(1)}{h} = \lim_{x \rightarrow 0} \frac{(1+x) - f(1)}{x} \\ &= \lim_{x \rightarrow 0} \frac{\ln(1+x) - \ln 1}{x} = \lim_{x \rightarrow 0} \frac{1}{x} \ln(1+x) \\ &= \lim_{x \rightarrow 0} \ln(1+x)^{1/x} \\ &= \ln \left( \lim_{x \rightarrow 0} (1+x)^{1/x} \right) \text{ since } \ln x \text{ is continuous.} \end{aligned}$$

Therefore since  $f'(1) = 1$  we have

$$\ln \left( \lim_{x \rightarrow 0} (1+x)^{1/x} \right) = 1.$$

Since  $\ln e = 1$  and  $\ln x$  is one-to-one,

$$\lim_{x \rightarrow 0} (1+x)^{1/x} = e.$$

*Q.E.D.*

**Note.** We can use the previous theorem to find that

$$e \approx 2.7 \ 1828 \ 1828 \ 45 \ 90 \ 45 \ 9.$$