

Chapter 3. Differentiation

3.8. Inverse Trigonometric Functions

Recall. The six inverse trigonometric functions are defined as follows:

1. $y = \cos^{-1} x$ if and only if $\cos y = x$ and $y \in [0, \pi]$.
2. $y = \sin^{-1} x$ if and only if $\sin y = x$ and $y \in [-\pi/2, \pi/2]$.
3. $y = \tan^{-1} x$ if and only if $\tan y = x$ and $y \in (-\pi/2, \pi/2)$.
4. $y = \sec^{-1} x$ if and only if $\sec y = x$ and $y \in [0, \pi/2) \cup (\pi/2, \pi]$.
5. $y = \csc^{-1} x$ if and only if $\csc y = x$ and $y \in [-\pi/2, 0) \cup (0, \pi/2]$.
6. $y = \cot^{-1} x$ if and only if $\cot y = x$ and $y \in (0, \pi)$.

For all appropriate x values:

$$\sec^{-1} x = \cos^{-1}(1/x)$$

$$\csc^{-1} x = \sin^{-1}(1/x)$$

$$\cot^{-1} x = \pi/2 - \tan^{-1} x.$$

Note. The graphs of the six inverse trig functions are:




Figure 3.47 Page 223

Example. Page 230 numbers 4, 14, 28 and 38.

Theorem. We differentiate \sin^{-1} as follows:

$$\frac{d}{dx} [\sin^{-1} u] = \frac{1}{\sqrt{1-u^2}} \frac{du}{dx}$$

where $|u| < 1$.

Proof. We know that if $y = \sin^{-1} x$ then (for appropriate domain and range values) $\sin y = x$ and so by implicit differentiation

$$\begin{aligned} \frac{d}{dx} [\sin y] &= \frac{d}{dx} [x] \\ \cos y \left[\frac{dy}{dx} \right] &= 1 \\ \frac{dy}{dx} &= \frac{1}{\cos y}. \end{aligned}$$

Since we have restricted y to the interval $[-\pi/2, \pi/2]$, we know that $\cos y \geq 0$ and so $\cos y = +\sqrt{1 - (\sin y)^2} = \sqrt{1 - x^2}$. Making this substitution we get

$$\frac{d}{dx} [\sin^{-1} x] = \frac{1}{\sqrt{1-x^2}}.$$

The theorem then follows from the Chain Rule.

Q.E.D.

Example. Page 231 number 58.

Theorem. We differentiate \tan^{-1} as follows:

$$\frac{d}{dx} [\tan^{-1} u] = \frac{1}{1 + u^2} \left[\frac{du}{dx} \right].$$

Proof. We know that if $y = \tan^{-1} x$ then (for appropriate domain and range values) $\tan y = x$ and so by implicit differentiation

$$\begin{aligned} \frac{d}{dx} [\tan y] &= \frac{d}{dx} [x] \\ \sec^2 y \left[\frac{dy}{dx} \right] &= 1 \\ \frac{dy}{dx} &= \frac{1}{\sec^2 y} \\ &= \frac{1}{1 + (\tan y)^2} \\ &= \frac{1}{1 + x^2}. \end{aligned}$$

The theorem then follows from the Chain Rule.

Q.E.D.

Example. Page 231 number 62.

Theorem. We differentiate \sec^{-1} as follows:

$$\frac{d}{dx} [\sec^{-1} u] = \frac{1}{|u|\sqrt{u^2 - 1}} \frac{du}{dx}$$

where $|u| > 1$.

Proof. We know that if $y = \sec^{-1} x$ then (for appropriate domain and range values) $\sec y = x$ and so by implicit differentiation

$$\begin{aligned} \frac{d}{dx} [\sec y] &= \frac{d}{dx} [x] \\ \sec y \tan y \left[\frac{dy}{dx} \right] &= 1 \\ \frac{dy}{dx} &= \frac{1}{\sec y \tan y}. \end{aligned}$$

We now need to express this last expression in terms of x . First, $\sec y = x$ and $\tan y = \pm\sqrt{\sec^2 y - 1} = \pm\sqrt{x^2 - 1}$. Therefore we have

$$\frac{d}{dx} [\sec^{-1}] = \pm \frac{1}{x\sqrt{x^2 - 1}}.$$

Notice from the graph of $y = \sec^{-1} x$ above, that the slope of this function is positive where ever it is defined. So

$$\frac{d}{dx} [\sec^{-1} x] = \begin{cases} +\frac{1}{x\sqrt{x^2-1}} & \text{if } x > 1 \\ -\frac{1}{x\sqrt{x^2-1}} & \text{if } x < -1. \end{cases}$$

Notice that if $x > 1$ then $x = |x|$ and if $x < -1$ then $-x = |x|$. Therefore

$$\frac{d}{dx} [\sec^{-1} x] = \frac{1}{|x|\sqrt{x^2 - 1}}.$$

The Theorem then follows from the Chain Rule.

Q.E.D.

Note. We can use the following identities to differentiate the other three inverse trig functions:

$$\cos^{-1} x = \pi/2 - \sin^{-1} x$$

$$\cot^{-1} x = \pi/2 - \tan^{-1} x$$

$$\csc^{-1} x = \pi/2 - \sec^{-1} x$$

We then see that the only difference in the derivative of an inverse trig function and the derivative of the inverse of its cofunction is a negative sign. In summary, that is (Table 3.1 page 229):

$$1. \frac{d}{dx} [\sin^{-1} u] = \frac{du/dx}{\sqrt{1-u^2}}, |u| < 1$$

$$2. \frac{d}{dx} [\cos^{-1} u] = -\frac{du/dx}{\sqrt{1-u^2}}, |u| < 1$$

$$3. \frac{d}{dx} [\tan^{-1} u] = \frac{du/dx}{1+u^2}$$

$$4. \frac{d}{dx} [\cot^{-1} u] = -\frac{du/dx}{1+u^2}$$

$$5. \frac{d}{dx} [\sec^{-1} u] = \frac{du/dx}{|u|\sqrt{u^2-1}}, |u| > 1$$

$$6. \frac{d}{dx} [\csc^{-1} u] = \frac{-du/dx}{|u|\sqrt{u^2-1}}, |u| > 1$$

Example. Page 231 numbers 60 and 90.