

Chapter 6. Applications of Definite Integrals

6.3 Lengths of Plane Curves

Definition. If a curve C is described by the parametric equations $x = f(t)$, $y = g(t)$, $\alpha \leq t \leq \beta$, where f' and g' are continuous and not simultaneously zero on $[\alpha, \beta]$ and if C is traversed exactly once as t increases from α to β , then the length of C is

$$\begin{aligned} L &= \int_{\alpha}^{\beta} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt. \\ &= \int_{\alpha}^{\beta} \sqrt{(f'(t))^2 + (g'(t))^2} dt. \end{aligned}$$

Examples. Page 452 number 2.

Definition. Function f is *smooth* if its derivative is continuous. If f is smooth on $[a, b]$, the *length* of the curve $y = f(x)$ from a to b is the number

$$L = \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx.$$

If g is smooth on $[c, d]$, the *length* of the curve $x = g(y)$ from c to d is the number

$$L = \int_c^d \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy.$$




Figure 5.27, page 415 of Edition 10




Figure 5.29, page 418 of Edition 10

Example. Page 452 number 12.

Examples. Page 452 number 26.

Examples. Page 453 number 33.