Chapter 8. Techniques of Integration8.1 Integration by Parts

Theorem. (Integration by Parts)If u = u(x) and v = v(x) are differentiable functions of x, then we have

$$\int u\,dv = uv - \int v\,du.$$

Proof. By the Product Rule we have

$$\frac{d}{dx}\left[uv\right] = \left[\frac{du}{dx}\right]v + u\left[\frac{dv}{dx}\right].$$

Integrating both sides with respect to x and rearranging leads to the integral equation:

$$\int \left(u\frac{dv}{dx}\right) dx = \int \left(\frac{d}{dx}[uv]\right) dx - \int \left(v\frac{du}{dx}\right) dx$$
$$= uv - \int \left(v\frac{du}{dx}\right) dx.$$
Q.E.D.

Note. Applying Integration by Parts to a definite integral, we have:

$$\int_{v_1}^{v_2} u \, dv = (u_2 v_2 - u_1 v_1) - \int_{u_1}^{u_2} v \, du.$$

In terms of areas, this gives the following figure.




Figure 7.1 from Edition 10

Example. Page 455 Example 2. Evaluate $\int \ln x \, dx$.

Example. Page 456 Example 4. Evaluate

$$\int e^x \cos x \, dx.$$

(This is sort of weird!)

Example. Page 457 Example 5. Express $\int \cos^n x \, dx$ in terms of an integral of a lower power of $\cos x$. This is called a "reduction formula."

Examples. Page 459 number 4, page 460 numbers 34 and 42, page 461 number 68.