

Linear Algebra

Chapter 1. Vectors, Matrices, and Linear Systems

Section 1.3. Matrices and Their Algebra—Proofs of Theorems




Table of contents

- 1 Page 46 Number 16
- 2 Page 46 Number 6
- 3 Page 47 Number 33. Associativity of Matrix Multiplication
- 4 Example 1.3.A

Page 46 Number 16

Page 46 Number 16. Let $B = \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix}$ and $C = \begin{bmatrix} 2 & -1 \\ 0 & 6 \\ -3 & 2 \end{bmatrix}$.

Compute BC and CB .

Solution. First, notice that B is 2×3 and C is 3×2 , so both products actually exist, BC is 2×2 , and CB is 3×3 . By Definition 1.8, “Matrix Product,” we have

Page 46 Number 16

Page 46 Number 16. Let $B = \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix}$ and $C = \begin{bmatrix} 2 & -1 \\ 0 & 6 \\ -3 & 2 \end{bmatrix}$.

Compute BC and CB .

Solution. First, notice that B is 2×3 and C is 3×2 , so both products actually exist, BC is 2×2 , and CB is 3×3 . By Definition 1.8, “Matrix Product,” we have

$$BC = \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix} \begin{bmatrix} 2 & -1 \\ 0 & 6 \\ -3 & 2 \end{bmatrix}$$

Page 46 Number 16

Page 46 Number 16. Let $B = \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix}$ and $C = \begin{bmatrix} 2 & -1 \\ 0 & 6 \\ -3 & 2 \end{bmatrix}$.

Compute BC and CB .

Solution. First, notice that B is 2×3 and C is 3×2 , so both products actually exist, BC is 2×2 , and CB is 3×3 . By Definition 1.8, “Matrix Product,” we have

$$\begin{aligned} BC &= \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix} \begin{bmatrix} 2 & -1 \\ 0 & 6 \\ -3 & 2 \end{bmatrix} \\ &= \begin{bmatrix} (4)(2) + (1)(0) + (-2)(-3) & (4)(-1) + (1)(6) + (-2)(2) \\ (5)(2) + (-1)(0) + (3)(-3) & (5)(-1) + (-1)(6) + (3)(2) \end{bmatrix} \end{aligned}$$

Page 46 Number 16

Page 46 Number 16. Let $B = \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix}$ and $C = \begin{bmatrix} 2 & -1 \\ 0 & 6 \\ -3 & 2 \end{bmatrix}$.

Compute BC and CB .

Solution. First, notice that B is 2×3 and C is 3×2 , so both products actually exist, BC is 2×2 , and CB is 3×3 . By Definition 1.8, “Matrix Product,” we have

$$\begin{aligned}
 BC &= \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix} \begin{bmatrix} 2 & -1 \\ 0 & 6 \\ -3 & 2 \end{bmatrix} \\
 &= \begin{bmatrix} (4)(2) + (1)(0) + (-2)(-3) & (4)(-1) + (1)(6) + (-2)(2) \\ (5)(2) + (-1)(0) + (3)(-3) & (5)(-1) + (-1)(6) + (3)(2) \end{bmatrix} \\
 &= \begin{bmatrix} 8 + 0 + 6 & -4 + 6 - 4 \\ 10 + 0 - 9 & -5 - 6 + 6 \end{bmatrix} = \boxed{\begin{bmatrix} 14 & -2 \\ 1 & -5 \end{bmatrix}}.
 \end{aligned}$$

Page 46 Number 16

Page 46 Number 16. Let $B = \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix}$ and $C = \begin{bmatrix} 2 & -1 \\ 0 & 6 \\ -3 & 2 \end{bmatrix}$.

Compute BC and CB .

Solution. First, notice that B is 2×3 and C is 3×2 , so both products actually exist, BC is 2×2 , and CB is 3×3 . By Definition 1.8, “Matrix Product,” we have

$$\begin{aligned}
 BC &= \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix} \begin{bmatrix} 2 & -1 \\ 0 & 6 \\ -3 & 2 \end{bmatrix} \\
 &= \begin{bmatrix} (4)(2) + (1)(0) + (-2)(-3) & (4)(-1) + (1)(6) + (-2)(2) \\ (5)(2) + (-1)(0) + (3)(-3) & (5)(-1) + (-1)(6) + (3)(2) \end{bmatrix} \\
 &= \begin{bmatrix} 8 + 0 + 6 & -4 + 6 - 4 \\ 10 + 0 - 9 & -5 - 6 + 6 \end{bmatrix} = \boxed{\begin{bmatrix} 14 & -2 \\ 1 & -5 \end{bmatrix}}.
 \end{aligned}$$

Page 46 Number 16 (continued)

Page 46 Number 16. Let $B = \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix}$ and $C = \begin{bmatrix} 2 & -1 \\ 0 & 6 \\ -3 & 2 \end{bmatrix}$.

Compute BC and CB .

Solution (continued). By Definition 1.8, “Matrix Product,” we have

Page 46 Number 16 (continued)

Page 46 Number 16. Let $B = \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix}$ and $C = \begin{bmatrix} 2 & -1 \\ 0 & 6 \\ -3 & 2 \end{bmatrix}$.

Compute BC and CB .

Solution (continued). By Definition 1.8, “Matrix Product,” we have

$$CB = \begin{bmatrix} 2 & -1 \\ 0 & 6 \\ -3 & 2 \end{bmatrix} \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix}$$

Page 46 Number 16 (continued)

Page 46 Number 16. Let $B = \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix}$ and $C = \begin{bmatrix} 2 & -1 \\ 0 & 6 \\ -3 & 2 \end{bmatrix}$.

Compute BC and CB .

Solution (continued). By Definition 1.8, "Matrix Product," we have

$$\begin{aligned}
 CB &= \begin{bmatrix} 2 & -1 \\ 0 & 6 \\ -3 & 2 \end{bmatrix} \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix} \\
 &= \begin{bmatrix} (2)(4) + (-1)(5) & (2)(1) + (-1)(-1) & (2)(-2) + (-1)(3) \\ (0)(4) + (6)(5) & (0)(1) + (6)(-1) & (0)(-2) + (6)(3) \\ (-3)(4) + (2)(5) & (-3)(1) + (2)(-1) & (-3)(-2) + (2)(3) \end{bmatrix}
 \end{aligned}$$

Page 46 Number 16 (continued)

Page 46 Number 16. Let $B = \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix}$ and $C = \begin{bmatrix} 2 & -1 \\ 0 & 6 \\ -3 & 2 \end{bmatrix}$.

Compute BC and CB .

Solution (continued). By Definition 1.8, "Matrix Product," we have

$$\begin{aligned}
 CB &= \begin{bmatrix} 2 & -1 \\ 0 & 6 \\ -3 & 2 \end{bmatrix} \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix} \\
 &= \begin{bmatrix} (2)(4) + (-1)(5) & (2)(1) + (-1)(-1) & (2)(-2) + (-1)(3) \\ (0)(4) + (6)(5) & (0)(1) + (6)(-1) & (0)(-2) + (6)(3) \\ (-3)(4) + (2)(5) & (-3)(1) + (2)(-1) & (-3)(-2) + (2)(3) \end{bmatrix} \\
 &= \begin{bmatrix} 8 - 5 & 2 + 1 & -4 - 3 \\ 0 + 30 & 0 - 6 & 0 + 18 \\ -12 + 10 & -3 - 2 & 6 + 6 \end{bmatrix} = \boxed{\begin{bmatrix} 3 & 3 & -7 \\ 30 & -6 & 18 \\ -2 & -5 & 12 \end{bmatrix}}. \quad \square
 \end{aligned}$$

Page 46 Number 16 (continued)

Page 46 Number 16. Let $B = \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix}$ and $C = \begin{bmatrix} 2 & -1 \\ 0 & 6 \\ -3 & 2 \end{bmatrix}$.

Compute BC and CB .

Solution (continued). By Definition 1.8, "Matrix Product," we have

$$\begin{aligned}
 CB &= \begin{bmatrix} 2 & -1 \\ 0 & 6 \\ -3 & 2 \end{bmatrix} \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix} \\
 &= \begin{bmatrix} (2)(4) + (-1)(5) & (2)(1) + (-1)(-1) & (2)(-2) + (-1)(3) \\ (0)(4) + (6)(5) & (0)(1) + (6)(-1) & (0)(-2) + (6)(3) \\ (-3)(4) + (2)(5) & (-3)(1) + (2)(-1) & (-3)(-2) + (2)(3) \end{bmatrix} \\
 &= \begin{bmatrix} 8 - 5 & 2 + 1 & -4 - 3 \\ 0 + 30 & 0 - 6 & 0 + 18 \\ -12 + 10 & -3 - 2 & 6 + 6 \end{bmatrix} = \boxed{\begin{bmatrix} 3 & 3 & -7 \\ 30 & -6 & 18 \\ -2 & -5 & 12 \end{bmatrix}}. \quad \square
 \end{aligned}$$

Page 46 Number 6

Page 46 Number 6. Let $A = \begin{bmatrix} -2 & 1 & 3 \\ 4 & 0 & -1 \end{bmatrix}$ and

$B = \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix}$. Compute $4A - 2B$.

Solution. Notice that A and B are both 2×3 matrices so the sum actually exists. By Definition 1.9/1.10, "Matrix Sum and Scalar Multiplication,"

Page 46 Number 6

Page 46 Number 6. Let $A = \begin{bmatrix} -2 & 1 & 3 \\ 4 & 0 & -1 \end{bmatrix}$ and

$B = \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix}$. Compute $4A - 2B$.

Solution. Notice that A and B are both 2×3 matrices so the sum actually exists. By Definition 1.9/1.10, "Matrix Sum and Scalar Multiplication,"

$$4A - 2B = 4 \begin{bmatrix} -2 & 1 & 3 \\ 4 & 0 & -1 \end{bmatrix} - 2 \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix}$$

Page 46 Number 6

Page 46 Number 6. Let $A = \begin{bmatrix} -2 & 1 & 3 \\ 4 & 0 & -1 \end{bmatrix}$ and

$B = \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix}$. Compute $4A - 2B$.

Solution. Notice that A and B are both 2×3 matrices so the sum actually exists. By Definition 1.9/1.10, "Matrix Sum and Scalar Multiplication,"

$$\begin{aligned} 4A - 2B &= 4 \begin{bmatrix} -2 & 1 & 3 \\ 4 & 0 & -1 \end{bmatrix} - 2 \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix} \\ &= \begin{bmatrix} 4(-2) & 4(1) & 4(3) \\ 4(4) & 4(0) & 4(-1) \end{bmatrix} + \begin{bmatrix} -2(4) & -2(1) & -2(-2) \\ -2(5) & -2(-1) & -2(3) \end{bmatrix} \end{aligned}$$

Page 46 Number 6

Page 46 Number 6. Let $A = \begin{bmatrix} -2 & 1 & 3 \\ 4 & 0 & -1 \end{bmatrix}$ and

$B = \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix}$. Compute $4A - 2B$.

Solution. Notice that A and B are both 2×3 matrices so the sum actually exists. By Definition 1.9/1.10, "Matrix Sum and Scalar Multiplication,"

$$\begin{aligned} 4A - 2B &= 4 \begin{bmatrix} -2 & 1 & 3 \\ 4 & 0 & -1 \end{bmatrix} - 2 \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix} \\ &= \begin{bmatrix} 4(-2) & 4(1) & 4(3) \\ 4(4) & 4(0) & 4(-1) \end{bmatrix} + \begin{bmatrix} -2(4) & -2(1) & -2(-2) \\ -2(5) & -2(-1) & -2(3) \end{bmatrix} \\ &= \begin{bmatrix} -8 & 4 & 12 \\ 16 & 0 & -4 \end{bmatrix} + \begin{bmatrix} -8 & -2 & 4 \\ -10 & 2 & -6 \end{bmatrix} \end{aligned}$$

Page 46 Number 6

Page 46 Number 6. Let $A = \begin{bmatrix} -2 & 1 & 3 \\ 4 & 0 & -1 \end{bmatrix}$ and

$B = \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix}$. Compute $4A - 2B$.

Solution. Notice that A and B are both 2×3 matrices so the sum actually exists. By Definition 1.9/1.10, "Matrix Sum and Scalar Multiplication,"

$$\begin{aligned}
 4A - 2B &= 4 \begin{bmatrix} -2 & 1 & 3 \\ 4 & 0 & -1 \end{bmatrix} - 2 \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix} \\
 &= \begin{bmatrix} 4(-2) & 4(1) & 4(3) \\ 4(4) & 4(0) & 4(-1) \end{bmatrix} + \begin{bmatrix} -2(4) & -2(1) & -2(-2) \\ -2(5) & -2(-1) & -2(3) \end{bmatrix} \\
 &= \begin{bmatrix} -8 & 4 & 12 \\ 16 & 0 & -4 \end{bmatrix} + \begin{bmatrix} -8 & -2 & 4 \\ -10 & 2 & -6 \end{bmatrix} \\
 &= \begin{bmatrix} -8 + (-8) & 4 + (-2) & 12 + 4 \\ 16 + (-10) & 0 + 2 & -4 + (-6) \end{bmatrix} = \boxed{\begin{bmatrix} -16 & 2 & 16 \\ 6 & 2 & -10 \end{bmatrix}}
 \end{aligned}$$

□

Page 46 Number 6

Page 46 Number 6. Let $A = \begin{bmatrix} -2 & 1 & 3 \\ 4 & 0 & -1 \end{bmatrix}$ and

$B = \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix}$. Compute $4A - 2B$.

Solution. Notice that A and B are both 2×3 matrices so the sum actually exists. By Definition 1.9/1.10, "Matrix Sum and Scalar Multiplication,"

$$\begin{aligned}
 4A - 2B &= 4 \begin{bmatrix} -2 & 1 & 3 \\ 4 & 0 & -1 \end{bmatrix} - 2 \begin{bmatrix} 4 & 1 & -2 \\ 5 & -1 & 3 \end{bmatrix} \\
 &= \begin{bmatrix} 4(-2) & 4(1) & 4(3) \\ 4(4) & 4(0) & 4(-1) \end{bmatrix} + \begin{bmatrix} -2(4) & -2(1) & -2(-2) \\ -2(5) & -2(-1) & -2(3) \end{bmatrix} \\
 &= \begin{bmatrix} -8 & 4 & 12 \\ 16 & 0 & -4 \end{bmatrix} + \begin{bmatrix} -8 & -2 & 4 \\ -10 & 2 & -6 \end{bmatrix} \\
 &= \begin{bmatrix} -8 + (-8) & 4 + (-2) & 12 + 4 \\ 16 + (-10) & 0 + 2 & -4 + (-6) \end{bmatrix} = \boxed{\begin{bmatrix} -16 & 2 & 16 \\ 6 & 2 & -10 \end{bmatrix}}
 \end{aligned}$$

□

Page 47 Number 33

Page 47 Number 33. Let A , B , and C be matrices where the products $(AB)C$ and $A(BC)$ are defined. Then matrix multiplication is associative: $A(BC) = (AB)C$.

Proof. Let $A = [a_{ij}]$ be $m \times n$, $B = [b_{ij}]$ be $n \times s$, and $C = [c_{ij}]$ be $s \times t$. The (i, j) entry of BC is $\sum_{k=1}^s b_{ik}c_{kj}$ and so the (k, j) entry of BC is $\sum_{\ell=1}^s b_{k\ell}c_{\ell j}$.

Page 47 Number 33

Page 47 Number 33. Let A , B , and C be matrices where the products $(AB)C$ and $A(BC)$ are defined. Then matrix multiplication is associative: $A(BC) = (AB)C$.

Proof. Let $A = [a_{ij}]$ be $m \times n$, $B = [b_{ij}]$ be $n \times s$, and $C = [c_{ij}]$ be $s \times t$. The (i, j) entry of BC is $\sum_{k=1}^s b_{ik}c_{kj}$ and so the (k, j) entry of BC is $\sum_{\ell=1}^s b_{k\ell}c_{\ell j}$. Therefore the (i, j) entry of $A(BC)$ is

$$\sum_{k=1}^n a_{ik} \left(\sum_{\ell=1}^s b_{k\ell} c_{\ell j} \right) = \sum_{\ell=1}^s \left(\sum_{k=1}^n a_{ik} b_{k\ell} \right) c_{\ell j} = \sum_{k=1}^s \left(\sum_{\ell=1}^n a_{i\ell} b_{\ell k} \right) c_{kj}$$

where the second equality holds by interchanging dummy variables ℓ and k .

Page 47 Number 33

Page 47 Number 33. Let A , B , and C be matrices where the products $(AB)C$ and $A(BC)$ are defined. Then matrix multiplication is associative: $A(BC) = (AB)C$.

Proof. Let $A = [a_{ij}]$ be $m \times n$, $B = [b_{ij}]$ be $n \times s$, and $C = [c_{ij}]$ be $s \times t$. The (i, j) entry of BC is $\sum_{k=1}^s b_{ik}c_{kj}$ and so the (k, j) entry of BC is $\sum_{\ell=1}^s b_{k\ell}c_{\ell j}$. Therefore the (i, j) entry of $A(BC)$ is

$$\sum_{k=1}^n a_{ik} \left(\sum_{\ell=1}^s b_{k\ell} c_{\ell j} \right) = \sum_{\ell=1}^s \left(\sum_{k=1}^n a_{ik} b_{k\ell} \right) c_{\ell j} = \sum_{k=1}^s \left(\sum_{\ell=1}^n a_{i\ell} b_{\ell k} \right) c_{kj}$$

where the second equality holds by interchanging dummy variables ℓ and k . Now $\sum_{\ell=1}^n a_{i\ell} b_{\ell k}$ is the (i, k) entry of AB , and so the last sum is the (i, j) entry of $(AB)C$. Therefore $A(BC) = (AB)C$. \square

Page 47 Number 33

Page 47 Number 33. Let A , B , and C be matrices where the products $(AB)C$ and $A(BC)$ are defined. Then matrix multiplication is associative: $A(BC) = (AB)C$.

Proof. Let $A = [a_{ij}]$ be $m \times n$, $B = [b_{ij}]$ be $n \times s$, and $C = [c_{ij}]$ be $s \times t$. The (i, j) entry of BC is $\sum_{k=1}^s b_{ik}c_{kj}$ and so the (k, j) entry of BC is $\sum_{\ell=1}^s b_{k\ell}c_{\ell j}$. Therefore the (i, j) entry of $A(BC)$ is

$$\sum_{k=1}^n a_{ik} \left(\sum_{\ell=1}^s b_{k\ell} c_{\ell j} \right) = \sum_{\ell=1}^s \left(\sum_{k=1}^n a_{ik} b_{k\ell} \right) c_{\ell j} = \sum_{k=1}^s \left(\sum_{\ell=1}^n a_{i\ell} b_{\ell k} \right) c_{kj}$$

where the second equality holds by interchanging dummy variables ℓ and k . Now $\sum_{\ell=1}^n a_{i\ell} b_{\ell k}$ is the (i, k) entry of AB , and so the last sum is the (i, j) entry of $(AB)C$. Therefore $A(BC) = (AB)C$. \square

Example 1.3.A

Example 1.3.A. Show that $\mathcal{I}A = A\mathcal{I} = A$ for $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$ and \mathcal{I} is 3×3 .

Solution. By Definition 1.8, "Matrix Product," we have

Example 1.3.A

Example 1.3.A. Show that $\mathcal{I}A = A\mathcal{I} = A$ for $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$ and \mathcal{I} is 3×3 .

Solution. By Definition 1.8, “Matrix Product,” we have

$$\mathcal{I}A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$$

Example 1.3.A

Example 1.3.A. Show that $\mathcal{I}A = A\mathcal{I} = A$ for $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$ and \mathcal{I} is 3×3 .

Solution. By Definition 1.8, “Matrix Product,” we have

$$\mathcal{I}A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$$

$$= \begin{bmatrix} (1)(1) + (0)(4) + (0)(7) & (1)(2) + (0)(5) + (0)(8) & (1)(3) + (0)(6) + (0)(9) \\ (0)(1) + (1)(4) + (0)(7) & (0)(2) + (1)(5) + (0)(8) & (0)(3) + (1)(6) + (0)(9) \\ (0)(1) + (0)(4) + (1)(7) & (0)(2) + (0)(5) + (1)(8) & (0)(3) + (0)(6) + (1)(9) \end{bmatrix}$$

Example 1.3.A

Example 1.3.A. Show that $\mathcal{I}A = A\mathcal{I} = A$ for $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$ and \mathcal{I} is 3×3 .

Solution. By Definition 1.8, “Matrix Product,” we have

$$\mathcal{I}A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$$

$$= \begin{bmatrix} (1)(1) + (0)(4) + (0)(7) & (1)(2) + (0)(5) + (0)(8) & (1)(3) + (0)(6) + (0)(9) \\ (0)(1) + (1)(4) + (0)(7) & (0)(2) + (1)(5) + (0)(8) & (0)(3) + (1)(6) + (0)(9) \\ (0)(1) + (0)(4) + (1)(7) & (0)(2) + (0)(5) + (1)(8) & (0)(3) + (0)(6) + (1)(9) \end{bmatrix}$$

$$= \begin{bmatrix} 1+0+0 & 2+0+0 & 3+0+0 \\ 0+4+0 & 0+5+0 & 0+6+0 \\ 0+0+7 & 0+0+8 & 0+0+9 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} = A.$$

Example 1.3.A

Example 1.3.A. Show that $\mathcal{I}A = A\mathcal{I} = A$ for $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$ and \mathcal{I} is 3×3 .

Solution. By Definition 1.8, “Matrix Product,” we have

$$\mathcal{I}A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$$

$$= \begin{bmatrix} (1)(1) + (0)(4) + (0)(7) & (1)(2) + (0)(5) + (0)(8) & (1)(3) + (0)(6) + (0)(9) \\ (0)(1) + (1)(4) + (0)(7) & (0)(2) + (1)(5) + (0)(8) & (0)(3) + (1)(6) + (0)(9) \\ (0)(1) + (0)(4) + (1)(7) & (0)(2) + (0)(5) + (1)(8) & (0)(3) + (0)(6) + (1)(9) \end{bmatrix}$$

$$= \begin{bmatrix} 1 + 0 + 0 & 2 + 0 + 0 & 3 + 0 + 0 \\ 0 + 4 + 0 & 0 + 5 + 0 & 0 + 6 + 0 \\ 0 + 0 + 7 & 0 + 0 + 8 & 0 + 0 + 9 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} = A.$$

Example 1.3.A (continued)

Example 1.3.A. Show that $\mathcal{I}A = A\mathcal{I} = A$ for $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$ and \mathcal{I} is 3×3 .

Solution (continued). By Definition 1.8, “Matrix Product,” we have

Example 1.3.A (continued)

Example 1.3.A. Show that $\mathcal{I}A = A\mathcal{I} = A$ for $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$ and \mathcal{I} is 3×3 .

Solution (continued). By Definition 1.8, “Matrix Product,” we have

$$A\mathcal{I} = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Example 1.3.A (continued)

Example 1.3.A. Show that $\mathcal{I}A = A\mathcal{I} = A$ for $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$ and \mathcal{I} is 3×3 .

Solution (continued). By Definition 1.8, “Matrix Product,” we have

$$A\mathcal{I} = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} (1)(1) + (2)(0) + (3)(0) & (1)(0) + (2)(1) + (3)(0) & (1)(0) + (2)(0) + (3)(1) \\ (4)(1) + (5)(0) + (6)(0) & (4)(0) + (5)(1) + (6)(0) & (4)(0) + (5)(0) + (6)(1) \\ (7)(1) + (8)(0) + (9)(0) & (7)(0) + (8)(1) + (9)(0) & (7)(0) + (8)(0) + (9)(1) \end{bmatrix}$$

Example 1.3.A (continued)

Example 1.3.A. Show that $\mathcal{I}A = A\mathcal{I} = A$ for $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$ and \mathcal{I} is 3×3 .

Solution (continued). By Definition 1.8, “Matrix Product,” we have

$$A\mathcal{I} = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} (1)(1) + (2)(0) + (3)(0) & (1)(0) + (2)(1) + (3)(0) & (1)(0) + (2)(0) + (3)(1) \\ (4)(1) + (5)(0) + (6)(0) & (4)(0) + (5)(1) + (6)(0) & (4)(0) + (5)(0) + (6)(1) \\ (7)(1) + (8)(0) + (9)(0) & (7)(0) + (8)(1) + (9)(0) & (7)(0) + (8)(0) + (9)(1) \end{bmatrix}$$

$$= \begin{bmatrix} 1+0+0 & 0+2+0 & 0+0+3 \\ 4+0+0 & 0+5+0 & 0+0+6 \\ 7+0+0 & 0+8+0 & 0+0+9 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} = A.$$

□

Example 1.3.A (continued)

Example 1.3.A. Show that $\mathcal{I}A = A\mathcal{I} = A$ for $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$ and \mathcal{I} is 3×3 .

Solution (continued). By Definition 1.8, “Matrix Product,” we have

$$A\mathcal{I} = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} (1)(1) + (2)(0) + (3)(0) & (1)(0) + (2)(1) + (3)(0) & (1)(0) + (2)(0) + (3)(1) \\ (4)(1) + (5)(0) + (6)(0) & (4)(0) + (5)(1) + (6)(0) & (4)(0) + (5)(0) + (6)(1) \\ (7)(1) + (8)(0) + (9)(0) & (7)(0) + (8)(1) + (9)(0) & (7)(0) + (8)(0) + (9)(1) \end{bmatrix}$$

$$= \begin{bmatrix} 1 + 0 + 0 & 0 + 2 + 0 & 0 + 0 + 3 \\ 4 + 0 + 0 & 0 + 5 + 0 & 0 + 0 + 6 \\ 7 + 0 + 0 & 0 + 8 + 0 & 0 + 0 + 9 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} = A.$$

□