


# Linear Algebra

## Chapter 3. Vector Spaces

### Section 3.5. Inner-Product Spaces—Proofs of Theorems



# Table of contents

- 1 Page 236 Number 2
- 2 Page 236 Number 10
- 3 Theorem 3.11. Schwarz Inequality
- 4 Page 236 Number 12
- 5 Page 237 Number 18
- 6 Page 237 Number 20
- 7 Page 237 Number 24
- 8 Page 237 Number 26

## Page 236 Number 2

**Page 236 Number 2.** For  $[x_1, x_2], [y_1, y_2] \in \mathbb{R}^2$ , define the quantity  $\langle [x_1, x_2], [y_1, y_2] \rangle = x_1x_2 + y_1y_2$ . Does this satisfy the conditions for an inner product on  $\mathbb{R}^2$ ?

**Solution.** We test the parts of Definition 3.12, "Inner-Product Space."

## Page 236 Number 2

**Page 236 Number 2.** For  $[x_1, x_2], [y_1, y_2] \in \mathbb{R}^2$ , define the quantity  $\langle [x_1, x_2], [y_1, y_2] \rangle = x_1x_2 + y_1y_2$ . Does this satisfy the conditions for an inner product on  $\mathbb{R}^2$ ?

**Solution.** We test the parts of Definition 3.12, “Inner-Product Space.”  
P1.

$$\langle [y_1, y_2], [x_1, x_2] \rangle = y_1y_2 + x_1x_2 = x_1x_2 + y_1y_2 = \langle [x_1, x_2], [y_1, y_2] \rangle.$$

So P1 is satisfied.

## Page 236 Number 2

**Page 236 Number 2.** For  $[x_1, x_2], [y_1, y_2] \in \mathbb{R}^2$ , define the quantity  $\langle [x_1, x_2], [y_1, y_2] \rangle = x_1x_2 + y_1y_2$ . Does this satisfy the conditions for an inner product on  $\mathbb{R}^2$ ?

**Solution.** We test the parts of Definition 3.12, “Inner-Product Space.”  
**P1.**

$$\langle [y_1, y_2], [x_1, x_2] \rangle = y_1y_2 + x_1x_2 = x_1x_2 + y_1y_2 = \langle [x_1, x_2], [y_1, y_2] \rangle.$$

So P1 is satisfied.

**P2.** Let  $[z_1, z_2] \in \mathbb{R}^2$ . Then

$$\begin{aligned} \langle [z_1, z_2], [x_1, x_2] + [y_1, y_2] \rangle &= \langle [z_1, z_2], [x_1 + y_1, x_2 + y_2] \rangle \\ &= z_1z_2 + (x_1 + y_1)(x_2 + y_2) = z_1z_2 + (x_1x_2 + y_1x_2 + x_1y_2 + y_1y_2) \end{aligned}$$

and

$$\langle [z_1, z_2], [x_1, x_2] \rangle + \langle [z_1, z_2], [y_1, y_2] \rangle = (z_1z_2 + x_1x_2) + (z_1z_2 + y_1y_2).$$

So these don't appear to be the same.

# Page 236 Number 2

**Page 236 Number 2.** For  $[x_1, x_2], [y_1, y_2] \in \mathbb{R}^2$ , define the quantity  $\langle [x_1, x_2], [y_1, y_2] \rangle = x_1x_2 + y_1y_2$ . Does this satisfy the conditions for an inner product on  $\mathbb{R}^2$ ?

**Solution.** We test the parts of Definition 3.12, “Inner-Product Space.”  
**P1.**

$$\langle [y_1, y_2], [x_1, x_2] \rangle = y_1y_2 + x_1x_2 = x_1x_2 + y_1y_2 = \langle [x_1, x_2], [y_1, y_2] \rangle.$$

So P1 is satisfied.

**P2.** Let  $[z_1, z_2] \in \mathbb{R}^2$ . Then

$$\begin{aligned} \langle [z_1, z_2], [x_1, x_2] + [y_1, y_2] \rangle &= \langle [z_1, z_2], [x_1 + y_1, x_2 + y_2] \rangle \\ &= z_1z_2 + (x_1 + y_1)(x_2 + y_2) = z_1z_2 + (x_1x_2 + y_1x_2 + x_1y_2 + y_1y_2) \end{aligned}$$

and

$$\langle [z_1, z_2], [x_1, x_2] \rangle + \langle [z_1, z_2], [y_1, y_2] \rangle = (z_1z_2 + x_1x_2) + (z_1z_2 + y_1y_2).$$

So these don't appear to be the same.

## Page 236 Number 2 (continued)

**Page 236 Number 2.** For  $[x_1, x_2], [y_1, y_2] \in \mathbb{R}^2$ , define the quantity  $\langle [x_1, x_2], [y_1, y_2] \rangle = x_1x_2 + y_1y_2$ . Does this satisfy the conditions for an inner product on  $\mathbb{R}^2$ ?

**Solution (continued).** We pick particular numbers (inspired by these equations) to show that P2 does not hold. Consider  $[z_1, z_2] = [0, 0]$ ,  $[x_1, x_2] = [1, 1]$ , and  $[y_1, y_2] = [2, 2]$ . Then

$$\langle [0, 0], [1, 1] + [2, 2] \rangle = \langle [0, 0], [3, 3] \rangle = (0)(0) + (3)(3) = 9$$

and

$$\begin{aligned} \langle [0, 0], [1, 1] \rangle + \langle [0, 0], [2, 2] \rangle &= ((0)(0) + (1)(1)) + ((0)(0) + (2)(2)) \\ &= 1 + 4 = 5 \neq 9 = \langle [0, 0], [1, 1] + [2, 2] \rangle. \end{aligned}$$

So P2 does not hold and this is not an inner product in  $\mathbb{R}^2$ .  $\square$

## Page 236 Number 2 (continued)

**Page 236 Number 2.** For  $[x_1, x_2], [y_1, y_2] \in \mathbb{R}^2$ , define the quantity  $\langle [x_1, x_2], [y_1, y_2] \rangle = x_1x_2 + y_1y_2$ . Does this satisfy the conditions for an inner product on  $\mathbb{R}^2$ ?

**Solution (continued).** We pick particular numbers (inspired by these equations) to show that P2 does not hold. Consider  $[z_1, z_2] = [0, 0]$ ,  $[x_1, x_2] = [1, 1]$ , and  $[y_1, y_2] = [2, 2]$ . Then

$$\langle [0, 0], [1, 1] + [2, 2] \rangle = \langle [0, 0], [3, 3] \rangle = (0)(0) + (3)(3) = 9$$

and

$$\begin{aligned} \langle [0, 0], [1, 1] \rangle + \langle [0, 0], [2, 2] \rangle &= ((0)(0) + (1)(1)) + ((0)(0) + (2)(2)) \\ &= 1 + 4 = 5 \neq 9 = \langle [0, 0], [1, 1] + [2, 2] \rangle. \end{aligned}$$

So P2 does not hold and this is not an inner product in  $\mathbb{R}^2$ .  $\square$



# Page 236 Number 10

**Page 236 Number 10.** Let  $C_{a,b}$  be the vector space of all continuous real valued functions with domain  $a \leq x \leq b$  (see Note 3.1.A for a related vector space). Define  $\langle f, g \rangle = \int_a^b f(x)g(x) dx$ . Prove that  $\langle \cdot, \cdot \rangle$  is an inner product on  $C_{a,b}$ .

**Proof.** We show that  $\langle \cdot, \cdot \rangle$  satisfies P1–P4 of Definition 3.12, “Inner-Product Space.” Let  $f, g, h \in C_{a,b}$  and  $r \in \mathbb{R}$ .

## Page 236 Number 10

**Page 236 Number 10.** Let  $C_{a,b}$  be the vector space of all continuous real valued functions with domain  $a \leq x \leq b$  (see Note 3.1.A for a related vector space). Define  $\langle f, g \rangle = \int_a^b f(x)g(x) dx$ . Prove that  $\langle \cdot, \cdot \rangle$  is an inner product on  $C_{a,b}$ .

**Proof.** We show that  $\langle \cdot, \cdot \rangle$  satisfies P1–P4 of Definition 3.12, “Inner-Product Space.” Let  $f, g, h \in C_{a,b}$  and  $r \in \mathbb{R}$ . Then

**P1.**  $\langle f, g \rangle = \int_a^b f(x)g(x) dx = \int_a^b g(x)f(x) dx = \langle g, f \rangle$ .

## Page 236 Number 10

**Page 236 Number 10.** Let  $C_{a,b}$  be the vector space of all continuous real valued functions with domain  $a \leq x \leq b$  (see Note 3.1.A for a related vector space). Define  $\langle f, g \rangle = \int_a^b f(x)g(x) dx$ . Prove that  $\langle \cdot, \cdot \rangle$  is an inner product on  $C_{a,b}$ .

**Proof.** We show that  $\langle \cdot, \cdot \rangle$  satisfies P1–P4 of Definition 3.12, “Inner-Product Space.” Let  $f, g, h \in C_{a,b}$  and  $r \in \mathbb{R}$ . Then

**P1.**  $\langle f, g \rangle = \int_a^b f(x)g(x) dx = \int_a^b g(x)f(x) dx = \langle g, f \rangle.$

**P2.**  $\langle f, g + h \rangle = \int_a^b f(x)(g(x) + h(x)) dx = \int_a^b (f(x)g(x) + f(x)h(x)) dx$   
 $= \int_a^b f(x)g(x) dx + \int_a^b f(x)h(x) dx = \langle f, g \rangle + \langle f, h \rangle.$

## Page 236 Number 10

**Page 236 Number 10.** Let  $C_{a,b}$  be the vector space of all continuous real valued functions with domain  $a \leq x \leq b$  (see Note 3.1.A for a related vector space). Define  $\langle f, g \rangle = \int_a^b f(x)g(x) dx$ . Prove that  $\langle \cdot, \cdot \rangle$  is an inner product on  $C_{a,b}$ .

**Proof.** We show that  $\langle \cdot, \cdot \rangle$  satisfies P1–P4 of Definition 3.12, “Inner-Product Space.” Let  $f, g, h \in C_{a,b}$  and  $r \in \mathbb{R}$ . Then

$$\mathbf{P1.} \quad \langle f, g \rangle = \int_a^b f(x)g(x) dx = \int_a^b g(x)f(x) dx = \langle g, f \rangle.$$

$$\mathbf{P2.} \quad \langle f, g + h \rangle = \int_a^b f(x)(g(x) + h(x)) dx = \int_a^b (f(x)g(x) + f(x)h(x)) dx \\ = \int_a^b f(x)g(x) dx + \int_a^b f(x)h(x) dx = \langle f, g \rangle + \langle f, h \rangle.$$

$$\mathbf{P3.} \quad r\langle f, g \rangle = r \int_a^b f(x)g(x) dx = \int_a^b rf(x)g(x) dx$$

$$= \begin{cases} \int_a^b (rf(x))g(x) dx = \langle rf, g \rangle \\ \int_a^b f(x)(rg(x)) dx = \langle f, rg \rangle. \end{cases}$$

## Page 236 Number 10

**Page 236 Number 10.** Let  $C_{a,b}$  be the vector space of all continuous real valued functions with domain  $a \leq x \leq b$  (see Note 3.1.A for a related vector space). Define  $\langle f, g \rangle = \int_a^b f(x)g(x) dx$ . Prove that  $\langle \cdot, \cdot \rangle$  is an inner product on  $C_{a,b}$ .

**Proof.** We show that  $\langle \cdot, \cdot \rangle$  satisfies P1–P4 of Definition 3.12, “Inner-Product Space.” Let  $f, g, h \in C_{a,b}$  and  $r \in \mathbb{R}$ . Then

$$\mathbf{P1.} \quad \langle f, g \rangle = \int_a^b f(x)g(x) dx = \int_a^b g(x)f(x) dx = \langle g, f \rangle.$$

$$\mathbf{P2.} \quad \langle f, g + h \rangle = \int_a^b f(x)(g(x) + h(x)) dx = \int_a^b (f(x)g(x) + f(x)h(x)) dx \\ = \int_a^b f(x)g(x) dx + \int_a^b f(x)h(x) dx = \langle f, g \rangle + \langle f, h \rangle.$$

$$\mathbf{P3.} \quad r\langle f, g \rangle = r \int_a^b f(x)g(x) dx = \int_a^b rf(x)g(x) dx$$

$$= \begin{cases} \int_a^b (rf(x))g(x) dx = \langle rf, g \rangle \\ \int_a^b f(x)(rg(x)) dx = \langle f, rg \rangle. \end{cases}$$

## Page 236 Number 10 (continued)

**Page 236 Number 10.** Let  $C_{a,b}$  be the vector space of all continuous real valued functions with domain  $a \leq x \leq b$  (see Note 3.1.A for a related vector space). Define  $\langle f, g \rangle = \int_a^b f(x)g(x) dx$ . Prove that  $\langle \cdot, \cdot \rangle$  is an inner product on  $C_{a,b}$ .

**Proof (continued).**

**P4.**  $\langle f, f \rangle = \int_a^b f(x)f(x) dx = \int_a^b (f(x))^2 dx \geq 0$ , and

$\langle f, f \rangle = \int_a^b (f(x))^2 dx = 0$  if and only if  $(f(x))^2 = 0$  for all  $a \leq x \leq b$ , that is if and only if  $f(x) = 0$  for  $a \leq x \leq b$  (in  $C_{a,b}$  this means that  $f$  is the zero vector).

So  $\langle \cdot, \cdot \rangle$  satisfies P1-P4 and hence is an inner product on  $C_{a,b}$ . □

## Page 236 Number 10 (continued)

**Page 236 Number 10.** Let  $C_{a,b}$  be the vector space of all continuous real valued functions with domain  $a \leq x \leq b$  (see Note 3.1.A for a related vector space). Define  $\langle f, g \rangle = \int_a^b f(x)g(x) dx$ . Prove that  $\langle \cdot, \cdot \rangle$  is an inner product on  $C_{a,b}$ .

**Proof (continued).**

**P4.**  $\langle f, f \rangle = \int_a^b f(x)f(x) dx = \int_a^b (f(x))^2 dx \geq 0$ , and

$\langle f, f \rangle = \int_a^b (f(x))^2 dx = 0$  if and only if  $(f(x))^2 = 0$  for all  $a \leq x \leq b$ , that is if and only if  $f(x) = 0$  for  $a \leq x \leq b$  (in  $C_{a,b}$  this means that  $f$  is the zero vector).

So  $\langle \cdot, \cdot \rangle$  satisfies P1-P4 and hence is an inner product on  $C_{a,b}$ . □

# Theorem 3.11

## Theorem 3.11. Schwarz Inequality.

Let  $V$  be an inner-product space, and let  $\vec{v}, \vec{w} \in V$ . Then  $\langle \vec{v}, \vec{w} \rangle \leq \|\vec{v}\| \|\vec{w}\|$ .

**Proof.** Let  $r, s \in \mathbb{R}$ . Then we have:

$$\begin{aligned} \|r\vec{v} + s\vec{w}\|^2 &= \langle r\vec{v} + s\vec{w}, r\vec{v} + s\vec{w} \rangle \text{ by Definition 3.13, "norm"} \\ &= \langle r\vec{v} + s\vec{w}, r\vec{v} \rangle + \langle r\vec{v} + s\vec{w}, s\vec{w} \rangle \text{ by P2} \end{aligned}$$



# Theorem 3.11

## Theorem 3.11. Schwarz Inequality.

Let  $V$  be an inner-product space, and let  $\vec{v}, \vec{w} \in V$ . Then  $\langle \vec{v}, \vec{w} \rangle \leq \|\vec{v}\| \|\vec{w}\|$ .

**Proof.** Let  $r, s \in \mathbb{R}$ . Then we have:

$$\begin{aligned}
 \|r\vec{v} + s\vec{w}\|^2 &= \langle r\vec{v} + s\vec{w}, r\vec{v} + s\vec{w} \rangle \text{ by Definition 3.13, "norm"} \\
 &= \langle r\vec{v} + s\vec{w}, r\vec{v} \rangle + \langle r\vec{v} + s\vec{w}, s\vec{w} \rangle \text{ by P2} \\
 &= \langle r\vec{v}, r\vec{v} + s\vec{w} \rangle + \langle s\vec{w}, r\vec{v} + s\vec{w} \rangle \text{ by P1} \\
 &= \langle r\vec{v}, r\vec{v} \rangle + \langle r\vec{v}, s\vec{w} \rangle + \langle s\vec{w}, r\vec{v} \rangle + \langle s\vec{w}, s\vec{w} \rangle \text{ by P2}
 \end{aligned}$$

# Theorem 3.11

## Theorem 3.11. Schwarz Inequality.

Let  $V$  be an inner-product space, and let  $\vec{v}, \vec{w} \in V$ . Then  $\langle \vec{v}, \vec{w} \rangle \leq \|\vec{v}\| \|\vec{w}\|$ .

**Proof.** Let  $r, s \in \mathbb{R}$ . Then we have:

$$\begin{aligned}
 \|r\vec{v} + s\vec{w}\|^2 &= \langle r\vec{v} + s\vec{w}, r\vec{v} + s\vec{w} \rangle \text{ by Definition 3.13, "norm"} \\
 &= \langle r\vec{v} + s\vec{w}, r\vec{v} \rangle + \langle r\vec{v} + s\vec{w}, s\vec{w} \rangle \text{ by P2} \\
 &= \langle r\vec{v}, r\vec{v} + s\vec{w} \rangle + \langle s\vec{w}, r\vec{v} + s\vec{w} \rangle \text{ by P1} \\
 &= \langle r\vec{v}, r\vec{v} \rangle + \langle r\vec{v}, s\vec{w} \rangle + \langle s\vec{w}, r\vec{v} \rangle + \langle s\vec{w}, s\vec{w} \rangle \text{ by P2} \\
 &= r^2 \langle \vec{v}, \vec{v} \rangle + 2rs \langle \vec{v}, \vec{w} \rangle + s^2 \langle \vec{w}, \vec{w} \rangle \text{ by P1 and P3} \\
 &\geq 0 \text{ by P4.}
 \end{aligned}$$

Since this equation holds for all  $r, s \in \mathbb{R}$ , we are free to choose particular values of  $r$  and  $s$ . We choose  $r = \langle \vec{w}, \vec{w} \rangle$  and  $s = -\langle \vec{v}, \vec{w} \rangle$ .

# Theorem 3.11

## Theorem 3.11. Schwarz Inequality.

Let  $V$  be an inner-product space, and let  $\vec{v}, \vec{w} \in V$ . Then  $\langle \vec{v}, \vec{w} \rangle \leq \|\vec{v}\| \|\vec{w}\|$ .

**Proof.** Let  $r, s \in \mathbb{R}$ . Then we have:

$$\begin{aligned}
 \|r\vec{v} + s\vec{w}\|^2 &= \langle r\vec{v} + s\vec{w}, r\vec{v} + s\vec{w} \rangle \text{ by Definition 3.13, "norm"} \\
 &= \langle r\vec{v} + s\vec{w}, r\vec{v} \rangle + \langle r\vec{v} + s\vec{w}, s\vec{w} \rangle \text{ by P2} \\
 &= \langle r\vec{v}, r\vec{v} + s\vec{w} \rangle + \langle s\vec{w}, r\vec{v} + s\vec{w} \rangle \text{ by P1} \\
 &= \langle r\vec{v}, r\vec{v} \rangle + \langle r\vec{v}, s\vec{w} \rangle + \langle s\vec{w}, r\vec{v} \rangle + \langle s\vec{w}, s\vec{w} \rangle \text{ by P2} \\
 &= r^2 \langle \vec{v}, \vec{v} \rangle + 2rs \langle \vec{v}, \vec{w} \rangle + s^2 \langle \vec{w}, \vec{w} \rangle \text{ by P1 and P3} \\
 &\geq 0 \text{ by P4.}
 \end{aligned}$$

Since this equation holds for all  $r, s \in \mathbb{R}$ , we are free to choose particular values of  $r$  and  $s$ . We choose  $r = \langle \vec{w}, \vec{w} \rangle$  and  $s = -\langle \vec{v}, \vec{w} \rangle$ .

## Theorem 3.11 (continued)

**Proof (continued).** ...

$$r^2 \langle \vec{v}, \vec{v} \rangle + 2rs \langle \vec{v}, \vec{w} \rangle + s^2 \langle \vec{w}, \vec{w} \rangle \geq 0$$

with  $r = \langle \vec{w}, \vec{w} \rangle$  and  $s = -\langle \vec{v}, \vec{w} \rangle$  we have

$$\langle \vec{w}, \vec{w} \rangle^2 \langle \vec{v}, \vec{v} \rangle - 2 \langle \vec{w}, \vec{w} \rangle \langle \vec{v}, \vec{w} \rangle^2 + \langle \vec{v}, \vec{w} \rangle^2 \langle \vec{w}, \vec{w} \rangle$$

$$= \langle \vec{w}, \vec{w} \rangle^2 \langle \vec{v}, \vec{v} \rangle - \langle \vec{w}, \vec{w} \rangle \langle \vec{v}, \vec{w} \rangle^2 = \langle \vec{w}, \vec{w} \rangle [\langle \vec{w}, \vec{w} \rangle \langle \vec{v}, \vec{v} \rangle - \langle \vec{v}, \vec{w} \rangle^2] \geq 0. \quad (13)$$

## Theorem 3.11 (continued)

**Proof (continued).** ...

$$r^2 \langle \vec{v}, \vec{v} \rangle + 2rs \langle \vec{v}, \vec{w} \rangle + s^2 \langle \vec{w}, \vec{w} \rangle \geq 0$$

with  $r = \langle \vec{w}, \vec{w} \rangle$  and  $s = -\langle \vec{v}, \vec{w} \rangle$  we have

$$\langle \vec{w}, \vec{w} \rangle^2 \langle \vec{v}, \vec{v} \rangle - 2 \langle \vec{w}, \vec{w} \rangle \langle \vec{v}, \vec{w} \rangle^2 + \langle \vec{v}, \vec{w} \rangle^2 \langle \vec{w}, \vec{w} \rangle$$

$$= \langle \vec{w}, \vec{w} \rangle^2 \langle \vec{v}, \vec{v} \rangle - \langle \vec{w}, \vec{w} \rangle \langle \vec{v}, \vec{w} \rangle^2 = \langle \vec{w}, \vec{w} \rangle [\langle \vec{w}, \vec{w} \rangle \langle \vec{v}, \vec{v} \rangle - \langle \vec{v}, \vec{w} \rangle^2] \geq 0. \quad (13)$$

If  $\langle \vec{w}, \vec{w} \rangle = 0$  then  $\vec{w} = \vec{0}$  by Theorem 3.12 Part (P4), and the Schwarz Inequality is proven (since it reduces to  $0 \geq 0$ ). If  $\|\vec{w}\|^2 = \langle \vec{w}, \vec{w} \rangle \neq 0$ , then by the above inequality the other factor of inequality (13) must also be nonnegative:

$$\langle \vec{w}, \vec{w} \rangle \langle \vec{v}, \vec{v} \rangle - \langle \vec{v}, \vec{w} \rangle^2 \geq 0.$$

## Theorem 3.11 (continued)

**Proof (continued).** ...

$$r^2 \langle \vec{v}, \vec{v} \rangle + 2rs \langle \vec{v}, \vec{w} \rangle + s^2 \langle \vec{w}, \vec{w} \rangle \geq 0$$

with  $r = \langle \vec{w}, \vec{w} \rangle$  and  $s = -\langle \vec{v}, \vec{w} \rangle$  we have

$$\langle \vec{w}, \vec{w} \rangle^2 \langle \vec{v}, \vec{v} \rangle - 2 \langle \vec{w}, \vec{w} \rangle \langle \vec{v}, \vec{w} \rangle^2 + \langle \vec{v}, \vec{w} \rangle^2 \langle \vec{w}, \vec{w} \rangle$$

$$= \langle \vec{w}, \vec{w} \rangle^2 \langle \vec{v}, \vec{v} \rangle - \langle \vec{w}, \vec{w} \rangle \langle \vec{v}, \vec{w} \rangle^2 = \langle \vec{w}, \vec{w} \rangle [\langle \vec{w}, \vec{w} \rangle \langle \vec{v}, \vec{v} \rangle - \langle \vec{v}, \vec{w} \rangle^2] \geq 0. \quad (13)$$

If  $\langle \vec{w}, \vec{w} \rangle = 0$  then  $\vec{w} = \vec{0}$  by Theorem 3.12 Part (P4), and the Schwarz Inequality is proven (since it reduces to  $0 \geq 0$ ). If  $\|\vec{w}\|^2 = \langle \vec{w}, \vec{w} \rangle \neq 0$ , then by the above inequality the other factor of inequality (13) must also be nonnegative:

$$\langle \vec{w}, \vec{w} \rangle \langle \vec{v}, \vec{v} \rangle - \langle \vec{v}, \vec{w} \rangle^2 \geq 0.$$

Therefore

$$\langle \vec{v}, \vec{w} \rangle^2 \leq \langle \vec{v}, \vec{v} \rangle \langle \vec{w}, \vec{w} \rangle = \|\vec{v}\|^2 \|\vec{w}\|^2.$$

Taking square roots, we get the Schwarz Inequality. □

## Theorem 3.11 (continued)

**Proof (continued).** ...

$$r^2 \langle \vec{v}, \vec{v} \rangle + 2rs \langle \vec{v}, \vec{w} \rangle + s^2 \langle \vec{w}, \vec{w} \rangle \geq 0$$

with  $r = \langle \vec{w}, \vec{w} \rangle$  and  $s = -\langle \vec{v}, \vec{w} \rangle$  we have

$$\langle \vec{w}, \vec{w} \rangle^2 \langle \vec{v}, \vec{v} \rangle - 2 \langle \vec{w}, \vec{w} \rangle \langle \vec{v}, \vec{w} \rangle^2 + \langle \vec{v}, \vec{w} \rangle^2 \langle \vec{w}, \vec{w} \rangle$$

$$= \langle \vec{w}, \vec{w} \rangle^2 \langle \vec{v}, \vec{v} \rangle - \langle \vec{w}, \vec{w} \rangle \langle \vec{v}, \vec{w} \rangle^2 = \langle \vec{w}, \vec{w} \rangle [\langle \vec{w}, \vec{w} \rangle \langle \vec{v}, \vec{v} \rangle - \langle \vec{v}, \vec{w} \rangle^2] \geq 0. \quad (13)$$

If  $\langle \vec{w}, \vec{w} \rangle = 0$  then  $\vec{w} = \vec{0}$  by Theorem 3.12 Part (P4), and the Schwarz Inequality is proven (since it reduces to  $0 \geq 0$ ). If  $\|\vec{w}\|^2 = \langle \vec{w}, \vec{w} \rangle \neq 0$ , then by the above inequality the other factor of inequality (13) must also be nonnegative:

$$\langle \vec{w}, \vec{w} \rangle \langle \vec{v}, \vec{v} \rangle - \langle \vec{v}, \vec{w} \rangle^2 \geq 0.$$

Therefore

$$\langle \vec{v}, \vec{w} \rangle^2 \leq \langle \vec{v}, \vec{v} \rangle \langle \vec{w}, \vec{w} \rangle = \|\vec{v}\|^2 \|\vec{w}\|^2.$$

Taking square roots, we get the Schwarz Inequality. □

## Page 236 Number 12

**Page 236 Number 12.** Show that  $\sin x$  and  $\cos x$  are orthogonal functions in the vector space  $C_{0,\pi}$  of continuous functions with domain  $0 \leq x \leq \pi$  where the inner product is defined as  $\langle f, g \rangle = \int_0^\pi f(x)g(x) dx$  (see Exercise 10).

**Solution.** We have (by  $u$ -substitution with  $u = \sin x$  and  $du = \cos x$ ):

$$\begin{aligned}\langle \cos x, \sin x \rangle &= \int_0^\pi \cos x \sin x dx \\ &= \frac{1}{2} \sin^2 x \Big|_0^\pi = \frac{1}{2} \sin^2 \pi - \frac{1}{2} \sin^2 0 = 0 - 0 = 0.\end{aligned}$$



## Page 236 Number 12

**Page 236 Number 12.** Show that  $\sin x$  and  $\cos x$  are orthogonal functions in the vector space  $C_{0,\pi}$  of continuous functions with domain  $0 \leq x \leq \pi$  where the inner product is defined as  $\langle f, g \rangle = \int_0^\pi f(x)g(x) dx$  (see Exercise 10).

**Solution.** We have (by  $u$ -substitution with  $u = \sin x$  and  $du = \cos x$ ):

$$\begin{aligned}\langle \cos x, \sin x \rangle &= \int_0^\pi \cos x \sin x dx \\ &= \frac{1}{2} \sin^2 x \Big|_0^\pi = \frac{1}{2} \sin^2 \pi - \frac{1}{2} \sin^2 0 = 0 - 0 = 0.\end{aligned}$$

So by the definition of orthogonal in an inner-product space,  $\cos x$  and  $\sin x$  are orthogonal.  $\square$

## Page 236 Number 12

**Page 236 Number 12.** Show that  $\sin x$  and  $\cos x$  are orthogonal functions in the vector space  $C_{0,\pi}$  of continuous functions with domain  $0 \leq x \leq \pi$  where the inner product is defined as  $\langle f, g \rangle = \int_0^\pi f(x)g(x) dx$  (see Exercise 10).

**Solution.** We have (by  $u$ -substitution with  $u = \sin x$  and  $du = \cos x$ ):

$$\begin{aligned}\langle \cos x, \sin x \rangle &= \int_0^\pi \cos x \sin x dx \\ &= \frac{1}{2} \sin^2 x \Big|_0^\pi = \frac{1}{2} \sin^2 \pi - \frac{1}{2} \sin^2 0 = 0 - 0 = 0.\end{aligned}$$

So by the definition of orthogonal in an inner-product space,  $\cos x$  and  $\sin x$  are orthogonal.  $\square$

# Page 237 Number 18

**Page 237 Number 18.** For vectors  $\vec{u}$ ,  $\vec{v}$ , and  $\vec{w}$  in an inner-product space and for scalars  $r$  and  $s$ , prove that if  $\vec{w}$  is perpendicular to both  $\vec{u}$  and  $\vec{v}$  then  $\vec{w}$  is perpendicular to  $r\vec{u} + s\vec{v}$ .

**Proof.** Since  $\vec{w}$  is perpendicular to both  $\vec{u}$  and  $\vec{v}$  then by the definition of perpendicular (or orthogonal),  $\langle \vec{w}, \vec{u} \rangle = \langle \vec{w}, \vec{v} \rangle = 0$ .

## Page 237 Number 18

**Page 237 Number 18.** For vectors  $\vec{u}$ ,  $\vec{v}$ , and  $\vec{w}$  in an inner-product space and for scalars  $r$  and  $s$ , prove that if  $\vec{w}$  is perpendicular to both  $\vec{u}$  and  $\vec{v}$  then  $\vec{w}$  is perpendicular to  $r\vec{u} + s\vec{v}$ .

**Proof.** Since  $\vec{w}$  is perpendicular to both  $\vec{u}$  and  $\vec{v}$  then by the definition of perpendicular (or orthogonal),  $\langle \vec{w}, \vec{u} \rangle = \langle \vec{w}, \vec{v} \rangle = 0$ . So

$$\begin{aligned}\langle \vec{w}, r\vec{u} + s\vec{v} \rangle &= \langle \vec{w}, r\vec{u} \rangle + \langle \vec{w}, s\vec{v} \rangle \text{ by P2} \\ &= r\langle \vec{w}, \vec{u} \rangle + s\langle \vec{w}, \vec{v} \rangle \text{ by P3} \\ &= r(0) + s(0) = 0.\end{aligned}$$

So  $\vec{w}$  is perpendicular to  $r\vec{u} + s\vec{v}$ , as claimed. □

## Page 237 Number 18

**Page 237 Number 18.** For vectors  $\vec{u}$ ,  $\vec{v}$ , and  $\vec{w}$  in an inner-product space and for scalars  $r$  and  $s$ , prove that if  $\vec{w}$  is perpendicular to both  $\vec{u}$  and  $\vec{v}$  then  $\vec{w}$  is perpendicular to  $r\vec{u} + s\vec{v}$ .

**Proof.** Since  $\vec{w}$  is perpendicular to both  $\vec{u}$  and  $\vec{v}$  then by the definition of perpendicular (or orthogonal),  $\langle \vec{w}, \vec{u} \rangle = \langle \vec{w}, \vec{v} \rangle = 0$ . So

$$\begin{aligned}\langle \vec{w}, r\vec{u} + s\vec{v} \rangle &= \langle \vec{w}, r\vec{u} \rangle + \langle \vec{w}, s\vec{v} \rangle \text{ by P2} \\ &= r\langle \vec{w}, \vec{u} \rangle + s\langle \vec{w}, \vec{v} \rangle \text{ by P3} \\ &= r(0) + s(0) = 0.\end{aligned}$$

So  $\vec{w}$  is perpendicular to  $r\vec{u} + s\vec{v}$ , as claimed. □

# Page 237 Number 20

**Page 237 Number 20.** Let  $V$  be an inner-product space and let  $S$  be a subset of  $V$ . Prove that

$$S^\perp = \{\vec{v} \in V \mid \vec{v} \text{ is orthogonal to each vector in } S\}$$

is a subspace of  $V$ .  $S^\perp$  is called the *perp space* of set  $S$ .

**Proof.** We apply Theorem 3.2, “Test for a Subspace,” and test  $S^\perp$  for closure under vector addition and closure under scalar multiplication. Let  $\vec{v}, \vec{w} \in S^\perp$  and let  $r \in \mathbb{R}$  be a scalar.

## Page 237 Number 20

**Page 237 Number 20.** Let  $V$  be an inner-product space and let  $S$  be a subset of  $V$ . Prove that

$$S^\perp = \{\vec{v} \in V \mid \vec{v} \text{ is orthogonal to each vector in } S\}$$

is a subspace of  $V$ .  $S^\perp$  is called the *perp space* of set  $S$ .

**Proof.** We apply Theorem 3.2, “Test for a Subspace,” and test  $S^\perp$  for closure under vector addition and closure under scalar multiplication. Let  $\vec{v}, \vec{w} \in S^\perp$  and let  $r \in \mathbb{R}$  be a scalar. Then by the definition of  $S^\perp$ , for every  $\vec{s} \in S$  we have  $\langle \vec{v}, \vec{s} \rangle = 0$  and  $\langle \vec{w}, \vec{s} \rangle = 0$ .

# Page 237 Number 20

**Page 237 Number 20.** Let  $V$  be an inner-product space and let  $S$  be a subset of  $V$ . Prove that

$$S^\perp = \{\vec{v} \in V \mid \vec{v} \text{ is orthogonal to each vector in } S\}$$

is a subspace of  $V$ .  $S^\perp$  is called the *perp space* of set  $S$ .

**Proof.** We apply Theorem 3.2, “Test for a Subspace,” and test  $S^\perp$  for closure under vector addition and closure under scalar multiplication. Let  $\vec{v}, \vec{w} \in S^\perp$  and let  $r \in \mathbb{R}$  be a scalar. Then by the definition of  $S^\perp$ , for every  $\vec{s} \in S$  we have  $\langle \vec{v}, \vec{s} \rangle = 0$  and  $\langle \vec{w}, \vec{s} \rangle = 0$ . Now

$$\begin{aligned} \langle \vec{v} + \vec{w}, \vec{s} \rangle &= \langle \vec{s}, \vec{v} + \vec{w} \rangle \text{ by P1} \\ &= \langle \vec{s}, \vec{v} \rangle + \langle \vec{s}, \vec{w} \rangle \text{ by P2} \\ &= 0 + 0 = 0 \end{aligned}$$

for every  $\vec{s} \in S$  and hence  $\vec{v} + \vec{w} \in S^\perp$  and  $S^\perp$  is closed under vector addition.



# Page 237 Number 20

**Page 237 Number 20.** Let  $V$  be an inner-product space and let  $S$  be a subset of  $V$ . Prove that

$$S^\perp = \{\vec{v} \in V \mid \vec{v} \text{ is orthogonal to each vector in } S\}$$

is a subspace of  $V$ .  $S^\perp$  is called the *perp space* of set  $S$ .

**Proof.** We apply Theorem 3.2, “Test for a Subspace,” and test  $S^\perp$  for closure under vector addition and closure under scalar multiplication. Let  $\vec{v}, \vec{w} \in S^\perp$  and let  $r \in \mathbb{R}$  be a scalar. Then by the definition of  $S^\perp$ , for every  $\vec{s} \in S$  we have  $\langle \vec{v}, \vec{s} \rangle = 0$  and  $\langle \vec{w}, \vec{s} \rangle = 0$ . Now

$$\begin{aligned} \langle \vec{v} + \vec{w}, \vec{s} \rangle &= \langle \vec{s}, \vec{v} + \vec{w} \rangle \text{ by P1} \\ &= \langle \vec{s}, \vec{v} \rangle + \langle \vec{s}, \vec{w} \rangle \text{ by P2} \\ &= 0 + 0 = 0 \end{aligned}$$

for every  $\vec{s} \in S$  and hence  $\vec{v} + \vec{w} \in S^\perp$  and  $S^\perp$  is closed under vector addition.

# Page 237 Number 20 (continued)

**Page 237 Number 20.** Let  $V$  be an inner-product space and let  $S$  be a subset of  $V$ . Prove that

$$S^\perp = \{\vec{v} \in V \mid \vec{v} \text{ is orthogonal to each vector in } S\}$$

is a subspace of  $V$ .  $S^\perp$  is called the *perp space* of set  $S$ .

**Proof (continued).** Next,

$$\begin{aligned} \langle r\vec{v}, \vec{s} \rangle &= r\langle \vec{v}, \vec{s} \rangle \text{ by P3} \\ &= r(0) = 0 \end{aligned}$$

for every  $\vec{s} \in S$  and hence  $r\vec{v} \in S^\perp$  and  $S^\perp$  is closed under scalar multiplication. Therefore Theorem 3.2, "Test for Subspace," implies that  $S^\perp$  is a subspace of  $V$ . □

# Page 237 Number 20 (continued)

**Page 237 Number 20.** Let  $V$  be an inner-product space and let  $S$  be a subset of  $V$ . Prove that

$$S^\perp = \{\vec{v} \in V \mid \vec{v} \text{ is orthogonal to each vector in } S\}$$

is a subspace of  $V$ .  $S^\perp$  is called the *perp space* of set  $S$ .

**Proof (continued).** Next,

$$\begin{aligned} \langle r\vec{v}, \vec{s} \rangle &= r\langle \vec{v}, \vec{s} \rangle \text{ by P3} \\ &= r(0) = 0 \end{aligned}$$

for every  $\vec{s} \in S$  and hence  $r\vec{v} \in S^\perp$  and  $S^\perp$  is closed under scalar multiplication. Therefore Theorem 3.2, “Test for Subspace,” implies that  $S^\perp$  is a subspace of  $V$ . □

## Page 237 Number 24

**Page 237 Number 24.** Use the Triangle Inequality to prove that for any  $\vec{v}, \vec{w}$  in an inner-product space  $V$ ,  $\|\vec{v} - \vec{w}\| \leq \|\vec{v}\| + \|\vec{w}\|$ .

**Proof.** Let  $\vec{v}, \vec{w} \in V$ . Then  $-\vec{w} \in V$  and so we consider the vector sum  $\vec{v} + (-\vec{w})$ . The Triangle Inequality implies that  $\|\vec{v} + (-\vec{w})\| \leq \|\vec{v}\| + \|-\vec{w}\|$ .

## Page 237 Number 24

**Page 237 Number 24.** Use the Triangle Inequality to prove that for any  $\vec{v}, \vec{w}$  in an inner-product space  $V$ ,  $\|\vec{v} - \vec{w}\| \leq \|\vec{v}\| + \|\vec{w}\|$ .

**Proof.** Let  $\vec{v}, \vec{w} \in V$ . Then  $-\vec{w} \in V$  and so we consider the vector sum  $\vec{v} + (-\vec{w})$ . The Triangle Inequality implies that  $\|\vec{v} + (-\vec{w})\| \leq \|\vec{v}\| + \|\vec{-w}\|$ . Now

$$\begin{aligned} \|\vec{-w}\| &= \sqrt{\langle \vec{-w}, \vec{-w} \rangle} \text{ by Definition 3.13,} \\ &\quad \text{Magnitude or Norm of a Vector''} \\ &= \sqrt{(-1)\langle \vec{w}, \vec{-w} \rangle} = \sqrt{(-1)(-1)\langle \vec{w}, \vec{w} \rangle} \text{ by P3} \\ &= \sqrt{\langle \vec{w}, \vec{w} \rangle} = \|\vec{w}\|. \end{aligned}$$

## Page 237 Number 24

**Page 237 Number 24.** Use the Triangle Inequality to prove that for any  $\vec{v}, \vec{w}$  in an inner-product space  $V$ ,  $\|\vec{v} - \vec{w}\| \leq \|\vec{v}\| + \|\vec{w}\|$ .

**Proof.** Let  $\vec{v}, \vec{w} \in V$ . Then  $-\vec{w} \in V$  and so we consider the vector sum  $\vec{v} + (-\vec{w})$ . The Triangle Inequality implies that  $\|\vec{v} + (-\vec{w})\| \leq \|\vec{v}\| + \|-\vec{w}\|$ . Now

$$\begin{aligned} \|-\vec{w}\| &= \sqrt{\langle -\vec{w}, -\vec{w} \rangle} \text{ by Definition 3.13,} \\ &\quad \text{Magnitude or Norm of a Vector''} \\ &= \sqrt{(-1)\langle \vec{w}, -\vec{w} \rangle} = \sqrt{(-1)(-1)\langle \vec{w}, \vec{w} \rangle} \text{ by P3} \\ &= \sqrt{\langle \vec{w}, \vec{w} \rangle} = \|\vec{w}\|. \end{aligned}$$

So  $\|\vec{v} - \vec{w}\| = \|\vec{v} + (-\vec{w})\| \leq \|\vec{v}\| + \|-\vec{w}\| = \|\vec{v}\| + \|\vec{w}\|$ , as claimed.  $\square$

## Page 237 Number 24

**Page 237 Number 24.** Use the Triangle Inequality to prove that for any  $\vec{v}, \vec{w}$  in an inner-product space  $V$ ,  $\|\vec{v} - \vec{w}\| \leq \|\vec{v}\| + \|\vec{w}\|$ .

**Proof.** Let  $\vec{v}, \vec{w} \in V$ . Then  $-\vec{w} \in V$  and so we consider the vector sum  $\vec{v} + (-\vec{w})$ . The Triangle Inequality implies that  $\|\vec{v} + (-\vec{w})\| \leq \|\vec{v}\| + \|-\vec{w}\|$ . Now

$$\begin{aligned} \|-\vec{w}\| &= \sqrt{\langle -\vec{w}, -\vec{w} \rangle} \text{ by Definition 3.13,} \\ &\quad \text{Magnitude or Norm of a Vector''} \\ &= \sqrt{(-1)\langle \vec{w}, -\vec{w} \rangle} = \sqrt{(-1)(-1)\langle \vec{w}, \vec{w} \rangle} \text{ by P3} \\ &= \sqrt{\langle \vec{w}, \vec{w} \rangle} = \|\vec{w}\|. \end{aligned}$$

So  $\|\vec{v} - \vec{w}\| = \|\vec{v} + (-\vec{w})\| \leq \|\vec{v}\| + \|-\vec{w}\| = \|\vec{v}\| + \|\vec{w}\|$ , as claimed.  $\square$

## Page 237 Number 26

**Page 237 Number 26.** Let  $\vec{v}$  and  $\vec{w}$  be vectors in an inner-product space  $V$ . Show that  $\|\vec{v}\|\vec{w} + \|\vec{w}\|\vec{v}$  is perpendicular to  $\|\vec{v}\|\vec{w} - \|\vec{w}\|\vec{v}$ .

**Solution.** Consider

$$\langle \|\vec{v}\|\vec{w} + \|\vec{w}\|\vec{v}, \|\vec{v}\|\vec{w} - \|\vec{w}\|\vec{v} \rangle$$



# Page 237 Number 26

**Page 237 Number 26.** Let  $\vec{v}$  and  $\vec{w}$  be vectors in an inner-product space  $V$ . Show that  $\|\vec{v}\|\vec{w} + \|\vec{w}\|\vec{v}$  is perpendicular to  $\|\vec{v}\|\vec{w} - \|\vec{w}\|\vec{v}$ .

**Solution.** Consider

$$\begin{aligned} & \langle \|\vec{v}\|\vec{w} + \|\vec{w}\|\vec{v}, \|\vec{v}\|\vec{w} - \|\vec{w}\|\vec{v} \rangle \\ = & \langle \|\vec{v}\|\vec{w} + \|\vec{w}\|\vec{v}, \|\vec{v}\|\vec{w} \rangle + \langle \|\vec{v}\|\vec{w} + \|\vec{w}\|\vec{v}, -\|\vec{w}\|\vec{v} \rangle \text{ by P2} \\ = & \langle \|\vec{v}\|\vec{w}, \|\vec{v}\|\vec{w} + \|\vec{w}\|\vec{v} \rangle + \langle -\|\vec{w}\|\vec{v}, \|\vec{v}\|\vec{w} + \|\vec{w}\|\vec{v} \rangle \text{ by P1} \end{aligned}$$

## Page 237 Number 26

**Page 237 Number 26.** Let  $\vec{v}$  and  $\vec{w}$  be vectors in an inner-product space  $V$ . Show that  $\|\vec{v}\|\vec{w} + \|\vec{w}\|\vec{v}$  is perpendicular to  $\|\vec{v}\|\vec{w} - \|\vec{w}\|\vec{v}$ .

**Solution.** Consider

$$\begin{aligned}
 & \langle \|\vec{v}\|\vec{w} + \|\vec{w}\|\vec{v}, \|\vec{v}\|\vec{w} - \|\vec{w}\|\vec{v} \rangle \\
 = & \langle \|\vec{v}\|\vec{w} + \|\vec{w}\|\vec{v}, \|\vec{v}\|\vec{w} \rangle + \langle \|\vec{v}\|\vec{w} + \|\vec{w}\|\vec{v}, -\|\vec{w}\|\vec{v} \rangle \text{ by P2} \\
 = & \langle \|\vec{v}\|\vec{w}, \|\vec{v}\|\vec{w} + \|\vec{w}\|\vec{v} \rangle + \langle -\|\vec{w}\|\vec{v}, \|\vec{v}\|\vec{w} + \|\vec{w}\|\vec{v} \rangle \text{ by P1} \\
 = & \langle \|\vec{v}\|\vec{w}, \|\vec{v}\|\vec{w} \rangle + \langle \|\vec{v}\|\vec{w}, \|\vec{w}\|\vec{v} \rangle + \langle -\|\vec{w}\|\vec{v}, \|\vec{v}\|\vec{w} \rangle \\
 & + \langle -\|\vec{w}\|\vec{v}, \|\vec{w}\|\vec{v} \rangle \text{ by P2} \\
 = & \|\vec{v}\|^2 \langle \vec{w}, \vec{w} \rangle + \|\vec{v}\|\|\vec{w}\| \langle \vec{w}, \vec{v} \rangle - \|\vec{v}\|\|\vec{w}\| \langle \vec{v}, \vec{w} \rangle - \|\vec{w}\|^2 \langle \vec{v}, \vec{v} \rangle \text{ by P3}
 \end{aligned}$$

## Page 237 Number 26

**Page 237 Number 26.** Let  $\vec{v}$  and  $\vec{w}$  be vectors in an inner-product space  $V$ . Show that  $\|\vec{v}\|\vec{w} + \|\vec{w}\|\vec{v}$  is perpendicular to  $\|\vec{v}\|\vec{w} - \|\vec{w}\|\vec{v}$ .

**Solution.** Consider

$$\begin{aligned}
 & \langle \|\vec{v}\|\vec{w} + \|\vec{w}\|\vec{v}, \|\vec{v}\|\vec{w} - \|\vec{w}\|\vec{v} \rangle \\
 = & \langle \|\vec{v}\|\vec{w} + \|\vec{w}\|\vec{v}, \|\vec{v}\|\vec{w} \rangle + \langle \|\vec{v}\|\vec{w} + \|\vec{w}\|\vec{v}, -\|\vec{w}\|\vec{v} \rangle \text{ by P2} \\
 = & \langle \|\vec{v}\|\vec{w}, \|\vec{v}\|\vec{w} + \|\vec{w}\|\vec{v} \rangle + \langle -\|\vec{w}\|\vec{v}, \|\vec{v}\|\vec{w} + \|\vec{w}\|\vec{v} \rangle \text{ by P1} \\
 = & \langle \|\vec{v}\|\vec{w}, \|\vec{v}\|\vec{w} \rangle + \langle \|\vec{v}\|\vec{w}, \|\vec{w}\|\vec{v} \rangle + \langle -\|\vec{w}\|\vec{v}, \|\vec{v}\|\vec{w} \rangle \\
 & + \langle -\|\vec{w}\|\vec{v}, \|\vec{w}\|\vec{v} \rangle \text{ by P2} \\
 = & \|\vec{v}\|^2 \langle \vec{w}, \vec{w} \rangle + \|\vec{v}\|\|\vec{w}\| \langle \vec{w}, \vec{v} \rangle - \|\vec{v}\|\|\vec{w}\| \langle \vec{v}, \vec{w} \rangle - \|\vec{w}\|^2 \langle \vec{v}, \vec{v} \rangle \text{ by P3} \\
 = & \|\vec{v}\|^2 \|\vec{w}\|^2 - \|\vec{w}\|^2 \|\vec{v}\|^2 \text{ by Definition 3.13,} \\
 & \text{“Magnitude and Norm of a Vector”} \\
 = & 0.
 \end{aligned}$$

Since the inner product is 0, the vectors are perpendicular.  $\square$

## Page 237 Number 26

**Page 237 Number 26.** Let  $\vec{v}$  and  $\vec{w}$  be vectors in an inner-product space  $V$ . Show that  $\|\vec{v}\|\vec{w} + \|\vec{w}\|\vec{v}$  is perpendicular to  $\|\vec{v}\|\vec{w} - \|\vec{w}\|\vec{v}$ .

**Solution.** Consider

$$\begin{aligned}
 & \langle \|\vec{v}\|\vec{w} + \|\vec{w}\|\vec{v}, \|\vec{v}\|\vec{w} - \|\vec{w}\|\vec{v} \rangle \\
 = & \langle \|\vec{v}\|\vec{w} + \|\vec{w}\|\vec{v}, \|\vec{v}\|\vec{w} \rangle + \langle \|\vec{v}\|\vec{w} + \|\vec{w}\|\vec{v}, -\|\vec{w}\|\vec{v} \rangle \text{ by P2} \\
 = & \langle \|\vec{v}\|\vec{w}, \|\vec{v}\|\vec{w} + \|\vec{w}\|\vec{v} \rangle + \langle -\|\vec{w}\|\vec{v}, \|\vec{v}\|\vec{w} + \|\vec{w}\|\vec{v} \rangle \text{ by P1} \\
 = & \langle \|\vec{v}\|\vec{w}, \|\vec{v}\|\vec{w} \rangle + \langle \|\vec{v}\|\vec{w}, \|\vec{w}\|\vec{v} \rangle + \langle -\|\vec{w}\|\vec{v}, \|\vec{v}\|\vec{w} \rangle \\
 & + \langle -\|\vec{w}\|\vec{v}, \|\vec{w}\|\vec{v} \rangle \text{ by P2} \\
 = & \|\vec{v}\|^2 \langle \vec{w}, \vec{w} \rangle + \|\vec{v}\| \|\vec{w}\| \langle \vec{w}, \vec{v} \rangle - \|\vec{v}\| \|\vec{w}\| \langle \vec{v}, \vec{w} \rangle - \|\vec{w}\|^2 \langle \vec{v}, \vec{v} \rangle \text{ by P3} \\
 = & \|\vec{v}\|^2 \|\vec{w}\|^2 - \|\vec{w}\|^2 \|\vec{v}\|^2 \text{ by Definition 3.13,} \\
 & \text{“Magnitude and Norm of a Vector”} \\
 = & 0.
 \end{aligned}$$

Since the inner product is 0, the vectors are perpendicular.  $\square$