


Theory of Matrices

Chapter 5. Matrix Transformations and Factorizations 5.7. QR Factorization—Proofs of Theorems



Theorem 5.7.A

Theorem 5.7.A. Let $\{a_1, a_2, \dots, a_k\}$ be a linearly independent set of vectors. There exists unique scalars x_{ij} where $1 \leq j \leq k$ and $0 < i < j$ such that the k vectors

$$\begin{aligned} b_1 &= a_1 \\ b_2 &= a_2 - x_{12}b_1 \\ &\vdots \\ b_j &= a_j - x_{j-1,j}b_{j-1} - x_{j-2,j}b_{j-2} - \dots - x_{1j}b_1 \\ &\vdots \\ b_m &= a_m - x_{m-1,m}b_{m-1} - x_{m-2,m}b_{m-2} - \dots - x_{1m}b_1 \end{aligned}$$

form an orthogonal set. The vectors b_1, b_2, \dots, b_k are nonzero and $x_{ij} = \langle a_j, b_i \rangle / \langle b_i, b_i \rangle$ for $1 \leq j \leq m$ and $0 < i < m$.

Theorem 5.7.A (continued 1)

Proof (continued). We prove this by induction on k . The result is trivial for $k = 1$ since we have $b_1 = a_1$ and there are no x_{ij} . Suppose the result holds for $m = k - 1$. That is, suppose there exists unique x_{ij} for $1 \leq j \leq k - 1$ and $0 < i < j$ such that b_1, b_2, \dots, b_{k-1} defined as given in the statement of the theorem form an orthogonal set of nonzero vectors where $x_{ij} = \langle a_j, b_i \rangle / \langle b_i, b_i \rangle$. We define b_k as given in the statement of the theorem and we must show that b_k is nonzero and that the coefficients x_{ij} are as claimed for $j = k$ and $0 < i < j$. Notice that for $0 < i < j$ we have

$$\begin{aligned} \langle b_k, b_i \rangle &= \langle a_k - x_{k-1,k}b_{k-1} - x_{k-2,k}b_{k-2} - \dots - x_{1k}b_1, b_i \rangle \\ &= \langle a_k, b_i \rangle - x_{k-1,k} \langle b_{k-1}, b_i \rangle - \dots - x_{i,k} \langle b_i, b_i \rangle - \dots - x_{1k} \langle b_1, b_i \rangle \\ &= \langle a_k, b_i \rangle - x_{ik} \langle b_i, b_i \rangle. \end{aligned}$$

So $\langle b_k, b_i \rangle = 0$ if and only if $\langle a_k, b_i \rangle - x_{ik} \langle b_i, b_i \rangle = 0$ or, since $b_i \neq 0$ by the induction hypothesis, $\langle b_k, b_i \rangle = 0$ (that is, $b_1, b_2, \dots, b_{k-1}, b_k$ form an orthogonal set) if and only if $x_{ik} = \langle a_k, b_i \rangle / \langle b_i, b_i \rangle$.

Theorem 5.7.A (continued 2)

Proof (continued). So the scalars x_{ij} for $j = k$ and $0 < i < j$ must satisfy $x_{ik} = \langle a_k, b_i \rangle / \langle b_i, b_i \rangle$. So by induction, the x_{ij} are as claimed for $1 \leq j \leq k$ and $0 < i < j$.

We just need to show that b_k is nonzero. From the definition of b_1, b_2, \dots, b_k we can express:

- b_1 in terms of a_1 with a coefficient of 1 for a_1 ,
- b_2 in terms of a_2 and a_1 with a coefficient of 1 for a_2 ,
- b_3 in terms of a_3, a_2 , and a_1 with a coefficient of 1 for a_3, \dots , and
- b_k in terms of a_k, a_{k-1}, \dots, a_1 with a coefficient of 1 for a_k .

Since $\{a_1, a_2, \dots, a_k\}$ is a linearly independent set and when expressing b_k as a linear combination of these vectors the coefficient of a_k is nonzero then b_k is nonzero. So the result holds for $m = k$ and therefore holds in general by induction. □

Theorem 5.7.B

Theorem 5.7.B. Let A be an $n \times m$ matrix of full column rank (that is, $\text{rank}(A) = m$). Then there is $n \times m$ matrix B , where the columns of B are mutually orthogonal and nonzero, and $m \times m$ upper triangular matrix X , with all diagonal entries 1, such that $A = BX$.

Proof. Denote the (linearly independent) columns of A as a_1, a_2, \dots, a_m . Recursively define (column) vectors b_1, b_2, \dots, b_m as

$$\begin{aligned} b_1 &= a_1 \\ b_2 &= a_2 - x_{12}b_1 \\ b_3 &= a_3 - x_{23}b_2 - x_{13}b_1 \\ &\vdots \\ b_j &= a_j - x_{j-1,j}b_{j-1} - x_{j-2,j}b_{j-2} - \dots - x_{1j}b_1 \\ &\vdots \\ b_m &= a_m - x_{m-1,m}b_{m-1} - x_{m-2,m}b_{m-2} - \dots - x_{1m}b_1 \end{aligned}$$

()

Theorem 5.7.B (continued 1)

Proof (continued). ... where $x_{ij} = \langle a_j, b_i \rangle / \langle b_i, b_i \rangle$. The b_1, b_2, \dots, b_m are mutually orthogonal and each is nonzero by Theorem 5.7.A. We can rearrange the equations defining the b_i 's to express the a_j 's as linear combinations of the b_i 's:

$$\begin{aligned} a_1 &= b_1 \\ a_2 &= b_2 + x_{12}b_1 \\ &\vdots \\ a_j &= b_j + x_{j-1,j}b_{j-1} + x_{j-2,j}b_{j-2} + \dots + x_{1j}b_1 \\ &\vdots \\ a_m &= b_m + x_{m-1,m}b_{m-1} + x_{m-2,m}b_{m-2} + \dots + x_{1m}b_1 \quad (*) \end{aligned}$$

where $x_{ij} = \langle a_j, b_i \rangle / \langle b_i, b_i \rangle$.

()

Theorem 5.7.B (continued 2)

Theorem 5.7.B. Let A be an $n \times m$ matrix of full column rank (that is, $\text{rank}(A) = m$). Then there is $n \times m$ matrix B , where the columns of B are mutually orthogonal and nonzero, and $m \times m$ upper triangular matrix X , with all diagonal entries 1, such that $A = BX$.

Proof (continued). Define $n \times m$ matrix B with columns b_1, b_2, \dots, b_m and $m \times m$ matrix X with entry (i, j) as x_{ij} for $i < j$, $x_{ii} = 1$, and $x_{ij} = 0$ for $i > j$. So the columns of B are orthogonal and X is upper triangular with diagonal entries of 1. From (*), we see that $A = BX$, as claimed. \square

()