


# Real Analysis

## Chapter 10. Metric Spaces: Three Fundamental Theorem

### 10.3. The Banach Contraction Principle—Proofs of Theorems



## Theorem 10.3.B. The Banach Contraction Principle

### Theorem 10.3.B. The Banach Contraction Principle.

Let  $X$  be a complete metric space and the mapping  $T : X \rightarrow X$  be a contract. Then  $T : X \rightarrow X$  has exactly one fixed point.

**Proof.** Let  $c$  be a number with  $0 \leq c < 1$  that is a Lipschitz constant for mapping  $T$ . Select any point in  $X$  and label it  $x_0$ . Define the sequence  $\{x_k\}$  inductively by defining  $x_1 = T(x_0)$  and, if  $k \in \mathbb{N}$  is such that  $x_k$  is defined, then define  $x_{k+1} = T(x_k)$ . The sequence  $\{x_n\}$  is properly defined since  $T(x)$  is a subset of  $X$ . We will show that  $\{x_n\}$  converges to a fixed point of  $T$ .

Since  $T$  is a contraction with Lipschitz constant  $c$ , we have

$$\rho(x_2, x_1) = \rho(T(x_1), T(x_0)) = \rho(T(T(x_0)), T(x_0)) \leq c \rho(T(x_0), x_0),$$

and similarly

$$\rho(x_{k+1}, x_k) = \rho(T(x_k), T(x_{k-1})) \leq c \rho(x_k, x_{k-1}) \text{ if } k \geq 2.$$

## Theorem 10.3.B (continued 1)

**Proof (continued).** These two inequalities imply, by induction, that

$$\rho(x_{k+1}, x_k) \leq c^k \rho(T(x_0), x_0) \text{ for every } k \in \mathbb{N}.$$

Hence if  $m, k \in \mathbb{N}$  with  $m > k$ , then

$$\begin{aligned} \rho(x_m, x_k) &\leq \rho(x_m, x_{m-1}) + \rho(x_{m-1}, x_{m-2}) + \cdots + \rho(x_{k+1}, x_k) \\ &\quad \text{by the Triangle Inequality for } \rho \\ &\leq (c^{m-1} + c^{m-2} + \cdots + c^k) \rho(T(x_0), x_0) \\ &= c^k (1 + c + \cdots + c^{m-1-k}) \rho(T(x_0), x_0) \\ &= c^k \frac{1 - c^{m-k}}{1 - c} \rho(T(x_0), x_0) \text{ since } \sum_{k=0}^n c^k = \frac{1 - c^{n+1}}{1 - c} \text{ for } c \neq 1. \end{aligned}$$

Since  $0 \leq c < 1$ , then  $\rho(x_m, x_k) \leq \frac{c^k}{1 - c} \rho(T(x_0), x_0)$  if  $m > k$ .

## Theorem 10.3.B (continued 2)

**Proof (continued).** Since  $0 \leq c < 1$ , then  $\rho(x_m, x_k) \leq \frac{c^k}{1 - c} \rho(T(x_0), x_0)$  if  $m > k$ . But  $\lim_{k \rightarrow \infty} c^k = 0$ , so that the inequality implies that  $\{x_k\}$  is a Cauchy sequence.

By hypothesis, the metric space  $S$  is complete. Thus there is a point  $x \in X$  to which the sequence  $\{x_k\}$  converges. Since  $T$  is Lipschitz, it is continuous (let  $\delta = \varepsilon/c$  in the  $\varepsilon/\delta$  Criterion for Continuity, Theorem 9.3.A). Therefore

$T(x) = \lim_{k \rightarrow \infty} T(x_k) = T(\lim_{k \rightarrow \infty} x_k) = \lim_{k \rightarrow \infty} x_{k+1} = x$ . Thus the mapping  $T : X \rightarrow X$  has at least one fixed point. ASSUME that  $u$  and  $v$  are two different fixed points such that  $T(u) = u$  and  $T(v) = v$ . Then  $0 < \rho(u, v) = \rho(T(u), T(v)) \leq c \rho(u, v)$ , so that since  $0 \leq c < 1$ , we must have  $\rho(u, v) = 0$ . But then  $u = v$ , a CONTRADICTION. So the assumption that there are two different fixed points is false, and hence  $T$  has exactly one fixed point, as claimed.  $\square$

## Theorem 10.3.C. The Picard Local Existence Theorem

**Theorem 10.3.C. The Picard Local Existence Theorem.**

Let  $\mathcal{O}$  be an open subset of the plane  $\mathbb{R}^2$  containing the point  $(x_0, y_0)$ . Suppose the function  $g : \mathcal{O} \rightarrow \mathbb{R}^2$  is continuous and there is a positive number  $M$  for which the following Lipschitz property in the second variable holds uniformly with respect to the first variable:

$$|g(x, y_1) - g(x, y_2)| \leq M|y_1 - y_2| \text{ for all points } (x, y_1) \text{ and } (x, y_2) \text{ in } \mathcal{O}. \quad (16)$$

Then there is an open interval  $I$  containing  $x_0$  on which the following differential equation has a unique solution:

$$\begin{aligned} f'(x) &= g(x, f(x)) \text{ for all } x \in I \\ f(x_0) &= y_0. \end{aligned} \quad (14)$$

**Proof.** For  $\ell$  a positive number, define  $I_\ell$  to be the closed interval  $[x_0 - \ell, x_0 + \ell]$ . By Note 10.3.C, we have a solution of differential equation (14) if and only if we have a solution of integral equation (15).

()

## Theorem 10.3.C (continued 2)

**Proof (continued).** A solution of the integral equation (15) is a fixed point of the mapping  $T : X_\ell \rightarrow C(I_\ell)$ . Since  $C(I_\ell)$  is a complete metric space by Proposition 9.10 (where the norm on  $C(I_\ell)$  has the max norm) and  $X_\ell$  is a closed subset of  $C(I_\ell)$  by Proposition 9.11, then  $X_\ell$  is a complete metric space by Proposition 9.12(iii). We will show that if  $\ell$  is chosen sufficiently small, then  $T(X_\ell) \subseteq X_\ell$  and  $T : X_\ell \rightarrow X_\ell$  is a contract. The Banach Contraction Principle will then imply that  $T : X_\ell \rightarrow X_\ell$  has a unique fixed point, and hence integral equation (15) and differential equation (14) have unique solutions.

Since rectangle  $R = [x_0 - a, x_0 + a] \times [y_0 - b, y_0 + b]$  is compact (by the Heine-Borel Theorem) and  $g$  is continuous by hypothesis, then  $g$  is the Extreme Value Theorem (Proposition 9.22) there is a positive number  $K$  such that  $|g(x, y)| \leq K$  for all  $(x, y) \in R$ .

()

## Theorem 10.3.C (continued 1)

**Proof (continued).** So, it suffices to show that  $\ell$  can be chosen so that there is exactly one continuous function  $f : I_\ell \rightarrow \mathbb{R}$  with the property that

$$f(x) = y_0 + \int_{x_0}^x g(t, f(t)) dt \text{ for all } x \in I_\ell.$$

Since  $\mathcal{O}$  is open, we may choose positive numbers  $a$  and  $b$  such that the closed rectangle  $R = [x_0 - a, x_0 + a] \times [y_0 - b, y_0 + b]$  is contained in  $\mathcal{O}$ . Now for each positive number  $\ell$  with  $\ell \leq a$ , define  $X_\ell$  to be the subspace of the metric space  $C(I_\ell)$  consisting of those continuous functions  $f : I_\ell \rightarrow \mathbb{R}$  such that  $|f(x) - y_0| \leq b$  for all  $x \in I_\ell$ . That is,  $X_\ell$  consists of all continuous functions on  $I_\ell$  (since  $X_\ell$  is a subspace of  $C(I_\ell)$ ) that have a graph contained in the rectangle  $I_\ell \times [y_0 - b, y_0 + b]$ .

For  $f \in X_\ell$ , define the function  $T(f) \in C(I_\ell)$  by

$$T(f)(x) = y_0 + \int_{x_0}^x g(t, f(t)) dt \text{ for all } x \in I_\ell.$$

()

## Theorem 10.3.C (continued 3)

**Proof (continued).** For  $f \in X_\ell$  and  $x \in I_\ell$  we have, by the definition of  $T$ ,

$$|T(f)(x) - y_0| = \left| \int_{x_0}^x g(t, f(t)) dt \right| \leq \ell K,$$

so that  $T(X_\ell) \subseteq X_\ell$  if  $\ell K \leq b$ . Observe that for functions  $f_1, f_2 \in X_\ell$  and  $x \in I_\ell$ , by hypothesized inequality (16) we have  $|g(x, f_1(x)) - g(x, f_2(x))| \leq M \rho_{\max}(f_1, f_2)$  since  $|y_1 - y_2| = |f(x_1) - f(x_2)| \leq \rho_{\max}(f_1, f_2)$ . So by the linearity and monotonicity of integrals, we have

$$\begin{aligned} |T(f_1)(x) - T(f_2)(x)| &= \left| \int_{x_0}^x [g(t, f_1(t)) - g(t, f_2(t))] dt \right| \\ &\leq |x - x_0| M \rho_{\max}(f_1, f_2) \text{ by the previous inequality} \\ &\leq \ell M \rho_{\max}(f_1, f_2) \text{ since } x \in I_\ell = [x_0 - \ell, x_0 + \ell]. \end{aligned}$$

This inequality, together with the inclusion  $T(X_\ell) \subseteq X_\ell$  if  $\ell K \leq b$ , implies that  $T : X_\ell \rightarrow X_\ell$  is a contraction if  $\ell K \leq b$  and  $\ell M < 1$ .

()

## Theorem 10.3.C (continued 4)

**Theorem 10.3.C. The Picard Local Existence Theorem.**

Let  $\mathcal{O}$  be an open subset of the plane  $\mathbb{R}^2$  containing the point  $(x_0, y_0)$ . Suppose the function  $g : \mathcal{O} \rightarrow \mathbb{R}^2$  is continuous and there is a positive number  $M$  for which the following Lipschitz property in the second variable holds uniformly with respect to the first variable:

$$|g(x, y_1) - g(x, y_2)| \leq M|y_1 - y_2| \text{ for all points } (x, y_1) \text{ and } (x, y_2) \text{ in } \mathcal{O}. \quad (16)$$

Then there is an open interval  $I$  containing  $x_0$  on which the following differential equation has a unique solution:

$$\begin{aligned} f'(x) &= g(x, f(x)) \text{ for all } x \in I \\ f(x_0) &= y_0. \end{aligned} \quad (14)$$

**Proof (continued).** So we define  $\ell = \min\{b/K, 1/(2M)\}$ . Then the Banach Contraction Principle applied to  $T : X_\ell \rightarrow X_\ell$  implies that  $T$  has a unique fixed point and, as described above, the integral equation (15) and the differential equation (14) each have a unique solution, as claimed.  $\square$