


# Real Analysis

## Chapter 11. Topological Spaces: General Properties

### 11.5. Compact Topological Spaces—Proofs of Theorems



# Table of contents

- 1 Proposition 11.14
- 2 Proposition 11.15
- 3 Proposition 11.16
- 4 Proposition 11.17
- 5 Theorem 11.18
- 6 Proposition 11.20
- 7 Corollary 11.21
- 8 Proposition 11.19

## Proposition 11.14

**Proposition 11.14.** A topological space  $(X, \mathcal{T})$  is compact if and only if every collection of closed subsets of  $X$  that possesses the finite intersection property has nonempty intersection.

**Proof.** Let  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  be an open cover of  $X$ . Define  $C_\lambda = X \setminus \mathcal{O}_\lambda$ . Then each  $C_\lambda$  is closed. Also, by DeMorgan's Laws,

$$X = \bigcup_{\lambda \in \Lambda} \mathcal{O}_\lambda \text{ implies } \emptyset = \bigcap_{\lambda \in \Lambda} C_\lambda.$$

## Proposition 11.14

**Proposition 11.14.** A topological space  $(X, \mathcal{T})$  is compact if and only if every collection of closed subsets of  $X$  that possesses the finite intersection property has nonempty intersection.

**Proof.** Let  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  be an open cover of  $X$ . Define  $C_\lambda = X \sim \mathcal{O}_\lambda$ . Then each  $C_\lambda$  is closed. Also, by DeMorgan's Laws,

$$X = \bigcup_{\lambda \in \Lambda} \mathcal{O}_\lambda \text{ implies } \emptyset = \bigcap_{\lambda \in \Lambda} C_\lambda.$$

Conversely, if  $\{C_\lambda\}_{\lambda \in \Lambda}$  is a collection of closed sets and we define  $\mathcal{O}_\lambda = X \sim C_\lambda$ , then by DeMorgan's Laws,

$$\emptyset = \bigcap_{\lambda \in \Lambda} C_\lambda \text{ implies } X = \bigcup_{\lambda \in \Lambda} \mathcal{O}_\lambda.$$

## Proposition 11.14

**Proposition 11.14.** A topological space  $(X, \mathcal{T})$  is compact if and only if every collection of closed subsets of  $X$  that possesses the finite intersection property has nonempty intersection.

**Proof.** Let  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  be an open cover of  $X$ . Define  $C_\lambda = X \sim \mathcal{O}_\lambda$ . Then each  $C_\lambda$  is closed. Also, by DeMorgan's Laws,

$$X = \bigcup_{\lambda \in \Lambda} \mathcal{O}_\lambda \text{ implies } \emptyset = \bigcap_{\lambda \in \Lambda} C_\lambda.$$

Conversely, if  $\{C_\lambda\}_{\lambda \in \Lambda}$  is a collection of closed sets and we define  $\mathcal{O}_\lambda = X \sim C_\lambda$ , then by DeMorgan's Laws,

$$\emptyset = \bigcap_{\lambda \in \Lambda} C_\lambda \text{ implies } X = \bigcup_{\lambda \in \Lambda} \mathcal{O}_\lambda.$$

Suppose  $X$  is compact and let  $\{C_\lambda\}_{\lambda \in \Lambda}$  be a collection of closed sets with the finite intersection property. ASSUME  $\bigcap_{\lambda \in \Lambda} C_\lambda = \emptyset$ . Define  $\mathcal{O}_\lambda$  as above.

# Proposition 11.14

**Proposition 11.14.** A topological space  $(X, \mathcal{T})$  is compact if and only if every collection of closed subsets of  $X$  that possesses the finite intersection property has nonempty intersection.

**Proof.** Let  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  be an open cover of  $X$ . Define  $C_\lambda = X \sim \mathcal{O}_\lambda$ . Then each  $C_\lambda$  is closed. Also, by DeMorgan's Laws,

$$X = \bigcup_{\lambda \in \Lambda} \mathcal{O}_\lambda \text{ implies } \emptyset = \bigcap_{\lambda \in \Lambda} C_\lambda.$$

Conversely, if  $\{C_\lambda\}_{\lambda \in \Lambda}$  is a collection of closed sets and we define  $\mathcal{O}_\lambda = X \sim C_\lambda$ , then by DeMorgan's Laws,

$$\emptyset = \bigcap_{\lambda \in \Lambda} C_\lambda \text{ implies } X = \bigcup_{\lambda \in \Lambda} \mathcal{O}_\lambda.$$

Suppose  $X$  is compact and let  $\{C_\lambda\}_{\lambda \in \Lambda}$  be a collection of closed sets with the finite intersection property. ASSUME  $\bigcap_{\lambda \in \Lambda} C_\lambda = \emptyset$ . Define  $\mathcal{O}_\lambda$  as above.

## Proposition 11.14 (continued)

**Proof (continued).** Then  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  is an open cover of  $X$ . Since  $X$  is compact, then for some  $\lambda = 1, 2, \dots, n$  we have  $X \subseteq \bigcup_{i=1}^n \mathcal{O}_i$ . So  $\bigcap_{i=1}^n C_i = \emptyset$ , but this contradicts the fact that  $\{C_\lambda\}_{\lambda \in \Lambda}$  has the finite intersection property. So the assumption that  $\bigcap_{\lambda \in \Lambda} C_\lambda = \emptyset$  is false and it must be that  $\bigcap_{\lambda \in \Lambda} C_\lambda \neq \emptyset$ . That is, every collection of closed sets with the finite intersection property has nonempty intersection.

## Proposition 11.14 (continued)

**Proof (continued).** Then  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  is an open cover of  $X$ . Since  $X$  is compact, then for some  $\lambda = 1, 2, \dots, n$  we have  $X \subseteq \cup_{i=1}^n \mathcal{O}_i$ . So  $\cap_{i=1}^n C_i = \emptyset$ , but this contradicts the fact that  $\{C_\lambda\}_{\lambda \in \Lambda}$  has the finite intersection property. So the assumption that  $\cap_{\lambda \in \Lambda} C_\lambda = \emptyset$  is false and it must be that  $\cap_{\lambda \in \Lambda} C_\lambda \neq \emptyset$ . That is, every collection of closed sets with the finite intersection property has nonempty intersection.

Suppose every collection of closed sets with the finite intersection property has nonempty intersection. Let  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  be an open cover of  $X$ .



## Proposition 11.14 (continued)

**Proof (continued).** Then  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  is an open cover of  $X$ . Since  $X$  is compact, then for some  $\lambda = 1, 2, \dots, n$  we have  $X \subseteq \cup_{i=1}^n \mathcal{O}_i$ . So  $\cap_{i=1}^n C_i = \emptyset$ , but this contradicts the fact that  $\{C_\lambda\}_{\lambda \in \Lambda}$  has the finite intersection property. So the assumption that  $\cap_{\lambda \in \Lambda} C_\lambda = \emptyset$  is false and it must be that  $\cap_{\lambda \in \Lambda} C_\lambda \neq \emptyset$ . That is, every collection of closed sets with the finite intersection property has nonempty intersection.

Suppose every collection of closed sets with the finite intersection property has nonempty intersection. Let  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  be an open cover of  $X$ . ASSUME  $X$  is not compact. Then no finite subcollection of  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  is a cover of  $X$ . Define  $C_\lambda = X \setminus \mathcal{O}_\lambda$ . Then any finite subcollection of  $\{C_\lambda\}_{\lambda \in \Lambda}$  is nonempty (and, of course, each  $C_\lambda$  is closed).

# Proposition 11.14 (continued)

**Proof (continued).** Then  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  is an open cover of  $X$ . Since  $X$  is compact, then for some  $\lambda = 1, 2, \dots, n$  we have  $X \subseteq \cup_{i=1}^n \mathcal{O}_i$ . So  $\cap_{i=1}^n C_i = \emptyset$ , but this contradicts the fact that  $\{C_\lambda\}_{\lambda \in \Lambda}$  has the finite intersection property. So the assumption that  $\cap_{\lambda \in \Lambda} C_\lambda = \emptyset$  is false and it must be that  $\cap_{\lambda \in \Lambda} C_\lambda \neq \emptyset$ . That is, every collection of closed sets with the finite intersection property has nonempty intersection.

Suppose every collection of closed sets with the finite intersection property has nonempty intersection. Let  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  be an open cover of  $X$ . ASSUME  $X$  is not compact. Then no finite subcollection of  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  is a cover of  $X$ . Define  $C_\lambda = X \setminus \mathcal{O}_\lambda$ . Then any finite subcollection of  $\{C_\lambda\}_{\lambda \in \Lambda}$  is nonempty (and, of course, each  $C_\lambda$  is closed). So, by hypotheses,  $\cap_{\lambda \in \Lambda} C_\lambda \neq \emptyset$ . But then  $\cup_{\lambda \in \Lambda} \mathcal{O}_\lambda \neq X$  and  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  is not a cover of  $X$ , a CONTRADICTION.

# Proposition 11.14 (continued)

**Proof (continued).** Then  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  is an open cover of  $X$ . Since  $X$  is compact, then for some  $\lambda = 1, 2, \dots, n$  we have  $X \subseteq \cup_{i=1}^n \mathcal{O}_i$ . So  $\cap_{i=1}^n C_i = \emptyset$ , but this contradicts the fact that  $\{C_\lambda\}_{\lambda \in \Lambda}$  has the finite intersection property. So the assumption that  $\cap_{\lambda \in \Lambda} C_\lambda = \emptyset$  is false and it must be that  $\cap_{\lambda \in \Lambda} C_\lambda \neq \emptyset$ . That is, every collection of closed sets with the finite intersection property has nonempty intersection.

Suppose every collection of closed sets with the finite intersection property has nonempty intersection. Let  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  be an open cover of  $X$ . ASSUME  $X$  is not compact. Then no finite subcollection of  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  is a cover of  $X$ . Define  $C_\lambda = X \setminus \mathcal{O}_\lambda$ . Then any finite subcollection of  $\{C_\lambda\}_{\lambda \in \Lambda}$  is nonempty (and, of course, each  $C_\lambda$  is closed). So, by hypotheses,  $\cap_{\lambda \in \Lambda} C_\lambda \neq \emptyset$ . But then  $\cup_{\lambda \in \Lambda} \mathcal{O}_\lambda \neq X$  and  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  is not a cover of  $X$ , a CONTRADICTION. So the assumption that  $X$  is not compact is false.  $\square$

# Proposition 11.14 (continued)

**Proof (continued).** Then  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  is an open cover of  $X$ . Since  $X$  is compact, then for some  $\lambda = 1, 2, \dots, n$  we have  $X \subseteq \cup_{i=1}^n \mathcal{O}_i$ . So  $\cap_{i=1}^n C_i = \emptyset$ , but this contradicts the fact that  $\{C_\lambda\}_{\lambda \in \Lambda}$  has the finite intersection property. So the assumption that  $\cap_{\lambda \in \Lambda} C_\lambda = \emptyset$  is false and it must be that  $\cap_{\lambda \in \Lambda} C_\lambda \neq \emptyset$ . That is, every collection of closed sets with the finite intersection property has nonempty intersection.

Suppose every collection of closed sets with the finite intersection property has nonempty intersection. Let  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  be an open cover of  $X$ . ASSUME  $X$  is not compact. Then no finite subcollection of  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  is a cover of  $X$ . Define  $C_\lambda = X \setminus \mathcal{O}_\lambda$ . Then any finite subcollection of  $\{C_\lambda\}_{\lambda \in \Lambda}$  is nonempty (and, of course, each  $C_\lambda$  is closed). So, by hypotheses,  $\cap_{\lambda \in \Lambda} C_\lambda \neq \emptyset$ . But then  $\cup_{\lambda \in \Lambda} \mathcal{O}_\lambda \neq X$  and  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  is not a cover of  $X$ , a CONTRADICTION. So the assumption that  $X$  is not compact is false.  $\square$

# Proposition 11.15

**Proposition 11.15.** A closed subset of a compact topological space  $(X, \mathcal{T})$  is compact.

**Proof.** Let  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  be an open cover of  $K$  with open sets in  $\mathcal{T}$ . Since  $X \sim K$  is open in  $\mathcal{T}$ , then  $\{X \sim K\} \cup \{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  is an open cover of  $X$ .

# Proposition 11.15

**Proposition 11.15.** A closed subset of a compact topological space  $(X, \mathcal{T})$  is compact.

**Proof.** Let  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  be an open cover of  $K$  with open sets in  $\mathcal{T}$ . Since  $X \sim K$  is open in  $\mathcal{T}$ , then  $\{X \sim K\} \cup \{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  is an open cover of  $X$ . Since  $X$  is compact, there is a finite subcover of  $X$ . This subcover of  $X$  is also a cover of  $K$ .

# Proposition 11.15

**Proposition 11.15.** A closed subset of a compact topological space  $(X, \mathcal{T})$  is compact.

**Proof.** Let  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  be an open cover of  $K$  with open sets in  $\mathcal{T}$ . Since  $X \sim K$  is open in  $\mathcal{T}$ , then  $\{X \sim K\} \cup \{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  is an open cover of  $X$ . Since  $X$  is compact, there is a finite subcover of  $X$ . This subcover of  $X$  is also a cover of  $K$ . If  $X \sim K$  is included in the subcover of  $K$ , then it can be omitted from the subcover of  $K$ , since it is disjoint from  $K$ . The resulting subcover of  $K$  is a subset of the original cover of  $K$  and hence  $K$  is compact.  $\square$

# Proposition 11.15

**Proposition 11.15.** A closed subset of a compact topological space  $(X, \mathcal{T})$  is compact.

**Proof.** Let  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  be an open cover of  $K$  with open sets in  $\mathcal{T}$ . Since  $X \sim K$  is open in  $\mathcal{T}$ , then  $\{X \sim K\} \cup \{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  is an open cover of  $X$ . Since  $X$  is compact, there is a finite subcover of  $X$ . This subcover of  $X$  is also a cover of  $K$ . If  $X \sim K$  is included in the subcover of  $K$ , then it can be omitted from the subcover of  $K$ , since it is disjoint from  $K$ . The resulting subcover of  $K$  is a subset of the original cover of  $K$  and hence  $K$  is compact.  $\square$



# Proposition 11.16

**Proposition 11.16.** A compact subspace  $K$  of a Hausdorff topological space  $(X, \mathcal{T})$  is a closed subset of  $X$ .

**Proof.** If  $K = X$ , then  $K$  is closed. Otherwise, let  $y \in X \setminus K$ . Since  $X$  is Hausdorff, for each  $x \in K$  there are disjoint neighborhoods  $\mathcal{O}_x$  and  $\mathcal{U}_x$  of  $x$  and  $y$  respectively. Then  $\{\mathcal{O}_x\}_{x \in K}$  is an open cover of  $K$ .

# Proposition 11.16

**Proposition 11.16.** A compact subspace  $K$  of a Hausdorff topological space  $(X, \mathcal{T})$  is a closed subset of  $X$ .

**Proof.** If  $K = X$ , then  $K$  is closed. Otherwise, let  $y \in X \setminus K$ . Since  $X$  is Hausdorff, for each  $x \in K$  there are disjoint neighborhoods  $\mathcal{O}_x$  and  $\mathcal{U}_x$  of  $x$  and  $y$  respectively. Then  $\{\mathcal{O}_x\}_{x \in K}$  is an open cover of  $K$ . Since  $K$  is compact, there is a finite subcover  $\{\mathcal{O}_{x_1}, \mathcal{O}_{x_2}, \dots, \mathcal{O}_{x_n}\}$ . Define  $\mathcal{N} = \bigcap_{i=1}^n \mathcal{U}_{x_i}$ . Then  $\mathcal{N}$  is a neighborhood of  $y$  which is disjoint from each  $\mathcal{O}_{x_i}$  (since  $\mathcal{O}_{x_i}$  and  $\mathcal{U}_{x_i}$  are disjoint).

# Proposition 11.16

**Proposition 11.16.** A compact subspace  $K$  of a Hausdorff topological space  $(X, \mathcal{T})$  is a closed subset of  $X$ .

**Proof.** If  $K = X$ , then  $K$  is closed. Otherwise, let  $y \in X \setminus K$ . Since  $X$  is Hausdorff, for each  $x \in K$  there are disjoint neighborhoods  $\mathcal{O}_x$  and  $\mathcal{U}_x$  of  $x$  and  $y$  respectively. Then  $\{\mathcal{O}_x\}_{x \in K}$  is an open cover of  $K$ . Since  $K$  is compact, there is a finite subcover  $\{\mathcal{O}_{x_1}, \mathcal{O}_{x_2}, \dots, \mathcal{O}_{x_n}\}$ . Define  $\mathcal{N} = \bigcap_{i=1}^n \mathcal{U}_{x_i}$ . Then  $\mathcal{N}$  is a neighborhood of  $y$  which is disjoint from each  $\mathcal{O}_{x_i}$  (since  $\mathcal{O}_{x_i}$  and  $\mathcal{U}_{x_i}$  are disjoint). Hence  $\mathcal{N} \subset X \setminus K$  since the  $\{\mathcal{O}_{x_i}\}_{i=1}^n$  cover  $K$ . Since  $y \in X \setminus K$  is arbitrary, then  $X \setminus K$  is open and so  $K$  is closed.  $\square$

# Proposition 11.16

**Proposition 11.16.** A compact subspace  $K$  of a Hausdorff topological space  $(X, \mathcal{T})$  is a closed subset of  $X$ .

**Proof.** If  $K = X$ , then  $K$  is closed. Otherwise, let  $y \in X \setminus K$ . Since  $X$  is Hausdorff, for each  $x \in K$  there are disjoint neighborhoods  $\mathcal{O}_x$  and  $\mathcal{U}_x$  of  $x$  and  $y$  respectively. Then  $\{\mathcal{O}_x\}_{x \in K}$  is an open cover of  $K$ . Since  $K$  is compact, there is a finite subcover  $\{\mathcal{O}_{x_1}, \mathcal{O}_{x_2}, \dots, \mathcal{O}_{x_n}\}$ . Define  $\mathcal{N} = \bigcap_{i=1}^n \mathcal{U}_{x_i}$ . Then  $\mathcal{N}$  is a neighborhood of  $y$  which is disjoint from each  $\mathcal{O}_{x_i}$  (since  $\mathcal{O}_{x_i}$  and  $\mathcal{U}_{x_i}$  are disjoint). Hence  $\mathcal{N} \subset X \setminus K$  since the  $\{\mathcal{O}_{x_i}\}_{i=1}^n$  cover  $K$ . Since  $y \in X \setminus K$  is arbitrary, then  $X \setminus K$  is open and so  $K$  is closed.  $\square$

# Proposition 11.17

**Proposition 11.17.** Let  $(X, \mathcal{T})$  be a second countable topological space. Then  $(X, \mathcal{T})$  is compact if and only if it is sequentially compact.

**Proof.** Let  $(X, \mathcal{T})$  be compact. Let  $\{x_n\}$  be a sequence in  $X$ . For each  $n \in \mathbb{N}$ , let  $F_n$  be the closure of  $\{x_k \mid k \geq n\}$ . Then  $\{F_n\}$  is a decreasing sequence of nonempty closed sets. So  $\{F_n\}$  has the finite intersection property.

## Proposition 11.17

**Proposition 11.17.** Let  $(X, \mathcal{T})$  be a second countable topological space. Then  $(X, \mathcal{T})$  is compact if and only if it is sequentially compact.

**Proof.** Let  $(X, \mathcal{T})$  be compact. Let  $\{x_n\}$  be a sequence in  $X$ . For each  $n \in \mathbb{N}$ , let  $F_n$  be the closure of  $\{x_k \mid k \geq n\}$ . Then  $\{F_n\}$  is a decreasing sequence of nonempty closed sets. So  $\{F_n\}$  has the finite intersection property. By Proposition 11.14,  $\bigcap_{n=1}^{\infty} F_n \neq \emptyset$ , so choose  $x_0 \in \bigcap_{n=1}^{\infty} F_n$ . Since  $X$  is second countable, it is first countable and so has a base  $\{B_n\}_{n=1}^{\infty}$  for the topology at point  $x_0$ . Without loss of generality we may assume  $B_{n+1} \subset B_n$  (or else we could replace  $B_n$  with  $\bigcap_{k=1}^n B_k$  and this then produces a base at  $x$  with this decreasing property). Since  $x_0 \in F_n$  for all  $n \in \mathbb{N}$ , then  $x_0$  is a point of closure of  $\{x_k \mid k \geq n\}$  for all  $n \in \mathbb{N}$ .

# Proposition 11.17

**Proposition 11.17.** Let  $(X, \mathcal{T})$  be a second countable topological space. Then  $(X, \mathcal{T})$  is compact if and only if it is sequentially compact.

**Proof.** Let  $(X, \mathcal{T})$  be compact. Let  $\{x_n\}$  be a sequence in  $X$ . For each  $n \in \mathbb{N}$ , let  $F_n$  be the closure of  $\{x_k \mid k \geq n\}$ . Then  $\{F_n\}$  is a decreasing sequence of nonempty closed sets. So  $\{F_n\}$  has the finite intersection property. By Proposition 11.14,  $\bigcap_{n=1}^{\infty} F_n \neq \emptyset$ , so choose  $x_0 \in \bigcap_{n=1}^{\infty} F_n$ . Since  $X$  is second countable, it is first countable and so has a base  $\{B_n\}_{n=1}^{\infty}$  for the topology at point  $x_0$ . Without loss of generality we may assume  $B_{n+1} \subset B_n$  (or else we could replace  $B_n$  with  $\bigcap_{k=1}^n B_k$  and this then produces a base at  $x$  with this decreasing property). Since  $x_0 \in F_n$  for all  $n \in \mathbb{N}$ , then  $x_0$  is a point of closure of  $\{x_k \mid k \geq n\}$  for all  $n \in \mathbb{N}$ . So neighborhood  $B_n$  of  $x_0$  has nonempty intersection with  $\{x_k \mid k \geq n\}$  for all  $n \in \mathbb{N}$ . So we can inductively choose  $x_{n_k}$  (where the sequence of subscripts  $n_k$  is strictly increasing) in  $B_k$ .

# Proposition 11.17


**Proposition 11.17.** Let  $(X, \mathcal{T})$  be a second countable topological space. Then  $(X, \mathcal{T})$  is compact if and only if it is sequentially compact.

**Proof.** Let  $(X, \mathcal{T})$  be compact. Let  $\{x_n\}$  be a sequence in  $X$ . For each  $n \in \mathbb{N}$ , let  $F_n$  be the closure of  $\{x_k \mid k \geq n\}$ . Then  $\{F_n\}$  is a decreasing sequence of nonempty closed sets. So  $\{F_n\}$  has the finite intersection property. By Proposition 11.14,  $\bigcap_{n=1}^{\infty} F_n \neq \emptyset$ , so choose  $x_0 \in \bigcap_{n=1}^{\infty} F_n$ . Since  $X$  is second countable, it is first countable and so has a base  $\{B_n\}_{n=1}^{\infty}$  for the topology at point  $x_0$ . Without loss of generality we may assume  $B_{n+1} \subset B_n$  (or else we could replace  $B_n$  with  $\bigcap_{k=1}^n B_k$  and this then produces a base at  $x$  with this decreasing property). Since  $x_0 \in F_n$  for all  $n \in \mathbb{N}$ , then  $x_0$  is a point of closure of  $\{x_k \mid k \geq n\}$  for all  $n \in \mathbb{N}$ . So neighborhood  $B_n$  of  $x_0$  has nonempty intersection with  $\{x_k \mid k \geq n\}$  for all  $n \in \mathbb{N}$ . So we can inductively choose  $x_{n_k}$  (where the sequence of subscripts  $n_k$  is strictly increasing) in  $B_k$ .



## Proposition 11.17 (continued 1)


Proof (continued).



Since for each neighborhood  $\mathcal{O}$  of  $x_0$ , there is an index  $N$  for which  $B_n \subset \mathcal{O}$  for  $n \geq N$  (by the definition of base at  $x_0$  and the nestedness of the  $B_n$ 's), the subsequence  $\{x_{n_k}\}$  converges to  $x_0$ . So  $X$  is sequentially compact.

## Proposition 11.17 (continued 1)

Proof (continued).



Since for each neighborhood  $\mathcal{O}$  of  $x_0$ , there is an index  $N$  for which  $B_n \subset \mathcal{O}$  for  $n \geq N$  (by the definition of base at  $x_0$  and the nestedness of the  $B_n$ 's), the subsequence  $\{x_{n_k}\}$  converges to  $x_0$ . So  $X$  is sequentially compact.

## Proposition 11.17 (continued 2)

**Proof (continued).** Suppose  $X$  is sequentially compact. Since  $X$  is second countable, every open cover has a countable subcover (by the definition of 2nd countable). So, to show that  $X$  is compact it suffices to show that every countable open cover of  $X$  has a finite subcover. Let  $\{\mathcal{O}_n\}_{n=1}^{\infty}$  be such a cover. ASSUME there is no finite subcover. Then for each  $n \in \mathbb{N}$ , there is an index  $m(n) > n$  for which  $\mathcal{O}_{m(n)} \not\sim (\cup_{i=1}^n \mathcal{O}_i) \neq \emptyset$  (or else  $\{\mathcal{O}_i\}_{i=1}^n$  is a finite subcover of  $X$ ). So for each  $n \in \mathbb{N}$ , choose  $x_n \in \mathcal{O}_{m(n)} \setminus (\cup_{i=1}^n \mathcal{O}_i)$ . Then since  $X$  is sequentially compact, a subsequence of  $\{x_n\}$  converges to some  $x_0 \in X$ .

## Proposition 11.17 (continued 2)

**Proof (continued).** Suppose  $X$  is sequentially compact. Since  $X$  is second countable, every open cover has a countable subcover (by the definition of 2nd countable). So, to show that  $X$  is compact it suffices to show that every countable open cover of  $X$  has a finite subcover. Let  $\{\mathcal{O}_n\}_{n=1}^{\infty}$  be such a cover. ASSUME there is no finite subcover. Then for each  $n \in \mathbb{N}$ , there is an index  $m(n) > n$  for which  $\mathcal{O}_{m(n)} \sim (\cup_{i=1}^n \mathcal{O}_i) \neq \emptyset$  (or else  $\{\mathcal{O}_i\}_{i=1}^n$  is a finite subcover of  $X$ ). So for each  $n \in \mathbb{N}$ , choose  $x_n \in \mathcal{O}_{m(n)} \cap (\cup_{i=1}^n \mathcal{O}_i)$ . Then since  $X$  is sequentially compact, a subsequence of  $\{x_n\}$  converges to some  $x_0 \in X$ . But  $\{\mathcal{O}_n\}_{n=1}^{\infty}$  is an open cover of  $X$ , so there is some  $\mathcal{O}_N$  that is a neighborhood of  $x_0$ . Therefore, there are infinitely many indices  $n$  for which  $x_n$  belongs to  $\mathcal{O}_N$  (these terms being in the subsequence of  $\{x_n\}$  which converges to  $x_0$ ). But by the construction of  $\{x_n\}$ ,  $x_n \notin \mathcal{O}_N$  for  $n > N$ .

## Proposition 11.17 (continued 2)

**Proof (continued).** Suppose  $X$  is sequentially compact. Since  $X$  is second countable, every open cover has a countable subcover (by the definition of 2nd countable). So, to show that  $X$  is compact it suffices to show that every countable open cover of  $X$  has a finite subcover. Let  $\{\mathcal{O}_n\}_{n=1}^{\infty}$  be such a cover. ASSUME there is no finite subcover. Then for each  $n \in \mathbb{N}$ , there is an index  $m(n) > n$  for which  $\mathcal{O}_{m(n)} \sim (\cup_{i=1}^n \mathcal{O}_i) \neq \emptyset$  (or else  $\{\mathcal{O}_i\}_{i=1}^n$  is a finite subcover of  $X$ ). So for each  $n \in \mathbb{N}$ , choose  $x_n \in \mathcal{O}_{m(n)} \cap (\cup_{i=1}^n \mathcal{O}_i)$ . Then since  $X$  is sequentially compact, a subsequence of  $\{x_n\}$  converges to some  $x_0 \in X$ . But  $\{\mathcal{O}_n\}_{n=1}^{\infty}$  is an open cover of  $X$ , so there is some  $\mathcal{O}_N$  that is a neighborhood of  $x_0$ . Therefore, there are infinitely many indices  $n$  for which  $x_n$  belongs to  $\mathcal{O}_N$  (these terms being in the subsequence of  $\{x_n\}$  which converges to  $x_0$ ). But by the construction of  $\{x_n\}$ ,  $x_n \notin \mathcal{O}_N$  for  $n > N$ . So this CONTRADICTION shows that the assumption that  $X$  is not compact is false.  $\square$

## Proposition 11.17 (continued 2)

**Proof (continued).** Suppose  $X$  is sequentially compact. Since  $X$  is second countable, every open cover has a countable subcover (by the definition of 2nd countable). So, to show that  $X$  is compact it suffices to show that every countable open cover of  $X$  has a finite subcover. Let  $\{\mathcal{O}_n\}_{n=1}^{\infty}$  be such a cover. ASSUME there is no finite subcover. Then for each  $n \in \mathbb{N}$ , there is an index  $m(n) > n$  for which  $\mathcal{O}_{m(n)} \sim (\cup_{i=1}^n \mathcal{O}_i) \neq \emptyset$  (or else  $\{\mathcal{O}_i\}_{i=1}^n$  is a finite subcover of  $X$ ). So for each  $n \in \mathbb{N}$ , choose  $x_n \in \mathcal{O}_{m(n)} \cap (\cup_{i=1}^n \mathcal{O}_i)$ . Then since  $X$  is sequentially compact, a subsequence of  $\{x_n\}$  converges to some  $x_0 \in X$ . But  $\{\mathcal{O}_n\}_{n=1}^{\infty}$  is an open cover of  $X$ , so there is some  $\mathcal{O}_N$  that is a neighborhood of  $x_0$ . Therefore, there are infinitely many indices  $n$  for which  $x_n$  belongs to  $\mathcal{O}_N$  (these terms being in the subsequence of  $\{x_n\}$  which converges to  $x_0$ ). But by the construction of  $\{x_n\}$ ,  $x_n \notin \mathcal{O}_N$  for  $n > N$ . So this CONTRADICTION shows that the assumption that  $X$  is not compact is false.  $\square$

# Theorem 11.18

**Theorem 11.18.** A compact Hausdorff space is normal.

**Proof.** Let  $(X, \mathcal{T})$  be compact and Hausdorff. Let  $F$  be a closed subset of  $X$  and let point  $x \in X \sim F$ . Since  $(X, \mathcal{T})$  is Hausdorff, for each  $y \in F$  there are disjoint neighborhoods  $\mathcal{O}_x$  and  $\mathcal{U}_y$  of  $x$  and  $y$ , respectively. Then  $\{\mathcal{U}_y\}_{y \in F}$  is an open cover of  $F$ .

# Theorem 11.18

**Theorem 11.18.** A compact Hausdorff space is normal.

**Proof.** Let  $(X, \mathcal{T})$  be compact and Hausdorff. Let  $F$  be a closed subset of  $X$  and let point  $x \in X \sim F$ . Since  $(X, \mathcal{T})$  is Hausdorff, for each  $y \in F$  there are disjoint neighborhoods  $\mathcal{O}_x$  and  $\mathcal{U}_y$  of  $x$  and  $y$ , respectively. Then  $\{\mathcal{U}_y\}_{y \in F}$  is an open cover of  $F$ . But  $F$  is a closed subset of a compact space and so by Proposition 11.15 is itself compact. So there is a finite subcover  $\{\mathcal{U}_{y_1}, \mathcal{U}_{y_2}, \dots, \mathcal{U}_{y_n}\}$  of  $F$ . Define  $\mathcal{N} = \bigcap_{i=1}^n \mathcal{O}_{y_i}$ . So  $\mathcal{N}$  is open and  $F \subset \mathcal{N}$  is disjoint from  $\bigcup_{i=1}^n \mathcal{U}_{y_i}$ , a neighborhood of  $F$ . Thus  $(X, \mathcal{T})$  is regular.



# Theorem 11.18

**Theorem 11.18.** A compact Hausdorff space is normal.

**Proof.** Let  $(X, \mathcal{T})$  be compact and Hausdorff. Let  $F$  be a closed subset of  $X$  and let point  $x \in X \sim F$ . Since  $(X, \mathcal{T})$  is Hausdorff, for each  $y \in F$  there are disjoint neighborhoods  $\mathcal{O}_x$  and  $\mathcal{U}_y$  of  $x$  and  $y$ , respectively. Then  $\{\mathcal{U}_y\}_{y \in F}$  is an open cover of  $F$ . But  $F$  is a closed subset of a compact space and so by Proposition 11.15 is itself compact. So there is a finite subcover  $\{\mathcal{U}_{y_1}, \mathcal{U}_{y_2}, \dots, \mathcal{U}_{y_n}\}$  of  $F$ . Define  $\mathcal{N} = \bigcap_{i=1}^n \mathcal{O}_{y_i}$ . So  $\mathcal{N}$  is open and  $F \subset \mathcal{N}$  is disjoint from  $\bigcup_{i=1}^n \mathcal{U}_{y_i}$ , a neighborhood of  $F$ . Thus  $(X, \mathcal{T})$  is regular.

Let  $F$  and  $G$  be disjoint closed sets. Since  $(X, \mathcal{T})$  is regular, for each  $g \in G$  there are disjoint  $\mathcal{V}_g$  and  $\mathcal{W}_g$  such that  $F \subset \mathcal{V}_g$  and  $g \in \mathcal{W}_g$ . Then  $\{\mathcal{V}_g\}_{g \in G}$  is an open cover of  $F$ . By Proposition 11.15,  $F$  is compact and so there is some finite  $\{\mathcal{V}_{g_i}\}_{i=1}^m$  subcover of  $F$ .

# Theorem 11.18

**Theorem 11.18.** A compact Hausdorff space is normal.

**Proof.** Let  $(X, \mathcal{T})$  be compact and Hausdorff. Let  $F$  be a closed subset of  $X$  and let point  $x \in X \sim F$ . Since  $(X, \mathcal{T})$  is Hausdorff, for each  $y \in F$  there are disjoint neighborhoods  $\mathcal{O}_x$  and  $\mathcal{U}_y$  of  $x$  and  $y$ , respectively. Then  $\{\mathcal{U}_y\}_{y \in F}$  is an open cover of  $F$ . But  $F$  is a closed subset of a compact space and so by Proposition 11.15 is itself compact. So there is a finite subcover  $\{\mathcal{U}_{y_1}, \mathcal{U}_{y_2}, \dots, \mathcal{U}_{y_n}\}$  of  $F$ . Define  $\mathcal{N} = \bigcap_{i=1}^n \mathcal{O}_{y_i}$ . So  $\mathcal{N}$  is open and  $F \subset \mathcal{N}$  is disjoint from  $\bigcup_{i=1}^n \mathcal{U}_{y_i}$ , a neighborhood of  $F$ . Thus  $(X, \mathcal{T})$  is regular.

Let  $F$  and  $G$  be disjoint closed sets. Since  $(X, \mathcal{T})$  is regular, for each  $g \in G$  there are disjoint  $\mathcal{V}_g$  and  $\mathcal{W}_g$  such that  $F \subset \mathcal{V}_g$  and  $g \in \mathcal{W}_g$ . Then  $\{\mathcal{V}_g\}_{g \in G}$  is an open cover of  $F$ . By Proposition 11.15,  $F$  is compact and so there is some finite  $\{\mathcal{V}_{g_i}\}_{i=1}^m$  subcover of  $F$ . Then (similar to above) the open sets  $\bigcup_{i=1}^m \mathcal{V}_{g_i}$  and  $\bigcap_{i=1}^m \mathcal{W}_{g_i}$  separates  $F$  and  $G$ . Therefore  $(X, \mathcal{T})$  is normal.  $\square$

# Theorem 11.18

**Theorem 11.18.** A compact Hausdorff space is normal.

**Proof.** Let  $(X, \mathcal{T})$  be compact and Hausdorff. Let  $F$  be a closed subset of  $X$  and let point  $x \in X \sim F$ . Since  $(X, \mathcal{T})$  is Hausdorff, for each  $y \in F$  there are disjoint neighborhoods  $\mathcal{O}_x$  and  $\mathcal{U}_y$  of  $x$  and  $y$ , respectively. Then  $\{\mathcal{U}_y\}_{y \in F}$  is an open cover of  $F$ . But  $F$  is a closed subset of a compact space and so by Proposition 11.15 is itself compact. So there is a finite subcover  $\{\mathcal{U}_{y_1}, \mathcal{U}_{y_2}, \dots, \mathcal{U}_{y_n}\}$  of  $F$ . Define  $\mathcal{N} = \bigcap_{i=1}^n \mathcal{O}_{y_i}$ . So  $\mathcal{N}$  is open and  $F \subset \mathcal{N}$  is disjoint from  $\bigcup_{i=1}^n \mathcal{U}_{y_i}$ , a neighborhood of  $F$ . Thus  $(X, \mathcal{T})$  is regular.

Let  $F$  and  $G$  be disjoint closed sets. Since  $(X, \mathcal{T})$  is regular, for each  $g \in G$  there are disjoint  $\mathcal{V}_g$  and  $\mathcal{W}_g$  such that  $F \subset \mathcal{V}_g$  and  $g \in \mathcal{W}_g$ . Then  $\{\mathcal{V}_g\}_{g \in G}$  is an open cover of  $F$ . By Proposition 11.15,  $F$  is compact and so there is some finite  $\{\mathcal{V}_{g_i}\}_{i=1}^m$  subcover of  $F$ . Then (similar to above) the open sets  $\bigcup_{i=1}^m \mathcal{V}_{g_i}$  and  $\bigcap_{i=1}^m \mathcal{W}_{g_i}$  separates  $F$  and  $G$ . Therefore  $(X, \mathcal{T})$  is normal. □

# Proposition 11.20

**Proposition 11.20.** The continuous image of a compact topological space is compact.

**Proof.** Let  $f$  be a continuous mapping of  $(X, \mathcal{T})$  to  $(Y, \mathcal{S})$ . Let  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  be an open covering of  $f(X)$ . Then since  $f$  is continuous, by Proposition 11.10,  $\{f^{-1}(\mathcal{O}_\lambda)\}_{\lambda \in \Lambda}$  is an open cover of  $X$ .

# Proposition 11.20

**Proposition 11.20.** The continuous image of a compact topological space is compact.

**Proof.** Let  $f$  be a continuous mapping of  $(X, \mathcal{T})$  to  $(Y, \mathcal{S})$ . Let  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  be an open covering of  $f(X)$ . Then since  $f$  is continuous, by Proposition 11.10,  $\{f^{-1}(\mathcal{O}_\lambda)\}_{\lambda \in \Lambda}$  is an open cover of  $X$ . Since  $X$  is compact, there is a finite subcover  $\{f^{-1}(\mathcal{O}_{\lambda_i})\}_{i=1}^n$  of  $X$ . Then the finite collection  $\{\mathcal{O}_{\lambda_i}\}_{i=1}^n$  is a cover of  $f(X)$  and  $f(X)$  is compact.  $\square$

# Proposition 11.20

**Proposition 11.20.** The continuous image of a compact topological space is compact.

**Proof.** Let  $f$  be a continuous mapping of  $(X, \mathcal{T})$  to  $(Y, \mathcal{S})$ . Let  $\{\mathcal{O}_\lambda\}_{\lambda \in \Lambda}$  be an open covering of  $f(X)$ . Then since  $f$  is continuous, by Proposition 11.10,  $\{f^{-1}(\mathcal{O}_\lambda)\}_{\lambda \in \Lambda}$  is an open cover of  $X$ . Since  $X$  is compact, there is a finite subcover  $\{f^{-1}(\mathcal{O}_{\lambda_i})\}_{i=1}^n$  of  $X$ . Then the finite collection  $\{\mathcal{O}_{\lambda_i}\}_{i=1}^n$  is a cover of  $f(X)$  and  $f(X)$  is compact.  $\square$

# Corollary 11.21

**Corollary 11.21.** A continuous real-valued function on a compact topological space takes a maximum and minimum functional value.

**Proof.** Let  $(X, \mathcal{T})$  be compact  $f : X \rightarrow \mathbb{R}$  be continuous. By Proposition 11.20,  $f(X)$  is a compact set of real numbers. So, by the Heine-Borel Theorem,  $f(X)$  is closed and bounded.

# Corollary 11.21

**Corollary 11.21.** A continuous real-valued function on a compact topological space takes a maximum and minimum functional value.

**Proof.** Let  $(X, \mathcal{T})$  be compact  $f : X \rightarrow \mathbb{R}$  be continuous. By Proposition 11.20,  $f(X)$  is a compact set of real numbers. So, by the Heine-Borel Theorem,  $f(X)$  is closed and bounded. So  $f$  attains a maximum and minimum value (namely,  $\sup f(X) = \max f(X)$  and  $\inf f(X) = \min f(X)$ ). □



# Corollary 11.21

**Corollary 11.21.** A continuous real-valued function on a compact topological space takes a maximum and minimum functional value.

**Proof.** Let  $(X, \mathcal{T})$  be compact  $f : X \rightarrow \mathbb{R}$  be continuous. By Proposition 11.20,  $f(X)$  is a compact set of real numbers. So, by the Heine-Borel Theorem,  $f(X)$  is closed and bounded. So  $f$  attains a maximum and minimum value (namely,  $\sup f(X) = \max f(X)$  and  $\inf f(X) = \min f(X)$ ). □

# Proposition 11.19

**Proposition 11.19.** A continuous one to one mapping  $f$  of a compact space  $(X, \mathcal{T})$  onto a Hausdorff space  $Y$  is a homeomorphism.

**Proof.** Since  $f$  is given to be continuous, one to one, and onto then we need only show that  $f^{-1}$  is continuous. This can be done by showing  $f$  maps open sets to closed sets or, equivalently,  $f$  maps closed sets to closed sets.

# Proposition 11.19

**Proposition 11.19.** A continuous one to one mapping  $f$  of a compact space  $(X, \mathcal{T})$  onto a Hausdorff space  $Y$  is a homeomorphism.

**Proof.** Since  $f$  is given to be continuous, one to one, and onto then we need only show that  $f^{-1}$  is continuous. This can be done by showing  $f$  maps open sets to closed sets or, equivalently,  $f$  maps closed sets to closed sets. Let  $F$  be a closed subset of  $X$ . Then  $F$  is compact by Proposition 11.15. By Proposition 11.20,  $f(F)$  is compact in  $(Y, \mathcal{S})$ . Hence by Proposition 11.16, since  $Y$  is Hausdorff,  $f(F)$  is closed.

# Proposition 11.19

**Proposition 11.19.** A continuous one to one mapping  $f$  of a compact space  $(X, \mathcal{T})$  onto a Hausdorff space  $Y$  is a homeomorphism.

**Proof.** Since  $f$  is given to be continuous, one to one, and onto then we need only show that  $f^{-1}$  is continuous. This can be done by showing  $f$  maps open sets to closed sets or, equivalently,  $f$  maps closed sets to closed sets. Let  $F$  be a closed subset of  $X$ . Then  $F$  is compact by Proposition 11.15. By Proposition 11.20,  $f(F)$  is compact in  $(Y, \mathcal{S})$ . Hence by Proposition 11.16, since  $Y$  is Hausdorff,  $f(F)$  is closed. Therefore  $f^{-1}$  is continuous and  $f$  is a homeomorphism.  $\square$

# Proposition 11.19

**Proposition 11.19.** A continuous one to one mapping  $f$  of a compact space  $(X, \mathcal{T})$  onto a Hausdorff space  $Y$  is a homeomorphism.

**Proof.** Since  $f$  is given to be continuous, one to one, and onto then we need only show that  $f^{-1}$  is continuous. This can be done by showing  $f$  maps open sets to closed sets or, equivalently,  $f$  maps closed sets to closed sets. Let  $F$  be a closed subset of  $X$ . Then  $F$  is compact by Proposition 11.15. By Proposition 11.20,  $f(F)$  is compact in  $(Y, \mathcal{S})$ . Hence by Proposition 11.16, since  $Y$  is Hausdorff,  $f(F)$  is closed. Therefore  $f^{-1}$  is continuous and  $f$  is a homeomorphism. □