

# Real Analysis 1, MATH 5210, Fall 2022

## Homework 1, Riemann-Lebesgue Theorem

Due Saturday, August 27, at 11:59 p.m.

**Write in complete sentences!!!** *Explain* what you are doing and convince me that you understand what you are doing and why. Justify all steps by quoting relevant results from the textbook, class notes, or hypotheses. Do not copy the work of others; **do your own work!!!**

**RI.B.** Prove Theorem 6-4: A bounded function  $f$  defined on  $[a, b]$  is Riemann integrable on  $[a, b]$  if and only if, given  $\varepsilon > 0$ , there is a partition  $P(\varepsilon)$  of  $[a, b]$  such that  $\overline{S}(f; P(\varepsilon)) - \underline{S}(f; P(\varepsilon)) < \varepsilon$ .  
HINT: If  $P$  and  $Q$  are partitions of  $[a, b]$  with  $P \subset Q$  then  $Q$  is a *refinement* of  $P$ . You may assume  $\underline{S}(f; P) \leq \underline{S}(f; Q)$  and  $\overline{S}(f; Q) \leq \overline{S}(f; P)$  (this is Exercise RI.G).

**RI.C.** Prove Theorem 6-5: Suppose  $f$  is a Riemann integrable function on  $[a, b]$ . If  $I$  is a number such that  $\underline{S}(f; P) \leq I \leq \overline{S}(f; P)$  for every partition  $P$  of  $[a, b]$ , then  $I = \int_a^b f$ .

**RI.D.** Prove that if bounded function  $f$  is Riemann integrable on  $[a, b]$  then there is a number  $I$  such that, given  $\varepsilon > 0$ , there is a partition  $P_\varepsilon$  such that, for any Riemann sum  $S(f; P_\varepsilon)$  of  $f$  with respect to  $P_\varepsilon$ ,  $|S(f; P_\varepsilon) - I| < \varepsilon$ . NOTE: The converse of this result also holds (see Exercise RI.E). Therefore this result and its converse could be used as the definition of the Riemann integral of  $f$  on  $[a, b]$  since this is an if and only if result. Notice that there is no explicit mention of suprema nor infima here, only function values (function values are in in the Riemann sum). This is the approach to Riemann integration taken in Calculus 1.

**RI.G. (a)** Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition of  $[a, b]$  and let

$$Q = P \cup \{x'\} = \{x_0, x_1, \dots, x_k, x', x_{k+1}, \dots, x_n\}$$

( $Q$  is a *one point refinement* of  $P$ ). Prove that  $\underline{S}(f; P) \leq \underline{S}(f; Q)$ . You may assume that  $A \subset B$  implies  $\inf(B) \leq \inf(A)$ .

**(b)** Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition of  $[a, b]$  and let  $Q$  be a partition of  $[a, b]$  such that  $P \subset Q$  ( $Q$  is called a *refinement* of  $P$ ). Prove that  $\underline{S}(f; P) \leq \underline{S}(f; Q)$ . NOTE: A similar argument shows that  $\overline{S}(f; P) \geq \overline{S}(f; Q)$ .