


Complex Variables

Chapter 4. Integrals

Section 4.49. Multiply Connected Domains—Proofs of Theorems



Theorem 4.49.A

Theorem 4.49.A. Suppose that

- (a) C is a simple closed contour, parameterized in the counterclockwise direction; and
- (b) C_k , for $k = 1, 2, \dots, n$, are simple closed contours interior to C , all parameterized in the clockwise direction, that are disjoint and whose interiors have no points in common (so one such contour cannot be inside another).

If a function f is analytic on all of these contours and throughout the multiply connected domain consisting of all points inside C and exterior to each C_k , then

$$\int_C f(z) dz + \sum_{k=1}^n \int_{C_k} f(z) dz = 0.$$

“Proof.” We assume some obvious properties of simple closed contours and assume that we can introduce a polygonal path L_1 consisting of a finite number of line segments joined end to end to connect the outer contour C to the inner contour C_1 .

Theorem 4.49.A (continued 1)

“Proof” continued. Introduce polygonal path L_2 connecting C_1 to C_2 , and in general polygonal path L_{k+1} connecting C_k to C_{k+1} for $k = 1, 2, \dots, n - 1$. Introduce polygonal path L_{n+1} connecting C_n to outer contour C . See Figure 60 (In Figure 60, each polygonal path is a single line segment and the L_k are disjoint; these need not be the case).




FIGURE 60

As shown in Figure 60, two simple closed contours Γ_1 and Γ_2 can be formed, each consisting of polygonal paths L_k or $-L_k$ and pieces of C and C_k and each described in such a direction that the points enclosed by them “lie to the left.” (This, granted, is an informal “proof by picture.”)

Theorem 4.49.A (continued 2)

“Proof” continued. By the Cauchy-Goursat Theorem (Theorem 4.46.A), $\int_{\Gamma_1} f(z) dz = \int_{\Gamma_2} f(z) dz = 0$. So (see Section 40)

$$\int_{\Gamma_1} f(z) dz + \int_{\Gamma_2} f(z) dz = 0 = \int_{\Gamma_1 \cup \Gamma_2} f(z) dz.$$

Now for each $k = 1, 2, \dots, n + 1$, the integral $\int_{L_k} f(z) dz$ and $\int_{-L_k} f(z) dz$ are parts of $\int_{\Gamma_1 \cup \Gamma_2} f(z) dz$, but $\int_{-L_k} f(z) dz = -\int_{L_k} f(z) dz$ (see Note 4.40.B) so these cancel each other and

$$0 = \int_{\Gamma_1 \cup \Gamma_2} f(z) dz = \int_C f(z) dz + \sum_{k=1}^n \int_{C_k} f(z) dz,$$

as claimed. □

Corollary 4.49.B

Corollary 4.49.B. Principle of Deformation.

Let C_1 and C_2 denote positively oriented simple closed contours, where C_1 is interior to C_2 . If a function f is analytic in the closed region consisting of those contours and all points between them, then


$$\int_{C_1} f(z) dz = \int_{C_2} f(z) dz.$$

Proof. By Theorem 4.49.A,

$$\int_{C_2} f(z) dz + \int_{-C_1} f(z) dz = 0$$

(we need to use $-C_1$ to get the inner contour parameterized in a clockwise direction; see Figure 61).

Corollary 4.49.B

Proof (continued).

So

$$\int_{C_2} f(z) dz = - \int_{-C_1} f(z) dz = \int_{C_1} f(z) dz$$

(see Note 4.40.B). □