


# Complex Variables

## Chapter 4. Integrals

### Section 4.50. Cauchy Integral Formula—Proofs of Theorems



## Theorem 4.50.A

### Theorem 4.50.A. Cauchy Integral Formula.

Let  $f$  be analytic everywhere inside and on simple closed contour  $C$ , parameterized in the positive sense. If  $z_0$  is any point interior to  $C$ , then

$$f(z_0) = \frac{1}{2\pi i} \int_C \frac{f(z) dz}{z - z_0}.$$

**Proof.** Let  $C_\rho$  denote the positively oriented circle  $|z - z_0| = \rho$ , where  $\rho$  is small enough the  $C_\rho$  is interior to  $C$  (which can be done since  $C$  is a closed set and so  $\mathbb{C} \setminus C$  is open with  $z_0$  as an interior point of the open set  $\mathbb{C} \setminus C$ ; see Figure 66).




FIGURE 66

## Theorem 4.50.A (continued 1)

**Proof (continued).** The function  $f(z)/(z - z_0)$  is analytic inside and on  $C$  except at  $z_0$ . So by the Principle of Deformation (Corollary 4.49.B),

$$\int_C \frac{f(z) dz}{z - z_0} = \int_{C_\rho} \frac{f(z) dz}{z - z_0}. \text{ So}$$

$$\int_C \frac{f(z) dz}{z - z_0} - f(z_0) \int_{C_\rho} \frac{dz}{z - z_0} = \int_{C_\rho} \frac{f(z) - f(z_0)}{z - z_0} dz. \text{ Next,}$$

$$\int_{C_\rho} \frac{dz}{z - z_0} = 2\pi i \text{ by Exercise 42.10(b), so}$$

$$\int_C \frac{f(z) dz}{z - z_0} - 2\pi i f(z_0) = \int_{C_\rho} \frac{f(z) - f(z_0)}{z - z_0} dz. \quad (4)$$

Since  $f$  is analytic, then it is continuous at  $z_0$  and so for all  $\varepsilon > 0$  there is  $\delta > 0$  such that if  $|z - z_0| < \delta$  then  $|f(z) - f(z_0)| < \varepsilon/(2\pi)$ . The only restriction on  $\rho$  above is that  $C_\rho$  is interior to  $C$ . Let  $\rho' = \min\{\rho, \delta/2\}$ . Then  $C_{\rho'}$  is interior to  $C$  and so the equations above involving  $C_\rho$  also hold for  $C_{\rho'}$ .

## Theorem 4.50.A (continued 2)

**Proof (continued).** Then for  $z$  on  $C_{\rho'}$  we have  $|z - z_0| = \rho' \leq \delta/2 < \delta$  and so  $|f(z) - f(z_0)| < \varepsilon/(2\pi)$ ; also the length of  $C_{\rho'}$  is  $2\pi\rho'$  and so by Theorem 4.43.A,

$$\left| \int_{C_{\rho'}} \frac{f(z) - f(z_0)}{z - z_0} dz \right| \leq \left( \frac{\varepsilon/(2\pi)}{\rho'} \right) (2\pi\rho') = \varepsilon.$$

So by equation (4),

$$\left| \int_{C_{\rho'}} \frac{f(z) - f(z_0)}{z - z_0} dz \right| = \left| \int_C \frac{f(z) dz}{z - z_0} - 2\pi i f(z_0) \right| < \varepsilon.$$

Since  $\varepsilon > 0$  is arbitrary, then the quantity  $\int_C \frac{f(z) dz}{z - z_0} - 2\pi i f(z_0)$  must be 0, and the result follows.  $\square$