

Introduction to Functional Analysis

Chapter 3. Major Banach Space Theorems

3.6. Uniform Boundedness Principle—Proofs of Theorems




Table of contents

1 Theorem 3.10. Uniform Boundedness Principle

2 Theorem 3.11

Theorem 3.10. Uniform Boundedness Principle

Theorem 3.10. Uniform Boundedness Principle.

If X is complete, then a pointwise bounded subset \mathcal{A} of $\mathcal{B}(X, Y)$ is bounded.

Proof. We replace Y with its completion using Theorem 2.22. We now show boundedness on the completion of Y , which certainly implies boundedness on Y itself. Let $\mathcal{A} \subseteq \mathcal{B}(X, Y)$ be a pointwise bounded set. Define the direct product \mathcal{Y} over set \mathcal{A} with each space equal to Y : $\mathcal{Y} = \prod_{T \in \mathcal{A}} Y$. Then the elements of the direct product are the bounded functions (under the sup norm) mapping \mathcal{A} to Y (see the definition of direct product in [Section 2.10. Direct Products and Sums](#)).

Theorem 3.10. Uniform Boundedness Principle

Theorem 3.10. Uniform Boundedness Principle.

If X is complete, then a pointwise bounded subset \mathcal{A} of $\mathcal{B}(X, Y)$ is bounded.

Proof. We replace Y with its completion using Theorem 2.22. We now show boundedness on the completion of Y , which certainly implies boundedness on Y itself. Let $\mathcal{A} \subseteq \mathcal{B}(X, Y)$ be a pointwise bounded set. Define the direct product \mathcal{Y} over set \mathcal{A} with each space equal to Y : $\mathcal{Y} = \prod_{T \in \mathcal{A}} Y$. Then the elements of the direct product are the bounded functions (under the sup norm) mapping \mathcal{A} to Y (see the definition of direct product in [Section 2.10. Direct Products and Sums](#)).

Define $\mathcal{T}x(T) = Tx$ for all $T \in \mathcal{A}$. So \mathcal{T} has input value x and produces $\mathcal{T}x \in \mathcal{Y}$, where $\mathcal{T}x \in \mathcal{Y}$ is a mapping of \mathcal{A} into Y :

$$\mathcal{T} : X \rightarrow \mathcal{Y} \text{ and for } x \in X \text{ we have } \mathcal{T}x : \mathcal{A} \rightarrow Y.$$

Theorem 3.10. Uniform Boundedness Principle

Theorem 3.10. Uniform Boundedness Principle.

If X is complete, then a pointwise bounded subset \mathcal{A} of $\mathcal{B}(X, Y)$ is bounded.

Proof. We replace Y with its completion using Theorem 2.22. We now show boundedness on the completion of Y , which certainly implies boundedness on Y itself. Let $\mathcal{A} \subseteq \mathcal{B}(X, Y)$ be a pointwise bounded set. Define the direct product \mathcal{Y} over set \mathcal{A} with each space equal to Y : $\mathcal{Y} = \prod_{T \in \mathcal{A}} Y$. Then the elements of the direct product are the bounded functions (under the sup norm) mapping \mathcal{A} to Y (see the definition of direct product in [Section 2.10. Direct Products and Sums](#)).

Define $\mathcal{T}x(T) = Tx$ for all $T \in \mathcal{A}$. So \mathcal{T} has input value x and produces $Tx \in \mathcal{Y}$, where $\mathcal{T}x \in \mathcal{Y}$ is a mapping of \mathcal{A} into Y :

$$\mathcal{T} : X \rightarrow \mathcal{Y} \text{ and for } x \in X \text{ we have } \mathcal{T}x : \mathcal{A} \rightarrow Y.$$

Theorem 3.10 (continued 1)

Proof (continued). Since \mathcal{A} is hypothesized to be pointwise bounded, for each $x \in X$, $\mathcal{O}_x = \{\|Tx\| \mid T \in \mathcal{A}\}$ is bounded, say by K_x . So for $Tx \in \mathcal{Y}$,

$$\begin{aligned}\|Tx\| &= \sup\{\|TxT\| \mid T \in \mathcal{A}, \|T\| = 1\} \\ &= \sup\{\|Tx\| \mid T \in \mathcal{A}, \|T\| = 1\} \leq K_x.\end{aligned}$$

So Tx is a bounded function on \mathcal{A} for each $x \in X$.

Now for the boundedness of \mathcal{A} . Suppose that $(x_n) \subseteq X$ converges to $x \in X$, and suppose that $(Tx_n) \subseteq \mathcal{A}$ converges to some $g : \mathcal{A} \rightarrow Y$. We now show that $g = Tx$ (by showing they agree for all $T \in \mathcal{A}$). Indeed,

$$\begin{aligned}g(T) &= \lim_{n \rightarrow \infty} (Tx_n)(T) \text{ by assumption on convergence of } (Tx_n) \\ &= \lim_{n \rightarrow \infty} Tx_n \text{ since } Tx_n(T) = Tx_n \text{ by the definition of } T \\ &= T(\lim x_n) \text{ since } T \in \mathcal{A} \subseteq \mathcal{B}(X, Y) \text{ is bounded and so} \\ &\quad \text{continuous by Theorem 2.6} \\ &= Tx \text{ since } (x_n) \rightarrow x \dots\end{aligned}$$

Theorem 3.10 (continued 1)

Proof (continued). Since \mathcal{A} is hypothesized to be pointwise bounded, for each $x \in X$, $\mathcal{O}_x = \{\|Tx\| \mid T \in \mathcal{A}\}$ is bounded, say by K_x . So for $Tx \in \mathcal{Y}$,

$$\begin{aligned}\|Tx\| &= \sup\{\|TxT\| \mid T \in \mathcal{A}, \|T\| = 1\} \\ &= \sup\{\|Tx\| \mid T \in \mathcal{A}, \|T\| = 1\} \leq K_x.\end{aligned}$$

So Tx is a bounded function on \mathcal{A} for each $x \in X$.

Now for the boundedness of \mathcal{A} . Suppose that $(x_n) \subseteq X$ converges to $x \in X$, and suppose that $(Tx_n) \subseteq \mathcal{A}$ converges to some $g : \mathcal{A} \rightarrow Y$. We now show that $g = Tx$ (by showing they agree for all $T \in \mathcal{A}$). Indeed,

$$\begin{aligned}g(T) &= \lim_{n \rightarrow \infty} (Tx_n)(T) \text{ by assumption on convergence of } (Tx_n) \\ &= \lim_{n \rightarrow \infty} Tx_n \text{ since } Tx_n(T) = Tx_n \text{ by the definition of } T \\ &= T(\lim x_n) \text{ since } T \in \mathcal{A} \subseteq \mathcal{B}(X, Y) \text{ is bounded and so} \\ &\quad \text{continuous by Theorem 2.6} \\ &= Tx \text{ since } (x_n) \rightarrow x \dots\end{aligned}$$

Theorem 3.10 (continued 2)

Proof (continued). ...

$$\begin{aligned} g(T) &= Tx \text{ since } (x_n) \rightarrow x \\ &= Tx(T) \text{ by definition of } T. \end{aligned}$$

So for any sequence $(x_n, Tx_n) \rightarrow (x, Tx)$ in $X \times \mathcal{Y}$ (with respect to the sup norm on $X \times \mathcal{Y}$), we have $(x_n) \rightarrow x$ and $(Tx_n) \rightarrow Tx$. Hence the graph of T is closed. So by the Closed Graph Theorem (Theorem 3.9), T is bounded. That is,

$$\|T\| = \sup\{\|Tx\| \mid x \in X, \|x\| = 1\} = K < \infty.$$

Hence, for any $T \in \mathcal{A}$ and for any $x \in X$ with $\|x\| = 1$, we have $\|Tx(T)\| = \|Tx\| \leq K$. Since $\|Tx\| \leq K$ for all unit vectors $x \in X$, then $\|T\| \leq K$. This holds for all $T \in \mathcal{A}$ so that \mathcal{A} is bounded (by K), as claimed. □

Theorem 3.10 (continued 2)

Proof (continued). ...

$$\begin{aligned} g(T) &= Tx \text{ since } (x_n) \rightarrow x \\ &= Tx(T) \text{ by definition of } T. \end{aligned}$$

So for any sequence $(x_n, Tx_n) \rightarrow (x, Tx)$ in $X \times Y$ (with respect to the sup norm on $X \times Y$), we have $(x_n) \rightarrow x$ and $(Tx_n) \rightarrow Tx$. Hence the graph of T is closed. So by the Closed Graph Theorem (Theorem 3.9), T is bounded. That is,

$$\|T\| = \sup\{\|Tx\| \mid x \in X, \|x\| = 1\} = K < \infty.$$

Hence, for any $T \in \mathcal{A}$ and for any $x \in X$ with $\|x\| = 1$, we have $\|Tx(T)\| = \|Tx\| \leq K$. Since $\|Tx\| \leq K$ for all unit vectors $x \in X$, then $\|T\| \leq K$. This holds for all $T \in \mathcal{A}$ so that \mathcal{A} is bounded (by K), as claimed. □

Theorem 3.11

Theorem 3.11. Suppose that (T_n) is a pointwise convergent sequence of bounded linear operators from Banach space X to normed linear space Y . That is, for each $x \in X$ the sequence $(T_n x)$ converges to an element $Tx \in Y$. Then T is linear and bounded.

Proof. For linearity, let $\alpha x + \beta y \in X$. Then

$$\begin{aligned} T(\alpha x + \beta y) &= \lim_{n \rightarrow \infty} T_n(\alpha x + \beta y) = \lim_{n \rightarrow \infty} (\alpha T_n(x) + \beta T_n(y)) \\ &= \alpha \lim_{n \rightarrow \infty} (T_n(x)) + \beta \lim_{n \rightarrow \infty} (T_n(y)) = \alpha T(x) + \beta T(y) \end{aligned}$$

and T is linear, as claimed.

Theorem 3.11

Theorem 3.11. Suppose that (T_n) is a pointwise convergent sequence of bounded linear operators from Banach space X to normed linear space Y . That is, for each $x \in X$ the sequence $(T_n x)$ converges to an element $Tx \in Y$. Then T is linear and bounded.

Proof. For linearity, let $\alpha x + \beta y \in X$. Then

$$\begin{aligned} T(\alpha x + \beta y) &= \lim_{n \rightarrow \infty} T_n(\alpha x + \beta y) = \lim_{n \rightarrow \infty} (\alpha T_n(x) + \beta T_n(y)) \\ &= \alpha \lim_{n \rightarrow \infty} (T_n(x)) + \beta \lim_{n \rightarrow \infty} (T_n(x)) = \alpha T(x) + \beta T(y) \end{aligned}$$

and T is linear, as claimed.

Now for boundedness. For $x \in X$, the sequence $(T_n x)$ converges and so is bounded by Theorem 2.9(a) and (b). So $\{T_n \mid n \in \mathbb{N}\}$ is a pointwise bounded set; that is, for each $x \in X$ the set $\mathcal{O}_x = \{T_n x \mid n \in \mathbb{N}\}$ is bounded. By the Uniform Boundedness Principle, there is $K > 0$ such that $\|T_n\| \leq K$ for all $n \in \mathbb{N}$.

Theorem 3.11

Theorem 3.11. Suppose that (T_n) is a pointwise convergent sequence of bounded linear operators from Banach space X to normed linear space Y . That is, for each $x \in X$ the sequence $(T_n x)$ converges to an element $Tx \in Y$. Then T is linear and bounded.

Proof. For linearity, let $\alpha x + \beta y \in X$. Then

$$\begin{aligned} T(\alpha x + \beta y) &= \lim_{n \rightarrow \infty} T_n(\alpha x + \beta y) = \lim_{n \rightarrow \infty} (\alpha T_n(x) + \beta T_n(y)) \\ &= \alpha \lim_{n \rightarrow \infty} (T_n(x)) + \beta \lim_{n \rightarrow \infty} (T_n(x)) = \alpha T(x) + \beta T(y) \end{aligned}$$

and T is linear, as claimed.

Now for boundedness. For $x \in X$, the sequence $(T_n x)$ converges and so is bounded by Theorem 2.9(a) and (b). So $\{T_n \mid n \in \mathbb{N}\}$ is a pointwise bounded set; that is, for each $x \in X$ the set $\mathcal{O}_x = \{T_n x \mid n \in \mathbb{N}\}$ is bounded. By the Uniform Boundedness Principle, there is $K > 0$ such that $\|T_n\| \leq K$ for all $n \in \mathbb{N}$.

Theorem 3.11 (continued)

Theorem 3.11. Suppose that (T_n) is a pointwise convergent sequence of bounded linear operators from Banach space X to normed linear space Y . That is, for each $x \in X$ the sequence $(T_n x)$ converges to an element $Tx \in Y$. Then T is linear and bounded.

Proof (continued). Then, for $x \in X$ a unit vector, we have

$$\begin{aligned} \|Tx\| &= \|\lim T_n x\| \text{ by the convergence hypothesis} \\ &= \lim \|T_n x\| \text{ since } \|\cdot\| \text{ is continuous by Theorem 2.3(c)} \\ &\leq K. \end{aligned}$$

So $\|T\| = \sup\{\|Tx\| \mid x \in X, \|x\| = 1\} \leq K$ and T is bounded, as claimed. □