


# Calculus 1

## Chapter 3. Derivatives

### 3.7. Implicit Differentiation—Examples and Proofs



# Table of contents

- 1 Example 3.7.A
- 2 Exercise 3.7.16
- 3 Exercise 3.7.20
- 4 Exercise 3.7.40
- 5 Exercise 3.7.44. Normals Parallel to a Line
- 6 Exercise 3.7.22
- 7 Exercise 3.7.48. The Folium of Descartes
- 8 Exercise 3.7.50. Power Rule for Rational Exponents

## Example 3.7.A

**Example 3.7.A.** Find the slope of the line tangent to  $x^2 + y^2 = 1$  at  $(x, y) = (\sqrt{2}/2, \sqrt{2}/2)$ . Do the same for the point  $(x, y) = (\sqrt{2}/2, -\sqrt{2}/2)$ .

**Solution.** We have just seen that implicit differentiation gives  $dy/dx = -x/y$ .

## Example 3.7.A

**Example 3.7.A.** Find the slope of the line tangent to  $x^2 + y^2 = 1$  at  $(x, y) = (\sqrt{2}/2, \sqrt{2}/2)$ . Do the same for the point  $(x, y) = (\sqrt{2}/2, -\sqrt{2}/2)$ .

**Solution.** We have just seen that implicit differentiation gives  $dy/dx = -x/y$ . So at  $(\sqrt{2}/2, \sqrt{2}/2)$ , the slope of a line tangent to  $x^2 + y^2 = 1$  is

$$\left. \frac{dy}{dx} \right|_{(x,y)=(\sqrt{2}/2, \sqrt{2}/2)} = -\frac{(\sqrt{2}/2)}{(\sqrt{2}/2)} = \boxed{-1}.$$

## Example 3.7.A

**Example 3.7.A.** Find the slope of the line tangent to  $x^2 + y^2 = 1$  at  $(x, y) = (\sqrt{2}/2, \sqrt{2}/2)$ . Do the same for the point  $(x, y) = (\sqrt{2}/2, -\sqrt{2}/2)$ .

**Solution.** We have just seen that implicit differentiation gives  $dy/dx = -x/y$ . So at  $(\sqrt{2}/2, \sqrt{2}/2)$ , the slope of a line tangent to  $x^2 + y^2 = 1$  is

$$\left. \frac{dy}{dx} \right|_{(x,y)=(\sqrt{2}/2, \sqrt{2}/2)} = -\frac{(\sqrt{2}/2)}{(\sqrt{2}/2)} = \boxed{-1}.$$

At  $(\sqrt{2}/2, -\sqrt{2}/2)$ , the slope of a line tangent to  $x^2 + y^2 = 1$  is

$$\left. \frac{dy}{dx} \right|_{(x,y)=(\sqrt{2}/2, -\sqrt{2}/2)} = -\frac{(\sqrt{2}/2)}{(-\sqrt{2}/2)} = \boxed{1}.$$

## Example 3.7.A

**Example 3.7.A.** Find the slope of the line tangent to  $x^2 + y^2 = 1$  at  $(x, y) = (\sqrt{2}/2, \sqrt{2}/2)$ . Do the same for the point  $(x, y) = (\sqrt{2}/2, -\sqrt{2}/2)$ .

**Solution.** We have just seen that implicit differentiation gives  $dy/dx = -x/y$ . So at  $(\sqrt{2}/2, \sqrt{2}/2)$ , the slope of a line tangent to  $x^2 + y^2 = 1$  is


$$\left. \frac{dy}{dx} \right|_{(x,y)=(\sqrt{2}/2, \sqrt{2}/2)} = -\frac{(\sqrt{2}/2)}{(\sqrt{2}/2)} = \boxed{-1}.$$

At  $(\sqrt{2}/2, -\sqrt{2}/2)$ , the slope of a line tangent to  $x^2 + y^2 = 1$  is

$$\left. \frac{dy}{dx} \right|_{(x,y)=(\sqrt{2}/2, -\sqrt{2}/2)} = -\frac{(\sqrt{2}/2)}{(-\sqrt{2}/2)} = \boxed{1}.$$

## Example 3.7.A (continued)

Solution (continued).



Since  $x^2 + y^2 = 1$  does not determine a single function (it fails the vertical line test), then to find the slope of a tangent to the graph of the equation, we need both an  $x$  value and a  $y$  for the point of tangency.  $\square$

## Exercise 3.7.16

**Exercise 3.7.16.** Find  $dy/dx$  for  $y$  an implicit function of  $x$  given by the equation  $e^{x^2y} = 2x + 2y$ .

**Solution.** Differentiating implicitly,  $\frac{d}{dx}[e^{x^2y}] = \frac{d}{dx}[2x + 2y]$  or

$e^{x^2y} \widehat{[2x]}(y) + (x^2)[dy/dx] = 2[1] + 2[dy/dx]$  (notice that since  $y$  is a function of  $x$  then we must use the Derivative Product Rule to differentiate  $x^2y$ ).



## Exercise 3.7.16

**Exercise 3.7.16.** Find  $dy/dx$  for  $y$  an implicit function of  $x$  given by the equation  $e^{x^2y} = 2x + 2y$ .

**Solution.** Differentiating implicitly,  $\frac{d}{dx}[e^{x^2y}] = \frac{d}{dx}[2x + 2y]$  or

$e^{x^2y} \overbrace{[[2x](y) + (x^2)[dy/dx]]} = 2[1] + 2[dy/dx]$  (notice that since  $y$  is a function of  $x$  then we must use the Derivative Product Rule to differentiate  $x^2y$ ). Solving for  $dy/dx$  we get  $e^{x^2y}(2xy + x^2(dy/dx)) = 2 + 2(dy/dx)$  or  $2xye^{x^2y} + x^2e^{x^2y}(dy/dx) = 2 + 2(dy/dx)$  or

$$x^2e^{x^2y}(dy/dx) - 2(dy/dx) = -2xye^{x^2y} + 2 \text{ or } \boxed{\frac{dy}{dx} = \frac{-2xye^{x^2y} + 2}{x^2e^{x^2y} - 2}}. \quad \square$$

## Exercise 3.7.16

**Exercise 3.7.16.** Find  $dy/dx$  for  $y$  an implicit function of  $x$  given by the equation  $e^{x^2y} = 2x + 2y$ .

**Solution.** Differentiating implicitly,  $\frac{d}{dx}[e^{x^2y}] = \frac{d}{dx}[2x + 2y]$  or

$e^{x^2y} \overbrace{[[2x](y) + (x^2)[dy/dx]]} = 2[1] + 2[dy/dx]$  (notice that since  $y$  is a function of  $x$  then we must use the Derivative Product Rule to differentiate  $x^2y$ ). Solving for  $dy/dx$  we get  $e^{x^2y}(2xy + x^2(dy/dx)) = 2 + 2(dy/dx)$  or  $2xye^{x^2y} + x^2e^{x^2y}(dy/dx) = 2 + 2(dy/dx)$  or

$$x^2e^{x^2y}(dy/dx) - 2(dy/dx) = -2xye^{x^2y} + 2 \text{ or } \boxed{\frac{dy}{dx} = \frac{-2xye^{x^2y} + 2}{x^2e^{x^2y} - 2}}. \quad \square$$

## Exercise 3.7.20

**Exercise 3.7.20.** Find  $dr/d\theta$  for  $r$  an implicit function of  $\theta$  given by the equation  $\cos r + \cot \theta = e^{r\theta}$ .

**Solution.** Differentiating implicitly,  $\frac{d}{d\theta}[\cos r + \cot \theta] = \frac{d}{d\theta}[e^{r\theta}]$  or  
 $-\sin r \overset{\curvearrowright}{[dr/d\theta]} - \csc^2 \theta = e^{r\theta} \overset{\curvearrowright}{[dr/d\theta]} (\theta) + (r)[1]$  (notice that since  $r$  is a function of  $\theta$  then we must use the Derivative Product Rule to differentiate  $r\theta$ ) or  $-\sin r(dr/d\theta) - \csc^2 \theta = \theta e^{r\theta}(dr/d\theta) + re^{r\theta}$ .

## Exercise 3.7.20

**Exercise 3.7.20.** Find  $dr/d\theta$  for  $r$  an implicit function of  $\theta$  given by the equation  $\cos r + \cot \theta = e^{r\theta}$ .

**Solution.** Differentiating implicitly,  $\frac{d}{d\theta}[\cos r + \cot \theta] = \frac{d}{d\theta}[e^{r\theta}]$  or  
 $-\sin r \widehat{[dr/d\theta]} - \csc^2 \theta = e^{r\theta} \widehat{[dr/d\theta](\theta) + (r)[1]}$  (notice that since  $r$  is a function of  $\theta$  then we must use the Derivative Product Rule to differentiate  $r\theta$ ) or  $-\sin r(dr/d\theta) - \csc^2 \theta = \theta e^{r\theta}(dr/d\theta) + re^{r\theta}$ . Solving for  $dr/d\theta$  we get  $(-\sin r - \theta e^{r\theta}) \frac{dr}{d\theta} = re^{r\theta} + \csc^2 \theta$  or  $\frac{dr}{d\theta} = \frac{re^{r\theta} + \csc^2 \theta}{-\sin r - \theta e^{r\theta}}$ .  $\square$

## Exercise 3.7.20

**Exercise 3.7.20.** Find  $dr/d\theta$  for  $r$  an implicit function of  $\theta$  given by the equation  $\cos r + \cot \theta = e^{r\theta}$ .

**Solution.** Differentiating implicitly,  $\frac{d}{d\theta}[\cos r + \cot \theta] = \frac{d}{d\theta}[e^{r\theta}]$  or

$-\sin r \widehat{[dr/d\theta]} - \csc^2 \theta = e^{r\theta} \widehat{[dr/d\theta](\theta) + (r)[1]}$  (notice that since  $r$  is a function of  $\theta$  then we must use the Derivative Product Rule to differentiate  $r\theta$ ) or  $-\sin r(dr/d\theta) - \csc^2 \theta = \theta e^{r\theta}(dr/d\theta) + re^{r\theta}$ . Solving for  $dr/d\theta$  we get  $(-\sin r - \theta e^{r\theta}) \frac{dr}{d\theta} = re^{r\theta} + \csc^2 \theta$  or  $\frac{dr}{d\theta} = \frac{re^{r\theta} + \csc^2 \theta}{-\sin r - \theta e^{r\theta}}$ .  $\square$

## Exercise 3.7.40

**Exercise 3.7.40.** Verify that the point  $(\pi/4, \pi/2)$  is on the curve  $x \sin 2y = y \cos 2x$  and find the equations of the lines that are **(a)** tangent and **(b)** normal to the curve at  $(\pi/4, \pi/2)$ .

**Solution.** First, with  $(x, y) = (\pi/4, \pi/2)$  the equation  $x \sin 2y = y \cos 2x$  becomes  $\left(\frac{\pi}{4}\right) \sin\left(2\left(\frac{\pi}{2}\right)\right) \stackrel{?}{=} \left(\frac{\pi}{2}\right) \cos\left(2\left(\frac{\pi}{4}\right)\right)$  or  $\left(\frac{\pi}{4}\right) \sin \pi \stackrel{?}{=} \left(\frac{\pi}{2}\right) \cos\left(\frac{\pi}{2}\right)$  or  $\left(\frac{\pi}{4}\right) (0) \stackrel{?}{=} \left(\frac{\pi}{2}\right) (0)$  or  $0 \stackrel{?}{=} 0$ , which is true and so  $(\pi/4, \pi/2)$  is on the curve  $x \sin 2y = y \cos 2x$ .

## Exercise 3.7.40

**Exercise 3.7.40.** Verify that the point  $(\pi/4, \pi/2)$  is on the curve  $x \sin 2y = y \cos 2x$  and find the equations of the lines that are **(a)** tangent and **(b)** normal to the curve at  $(\pi/4, \pi/2)$ .

**Solution.** First, with  $(x, y) = (\pi/4, \pi/2)$  the equation  $x \sin 2y = y \cos 2x$  becomes  $\left(\frac{\pi}{4}\right) \sin\left(2\left(\frac{\pi}{2}\right)\right) \stackrel{?}{=} \left(\frac{\pi}{2}\right) \cos\left(2\left(\frac{\pi}{4}\right)\right)$  or  $\left(\frac{\pi}{4}\right) \sin \pi \stackrel{?}{=} \left(\frac{\pi}{2}\right) \cos\left(\frac{\pi}{2}\right)$  or  $\left(\frac{\pi}{4}\right) (0) \stackrel{?}{=} \left(\frac{\pi}{2}\right) (0)$  or  $0 \stackrel{?}{=} 0$ , which is true and so  $(\pi/4, \pi/2)$  is on the curve  $x \sin 2y = y \cos 2x$ .

**(a)** We differentiate the equation implicitly to find  $dy/dx$  and we get

$$\frac{d}{dx}[x \sin 2y] = \frac{d}{dx}[y \cos 2x] \text{ or}$$

$$[1](\sin 2y) + (x)[\cos(2y)[2dy/dx]] = [dy/dx](\cos 2x) + (y)[- \sin(2x)[2]]$$

or  $\sin 2y + 2x(\cos 2y)(dy/dx) = (\cos 2x)(dy/dx) - 2y \sin 2x$ .

## Exercise 3.7.40

**Exercise 3.7.40.** Verify that the point  $(\pi/4, \pi/2)$  is on the curve  $x \sin 2y = y \cos 2x$  and find the equations of the lines that are **(a)** tangent and **(b)** normal to the curve at  $(\pi/4, \pi/2)$ .

**Solution.** First, with  $(x, y) = (\pi/4, \pi/2)$  the equation  $x \sin 2y = y \cos 2x$  becomes  $\left(\frac{\pi}{4}\right) \sin\left(2\left(\frac{\pi}{2}\right)\right) \stackrel{?}{=} \left(\frac{\pi}{2}\right) \cos\left(2\left(\frac{\pi}{4}\right)\right)$  or  $\left(\frac{\pi}{4}\right) \sin \pi \stackrel{?}{=} \left(\frac{\pi}{2}\right) \cos\left(\frac{\pi}{2}\right)$  or  $\left(\frac{\pi}{4}\right) (0) \stackrel{?}{=} \left(\frac{\pi}{2}\right) (0)$  or  $0 \stackrel{?}{=} 0$ , which is true and so  $(\pi/4, \pi/2)$  is on the curve  $x \sin 2y = y \cos 2x$ .

**(a)** We differentiate the equation implicitly to find  $dy/dx$  and we get

$$\frac{d}{dx}[x \sin 2y] = \frac{d}{dx}[y \cos 2x] \text{ or}$$

$$[1](\sin 2y) + (x)[\cos(2y)[2dy/dx]] = [dy/dx](\cos 2x) + (y)[- \sin(2x)[2]]$$

or  $\sin 2y + 2x(\cos 2y)(dy/dx) = (\cos 2x)(dy/dx) - 2y \sin 2x$ .



## Exercise 3.7.40 (continued 1)

**Exercise 3.7.40.** Verify that the point  $(\pi/4, \pi/2)$  is on the curve  $x \sin 2y = y \cos 2x$  and find the equations of the lines that are **(a)** tangent and **(b)** normal to the curve at  $(\pi/4, \pi/2)$ .

**Solution (continued).**

...  $\sin 2y + 2x(\cos 2y)(dy/dx) = (\cos 2x)(dy/dx) - 2y \sin 2x$ . Solving for  $dy/dx$  we get  $(2x \cos 2y - \cos 2x)(dy/dx) = -2y \sin 2x - \sin 2y$  or

$$\frac{dy}{dx} = \frac{-2y \sin 2x - \sin 2y}{2x \cos 2y - \cos 2x}.$$

## Exercise 3.7.40 (continued 1)

**Exercise 3.7.40.** Verify that the point  $(\pi/4, \pi/2)$  is on the curve  $x \sin 2y = y \cos 2x$  and find the equations of the lines that are **(a)** tangent and **(b)** normal to the curve at  $(\pi/4, \pi/2)$ .

**Solution (continued).**

...  $\sin 2y + 2x(\cos 2y)(dy/dx) = (\cos 2x)(dy/dx) - 2y \sin 2x$ . Solving for  $dy/dx$  we get  $(2x \cos 2y - \cos 2x)(dy/dx) = -2y \sin 2x - \sin 2y$  or  $\frac{dy}{dx} = \frac{-2y \sin 2x - \sin 2y}{2x \cos 2y - \cos 2x}$ . So at  $(\pi/4, \pi/2)$  we have the slope of the tangent line as

$$m = \left. \frac{dy}{dx} \right|_{(x,y)=(\pi/4,\pi/2)} = \frac{-2(\pi/2) \sin 2(\pi/4) - \sin 2(\pi/2)}{2(\pi/4) \cos 2(\pi/2) - \cos 2(\pi/4)} = \frac{-\pi \sin(\pi/2) - \sin \pi}{(\pi/2) \cos \pi - \cos \pi/2} = \frac{-\pi(1) - (0)}{(\pi/2)(-1) - (0)} = \frac{-\pi}{-\pi/2} = 2.$$

## Exercise 3.7.40 (continued 1)

**Exercise 3.7.40.** Verify that the point  $(\pi/4, \pi/2)$  is on the curve  $x \sin 2y = y \cos 2x$  and find the equations of the lines that are **(a)** tangent and **(b)** normal to the curve at  $(\pi/4, \pi/2)$ .

**Solution (continued).**

...  $\sin 2y + 2x(\cos 2y)(dy/dx) = (\cos 2x)(dy/dx) - 2y \sin 2x$ . Solving for  $dy/dx$  we get  $(2x \cos 2y - \cos 2x)(dy/dx) = -2y \sin 2x - \sin 2y$  or  $\frac{dy}{dx} = \frac{-2y \sin 2x - \sin 2y}{2x \cos 2y - \cos 2x}$ . So at  $(\pi/4, \pi/2)$  we have the slope of the tangent line as

$$m = \left. \frac{dy}{dx} \right|_{(x,y)=(\pi/4,\pi/2)} = \frac{-2(\pi/2) \sin 2(\pi/4) - \sin 2(\pi/2)}{2(\pi/4) \cos 2(\pi/2) - \cos 2(\pi/4)} = \frac{-\pi \sin(\pi/2) - \sin \pi}{(\pi/2) \cos \pi - \cos \pi/2} = \frac{-\pi(1) - (0)}{(\pi/2)(-1) - (0)} = \frac{-\pi}{-\pi/2} = 2.$$

So the equation of the line tangent to the curve at  $(x_1, y_1) = (\pi/4, \pi/2)$  is, by the point-slope formula,  $y - y_1 = m(x - x_1)$  or  $y - (\pi/2) = (2)(x - (\pi/4))$  or  $y = 2x - \pi/2 + \pi/2$  or  $y = 2x$ .

## Exercise 3.7.40 (continued 1)

**Exercise 3.7.40.** Verify that the point  $(\pi/4, \pi/2)$  is on the curve  $x \sin 2y = y \cos 2x$  and find the equations of the lines that are **(a)** tangent and **(b)** normal to the curve at  $(\pi/4, \pi/2)$ .

**Solution (continued).**

...  $\sin 2y + 2x(\cos 2y)(dy/dx) = (\cos 2x)(dy/dx) - 2y \sin 2x$ . Solving for  $dy/dx$  we get  $(2x \cos 2y - \cos 2x)(dy/dx) = -2y \sin 2x - \sin 2y$  or  $\frac{dy}{dx} = \frac{-2y \sin 2x - \sin 2y}{2x \cos 2y - \cos 2x}$ . So at  $(\pi/4, \pi/2)$  we have the slope of the tangent line as

$$m = \left. \frac{dy}{dx} \right|_{(x,y)=(\pi/4,\pi/2)} = \frac{-2(\pi/2) \sin 2(\pi/4) - \sin 2(\pi/2)}{2(\pi/4) \cos 2(\pi/2) - \cos 2(\pi/4)} = \frac{-\pi \sin(\pi/2) - \sin \pi}{(\pi/2) \cos \pi - \cos \pi/2} = \frac{-\pi(1) - (0)}{(\pi/2)(-1) - (0)} = \frac{-\pi}{-\pi/2} = 2.$$

So the equation of the line tangent to the curve at  $(x_1, y_1) = (\pi/4, \pi/2)$  is, by the point-slope formula,  $y - y_1 = m(x - x_1)$  or  $y - (\pi/2) = (2)(x - (\pi/4))$  or  $y = 2x - \pi/2 + \pi/2$  or  $y = 2x$ .

## Exercise 3.7.40 (continued 2)

**Exercise 3.7.40.** Verify that the point  $(\pi/4, \pi/2)$  is on the curve  $x \sin 2y = y \cos 2x$  and find the equations of the lines that are **(a)** tangent and **(b)** normal to the curve at  $(\pi/4, \pi/2)$ .

**Solution (continued).** Since the slope of a tangent line to the curve at  $(x_1, y_1) = (\pi/4, \pi/2)$  is 2, then the slope of a line perpendicular (or normal) to the curve at this point is the negative reciprocal of 2, and normal to the curve has slope  $m = -1/2$ . So by the point-slope formula,  $y - y_1 = m(x - x_1)$ , the normal line has formula  $y - (\pi/2) = (-1/2)(x - (\pi/4))$  or  $y = -x/2 + \pi/8 + \pi/2$  or

$$y = -x/2 + 5\pi/8. \quad \square$$

## Exercise 3.7.44

### Exercise 3.7.44. Normals Parallel to a Line.

Find the normals to the curve  $xy + 2x - y = 0$  that are parallel to the line  $2x + y = 0$ .

**Solution.** First we find  $dy/dx$  by differentiating implicitly to get:

$$\frac{d}{dx}[xy + 2x - y] = \frac{d}{dx}[0] \text{ or } [1](y) + (x)[dy/dx] + 2[1] - [dy/dx] = 0 \text{ or}$$

$$y + 2 + (x - 1)dy/dx = 0 \text{ or } \frac{dy}{dx} = \frac{-y - 2}{x - 1} = \frac{y + 2}{1 - x}.$$

So at a point  $(x_1, y_1)$  on the curve  $xy + 2x - y = 0$ , the slope of a normal line is

$$-1 / \left( \frac{dy}{dx} \right) \Big|_{(x_1, y_1)} = -\frac{1 - x_1}{y_1 + 2}.$$

## Exercise 3.7.44

### Exercise 3.7.44. Normals Parallel to a Line.

Find the normals to the curve  $xy + 2x - y = 0$  that are parallel to the line  $2x + y = 0$ .

**Solution.** First we find  $dy/dx$  by differentiating implicitly to get:

$$\frac{d}{dx}[xy + 2x - y] = \frac{d}{dx}[0] \text{ or } [1](y) + (x)[dy/dx] + 2[1] - [dy/dx] = 0 \text{ or}$$

$$y + 2 + (x - 1)dy/dx = 0 \text{ or } \frac{dy}{dx} = \frac{-y - 2}{x - 1} = \frac{y + 2}{1 - x}.$$

So at a point  $(x_1, y_1)$  on the curve  $xy + 2x - y = 0$ , the slope of a normal line is

$$-1 / \left( \frac{dy}{dx} \right) \Big|_{(x_1, y_1)} = -\frac{1 - x_1}{y_1 + 2}.$$

Now the slope of line  $2x + y = 0$  is  $m = -2$  (“ $-A/B$ ,” if you like). So we look for a point  $(x_1, y_1)$  such that

$$-\frac{1 - x_1}{y_1 + 2} = -2 \text{ or } 1 - x_1 = 2(y_1 + 2) = 2y_1 + 4 \text{ or } x_1 = -2y_1 - 3.$$

## Exercise 3.7.44

### Exercise 3.7.44. Normals Parallel to a Line.

Find the normals to the curve  $xy + 2x - y = 0$  that are parallel to the line  $2x + y = 0$ .

**Solution.** First we find  $dy/dx$  by differentiating implicitly to get:

$$\frac{d}{dx}[xy + 2x - y] = \frac{d}{dx}[0] \text{ or } [1](y) + (x)[dy/dx] + 2[1] - [dy/dx] = 0 \text{ or}$$

$$y + 2 + (x - 1)dy/dx = 0 \text{ or } \frac{dy}{dx} = \frac{-y - 2}{x - 1} = \frac{y + 2}{1 - x}.$$

So at a point  $(x_1, y_1)$  on the curve  $xy + 2x - y = 0$ , the slope of a normal line is

$$-1 / \left( \frac{dy}{dx} \right) \Big|_{(x_1, y_1)} = -\frac{1 - x_1}{y_1 + 2}.$$

Now the slope of line  $2x + y = 0$  is

$$m = -2 \text{ ("}-A/B\text{," if you like). So we look for a point } (x_1, y_1) \text{ such that}$$

$$-\frac{1 - x_1}{y_1 + 2} = -2 \text{ or } 1 - x_1 = 2(y_1 + 2) = 2y_1 + 4 \text{ or } x_1 = -2y_1 - 3.$$



## Exercise 3.7.44 (continued)

### Exercise 3.7.44. Normals Parallel to a Line.

Find the normals to the curve  $xy + 2x - y = 0$  that are parallel to the line  $2x + y = 0$ .

**Solution (continued).** So we look for a point  $(x_1, y_1)$  such that  $\dots x_1 = -2y_1 - 3$ . Since  $(x_1, y_1)$  lies on the curve  $xy + 2x - y = 0$ , then we need  $(-2y_1 - 3)y_1 + 2(-2y_1 - 3) - y_1 = 0$  or  $-2y_1^2 - 3y_1 - 4y_1 - 6 - y_1 = 0$  or  $-2y_1^2 - 8y_1 - 6 = 0$  or  $y_1^2 + 4y_1 + 3 = 0$  or  $(y_1 + 3)(y_1 + 1) = 0$ . So we need  $y_1 = -3$  **or**  $y_1 = -1$  and then (from  $xy + 2x - y = 0$ ) we have  $x_1(-3) + 2x_1 - (-3) = 0$  and so  $x_1 = 3$  **or**  $x_1(-1) + 2x_1 - (-1) = 0$  and so  $x_1 = -1$ , respectively.

## Exercise 3.7.44 (continued)

### Exercise 3.7.44. Normals Parallel to a Line.

Find the normals to the curve  $xy + 2x - y = 0$  that are parallel to the line  $2x + y = 0$ .

**Solution (continued).** So we look for a point  $(x_1, y_1)$  such that  $\dots x_1 = -2y_1 - 3$ . Since  $(x_1, y_1)$  lies on the curve  $xy + 2x - y = 0$ , then we need  $(-2y_1 - 3)y_1 + 2(-2y_1 - 3) - y_1 = 0$  or  $-2y_1^2 - 3y_1 - 4y_1 - 6 - y_1 = 0$  or  $-2y_1^2 - 8y_1 - 6 = 0$  or  $y_1^2 + 4y_1 + 3 = 0$  or  $(y_1 + 3)(y_1 + 1) = 0$ . So we need  $y_1 = -3$  **or**  $y_1 = -1$  and then (from  $xy + 2x - y = 0$ ) we have  $x_1(-3) + 2x_1 - (-3) = 0$  and so  $x_1 = 3$  **or**  $x_1(-1) + 2x_1 - (-1) = 0$  and so  $x_1 = -1$ , respectively. That is, the desired normal lines occur at the points  $(3, -3)$  and  $(-1, -1)$ . Since the slope of the normal line is  $m = -2$  then by the point-slope formula the two normal lines are  $y - (-3) = (-2)(x - 3)$  or  $y = -2x + 3$  and  $y - (-1) = (-2)(x - (-1))$  or  $y = -2x - 3$ .  $\square$

## Exercise 3.7.44 (continued)

### Exercise 3.7.44. Normals Parallel to a Line.

Find the normals to the curve  $xy + 2x - y = 0$  that are parallel to the line  $2x + y = 0$ .

**Solution (continued).** So we look for a point  $(x_1, y_1)$  such that  $\dots x_1 = -2y_1 - 3$ . Since  $(x_1, y_1)$  lies on the curve  $xy + 2x - y = 0$ , then we need  $(-2y_1 - 3)y_1 + 2(-2y_1 - 3) - y_1 = 0$  or  $-2y_1^2 - 3y_1 - 4y_1 - 6 - y_1 = 0$  or  $-2y_1^2 - 8y_1 - 6 = 0$  or  $y_1^2 + 4y_1 + 3 = 0$  or  $(y_1 + 3)(y_1 + 1) = 0$ . So we need  $y_1 = -3$  **or**  $y_1 = -1$  and then (from  $xy + 2x - y = 0$ ) we have  $x_1(-3) + 2x_1 - (-3) = 0$  and so  $x_1 = 3$  **or**  $x_1(-1) + 2x_1 - (-1) = 0$  and so  $x_1 = -1$ , respectively. That is, the desired normal lines occur at the points  $(3, -3)$  and  $(-1, -1)$ . Since the slope of the normal line is  $m = -2$  then by the point-slope formula the two normal lines are  $y - (-3) = (-2)(x - 3)$  or  $y = -2x + 3$  and  $y - (-1) = (-2)(x - (-1))$  or  $y = -2x - 3$ .  $\square$

## Exercise 3.7.22

**Exercise 3.7.22.** Find  $dy/dx$  and  $d^2y/dx^2$  for  $y$  an implicit function of  $x$  given by the equation  $x^{2/3} + y^{2/3} = 1$ .

**Solution.** We differentiate implicitly to get  $\frac{d}{dx}[x^{2/3} + y^{2/3}] = \frac{d}{dx}[1]$  or

$$\frac{2}{3}x^{-1/3} + \frac{2}{3}y^{-1/3} \left[ \frac{dy}{dx} \right] = 0 \text{ or } \frac{2}{3}y^{-1/3} \frac{dy}{dx} = -\frac{2}{3}x^{-1/3} \text{ or } \frac{dy}{dx} = -\frac{x^{-1/3}}{y^{-1/3}}$$

or  $\boxed{\frac{dy}{dx} = -\frac{y^{1/3}}{x^{1/3}}}$ .

## Exercise 3.7.22

**Exercise 3.7.22.** Find  $dy/dx$  and  $d^2y/dx^2$  for  $y$  an implicit function of  $x$  given by the equation  $x^{2/3} + y^{2/3} = 1$ .

**Solution.** We differentiate implicitly to get  $\frac{d}{dx}[x^{2/3} + y^{2/3}] = \frac{d}{dx}[1]$  or

$$\frac{2}{3}x^{-1/3} + \frac{2}{3}y^{-1/3} \left[ \frac{dy}{dx} \right] = 0 \text{ or } \frac{2}{3}y^{-1/3} \frac{dy}{dx} = -\frac{2}{3}x^{-1/3} \text{ or } \frac{dy}{dx} = -\frac{x^{-1/3}}{y^{-1/3}}$$

or  $\boxed{\frac{dy}{dx} = -\frac{y^{1/3}}{x^{1/3}}}$ .

Next we have, differentiating implicitly again, that

$$\frac{d}{dx} \left[ \frac{dy}{dx} \right] = \frac{d}{dx} \left[ -\frac{y^{1/3}}{x^{1/3}} \right] \text{ or}$$

$$\frac{d^2y}{dx^2} = -\frac{[(1/3)y^{-2/3}][dy/dx](x^{1/3}) - (y^{1/3})[(1/3)x^{-2/3}]}{(x^{1/3})^2} \text{ or } \dots$$

## Exercise 3.7.22

**Exercise 3.7.22.** Find  $dy/dx$  and  $d^2y/dx^2$  for  $y$  an implicit function of  $x$  given by the equation  $x^{2/3} + y^{2/3} = 1$ .

**Solution.** We differentiate implicitly to get  $\frac{d}{dx}[x^{2/3} + y^{2/3}] = \frac{d}{dx}[1]$  or

$$\frac{2}{3}x^{-1/3} + \frac{2}{3}y^{-1/3} \left[ \frac{dy}{dx} \right] = 0 \text{ or } \frac{2}{3}y^{-1/3} \frac{dy}{dx} = -\frac{2}{3}x^{-1/3} \text{ or } \frac{dy}{dx} = -\frac{x^{-1/3}}{y^{-1/3}}$$

or  $\boxed{\frac{dy}{dx} = -\frac{y^{1/3}}{x^{1/3}}}$ .

Next we have, differentiating implicitly again, that

$$\frac{d}{dx} \left[ \frac{dy}{dx} \right] = \frac{d}{dx} \left[ -\frac{y^{1/3}}{x^{1/3}} \right] \text{ or}$$

$$\frac{d^2y}{dx^2} = -\frac{[(1/3)y^{-2/3}][dy/dx](x^{1/3}) - (y^{1/3})[(1/3)x^{-2/3}]}{(x^{1/3})^2} \text{ or } \dots$$

# Exercise 3.7.22 (continued)

**Exercise 3.7.22.** Find  $dy/dx$  and  $d^2y/dx^2$  for  $y$  an implicit function of  $x$  given by the equation  $x^{2/3} + y^{2/3} = 1$ .

**Solution.** ...  $\frac{d^2y}{dx^2} = -\frac{[(1/3)y^{-2/3}[dy/dx]](x^{1/3}) - (y^{1/3})[(1/3)x^{-2/3}]}{(x^{1/3})^2}$  or

$$\frac{d^2y}{dx^2} = -\frac{(1/3)x^{1/3}y^{-2/3}(dy/dx) - (1/3)x^{-2/3}y^{1/3}}{x^{2/3}}$$
 or

$$\frac{d^2y}{dx^2} = -\frac{x^{1/3}y^{-2/3}(-y^{1/3}/x^{1/3}) - x^{-2/3}y^{1/3}}{3x^{2/3}}$$
 or

$$\frac{d^2y}{dx^2} = \frac{y^{-1/3} + x^{-2/3}y^{1/3}}{3x^{2/3}}. \quad \square$$

## Exercise 3.7.22 (continued)

**Exercise 3.7.22.** Find  $dy/dx$  and  $d^2y/dx^2$  for  $y$  an implicit function of  $x$  given by the equation  $x^{2/3} + y^{2/3} = 1$ .

**Solution.** ...  $\frac{d^2y}{dx^2} = -\frac{[(1/3)y^{-2/3}[dy/dx]](x^{1/3}) - (y^{1/3})[(1/3)x^{-2/3}]}{(x^{1/3})^2}$  or

$$\frac{d^2y}{dx^2} = -\frac{(1/3)x^{1/3}y^{-2/3}(dy/dx) - (1/3)x^{-2/3}y^{1/3}}{x^{2/3}}$$
 or

$$\frac{d^2y}{dx^2} = -\frac{x^{1/3}y^{-2/3}(-y^{1/3}/x^{1/3}) - x^{-2/3}y^{1/3}}{3x^{2/3}}$$
 or

$$\frac{d^2y}{dx^2} = \frac{y^{-1/3} + x^{-2/3}y^{1/3}}{3x^{2/3}}. \quad \square$$



# Exercise 3.7.48

## Exercise 3.7.48. The Folium of Descartes.

- (a) Find the slope of the folium of Descartes  $x^3 + y^3 - 9xy = 0$  at the points  $(4, 2)$  and  $(2, 4)$ . (b) At what point other than the origin does the folium have a horizontal tangent? (c) Find the coordinates of the point  $A$  in Figure 3.29 where the folium has a vertical tangent.




Figure 3.29

**Solution.** (a) Differentiating implicitly we have

$$\frac{d}{dx}[x^3 + y^3 - 9xy] = \frac{d}{dx}[0] \text{ or}$$

$$3x^2 + 3y^2 \left[ \frac{dy}{dx} \right] - 9 \left[ [1](y) + (x) \left[ \frac{dy}{dx} \right] \right] = 0 \text{ or } (3y^2 - 9x) \frac{dy}{dx} = 9y - 3x^2$$

...

# Exercise 3.7.48

## Exercise 3.7.48. The Folium of Descartes.

- (a) Find the slope of the folium of Descartes  $x^3 + y^3 - 9xy = 0$  at the points  $(4, 2)$  and  $(2, 4)$ . (b) At what point other than the origin does the folium have a horizontal tangent? (c) Find the coordinates of the point  $A$  in Figure 3.29 where the folium has a vertical tangent.




Figure 3.29

**Solution.** (a) Differentiating implicitly we have

$$\frac{d}{dx}[x^3 + y^3 - 9xy] = \frac{d}{dx}[0] \text{ or}$$

$$3x^2 + 3y^2 \left[ \frac{dy}{dx} \right] - 9 \left[ [1](y) + (x) \left[ \frac{dy}{dx} \right] \right] = 0 \text{ or } (3y^2 - 9x) \frac{dy}{dx} = 9y - 3x^2$$

...

## Exercise 3.7.48 (continued 1)

(a) Find the slope of the folium of Descartes  $x^3 + y^3 - 9xy = 0$  at the points  $(4, 2)$  and  $(2, 4)$ .

**Solution (continued).** ...  $(3y^2 - 9x) \frac{dy}{dx} = 9y - 3x^2$  or  $\frac{dy}{dx} = \frac{9y - 3x^2}{3y^2 - 9x}$

if  $y^2 \neq 3x$ . So the slope of the curve at  $(4, 2)$  is

$$\left. \frac{dy}{dx} \right|_{(x,y)=(4,2)} = \frac{9(2) - 3(4)^2}{3(2)^2 - 9(4)} = \frac{18 - 48}{12 - 36} = \frac{-30}{-24} = \boxed{\frac{5}{4}}. \text{ The slope of the}$$

curve at  $(4, 2)$  is  $\left. \frac{dy}{dx} \right|_{(x,y)=(2,4)} = \frac{9(4) - 3(2)^2}{3(4)^2 - 9(2)} = \frac{36 - 12}{48 - 18} = \frac{24}{30} = \boxed{\frac{4}{5}}. \square$

# Exercise 3.7.48 (continued 1)

(a) Find the slope of the folium of Descartes  $x^3 + y^3 - 9xy = 0$  at the points  $(4, 2)$  and  $(2, 4)$ .

**Solution (continued).** ...  $(3y^2 - 9x) \frac{dy}{dx} = 9y - 3x^2$  or  $\frac{dy}{dx} = \frac{9y - 3x^2}{3y^2 - 9x}$

if  $y^2 \neq 3x$ . So the slope of the curve at  $(4, 2)$  is

$$\left. \frac{dy}{dx} \right|_{(x,y)=(4,2)} = \frac{9(2) - 3(4)^2}{3(2)^2 - 9(4)} = \frac{18 - 48}{12 - 36} = \frac{-30}{-24} = \boxed{\frac{5}{4}}. \text{ The slope of the}$$

$$\text{curve at } (4, 2) \text{ is } \left. \frac{dy}{dx} \right|_{(x,y)=(2,4)} = \frac{9(4) - 3(2)^2}{3(4)^2 - 9(2)} = \frac{36 - 12}{48 - 18} = \frac{24}{30} = \boxed{\frac{4}{5}}. \quad \square$$

## Exercise 3.7.48 (continued 2)

**(b)** At what point other than the origin does the folium have a horizontal tangent?

**Solution (continued).** **(b)** If the folium has a horizontal tangent at  $(x, y) = (x_1, y_1)$  then

$$\left. \frac{dy}{dx} \right|_{(x,y)=(x_1,y_1)} = \left. \frac{9y - 3x^2}{3y^2 - 9x} \right|_{(x,y)=(x_1,y_1)} = \frac{9y_1 - 3x_1^2}{3y_1^2 - 9x_1} = 0. \text{ This implies}$$

$9y_1 - 3x_1^2 = 0$ , or  $y_1 = x_1^2/3$ . Since  $(x_1, y_1)$  also lies on the folium  $x^3 + y^3 - 9xy = 0$ , then we must have  $x_1^3 + (x_1^2/3)^3 - 9x_1(x_1^2/3) = 0$  or  $x_1^3 + x_1^6/27 - 3x_1^3 = 0$  or  $x_1^6/27 - 2x_1^3 = 0$  or  $x_1^3(x_1^3/27 - 2) = 0$ . So we need either  $x_1 = 0$  or  $x_1 = 3\sqrt[3]{2}$ . When  $x_1 = 0$  then  $y_1 = 0$  and this is the horizontal tangent at the origin. When  $x_1 = 3\sqrt[3]{2}$  we have  $y_1 = (3\sqrt[3]{2})^2/3 = 9(2^{2/3})/3 = 3(\sqrt[3]{2^2}) = 3\sqrt[3]{4}$ . So the other horizontal tangent occurs at  $(x_1, y_1) = (3\sqrt[3]{2}, 3\sqrt[3]{4})$ .  $\square$

## Exercise 3.7.48 (continued 2)

**(b)** At what point other than the origin does the folium have a horizontal tangent?

**Solution (continued).** **(b)** If the folium has a horizontal tangent at  $(x, y) = (x_1, y_1)$  then

$$\left. \frac{dy}{dx} \right|_{(x,y)=(x_1,y_1)} = \left. \frac{9y - 3x^2}{3y^2 - 9x} \right|_{(x,y)=(x_1,y_1)} = \frac{9y_1 - 3x_1^2}{3y_1^2 - 9x_1} = 0. \text{ This implies}$$

$9y_1 - 3x_1^2 = 0$ , or  $y_1 = x_1^2/3$ . Since  $(x_1, y_1)$  also lies on the folium  $x^3 + y^3 - 9xy = 0$ , then we must have  $x_1^3 + (x_1^2/3)^3 - 9x_1(x_1^2/3) = 0$  or  $x_1^3 + x_1^6/27 - 3x_1^3 = 0$  or  $x_1^6/27 - 2x_1^3 = 0$  or  $x_1^3(x_1^3/27 - 2) = 0$ . So we need either  $x_1 = 0$  or  $x_1 = 3\sqrt[3]{2}$ . When  $x_1 = 0$  then  $y_1 = 0$  and this is the horizontal tangent at the origin. When  $x_1 = 3\sqrt[3]{2}$  we have

$y_1 = (3\sqrt[3]{2})^2/3 = 9(2^{2/3})/3 = 3(\sqrt[3]{2^2}) = 3\sqrt[3]{4}$ . So the other horizontal

tangent occurs at  $\boxed{(x_1, y_1) = (3\sqrt[3]{2}, 3\sqrt[3]{4})}$ .  $\square$

## Exercise 3.7.48 (continued 3)

(c) Find the coordinates of the point  $A$  in Figure 3.29 where the folium has a vertical tangent.

**Solution (continued).** (c) Since  $\frac{dy}{dx} = \frac{9y - 3x^2}{3y^2 - 9x}$ , then we look for a vertical tangent where the denominator is 0 and the numerator is not (see Dr. Bob's Infinite Limits Theorem in [2.6. Limits Involving Infinity; Asymptotes of Graphs](#) and the definition of vertical tangent line in [3.1. Tangent Lines and the Derivative at a Point](#)). So if the folium has a vertical tangent at  $(x_1, y_1)$  then  $3y_1^2 - 9x_1 = 0$ , or  $x_1 = y_1^2/3$ . Since  $(x_1, y_1)$  also lies on the folium  $x^3 + y^3 - 9xy = 0$ , then we must have  $(y_1^2/3)^3 + y_1^3 - 9(y_1^2/3)y_1 = 0$  or  $y_1^6/27 + y_1^3 - 3y_1^3 = 0$  or  $y_1^6/27 - 2y_1^3 = 0$  or  $y_1^3(y_1^3/27 - 2) = 0$ . So we need either  $y_1 = 0$  or  $y_1 = 3\sqrt[3]{2}$ . When  $y_1 = 0$  then  $x_1 = 0$  so that  $(x_1, y_1) = (0, 0)$  and we see that the folium has both a horizontal tangent and a vertical tangent at the origin. When  $y_1 = 3\sqrt[3]{2}$  we have  $x_1 = (3\sqrt[3]{2})^2/3 = 3\sqrt[3]{2}^2 = 3\sqrt[3]{4}$ .

## Exercise 3.7.48 (continued 3)

(c) Find the coordinates of the point  $A$  in Figure 3.29 where the folium has a vertical tangent.

**Solution (continued).** (c) Since  $\frac{dy}{dx} = \frac{9y - 3x^2}{3y^2 - 9x}$ , then we look for a vertical tangent where the denominator is 0 and the numerator is not (see Dr. Bob's Infinite Limits Theorem in [2.6. Limits Involving Infinity; Asymptotes of Graphs](#) and the definition of vertical tangent line in [3.1. Tangent Lines and the Derivative at a Point](#)). So if the folium has a vertical tangent at  $(x_1, y_1)$  then  $3y_1^2 - 9x_1 = 0$ , or  $x_1 = y_1^2/3$ . Since  $(x_1, y_1)$  also lies on the folium  $x^3 + y^3 - 9xy = 0$ , then we must have  $(y_1^2/3)^3 + y_1^3 - 9(y_1^2/3)y_1 = 0$  or  $y_1^6/27 + y_1^3 - 3y_1^3 = 0$  or  $y_1^6/27 - 2y_1^3 = 0$  or  $y_1^3(y_1^3/27 - 2) = 0$ . So we need either  $y_1 = 0$  or  $y_1 = 3\sqrt[3]{2}$ . When  $y_1 = 0$  then  $x_1 = 0$  so that  $(x_1, y_1) = (0, 0)$  and we see that the folium has both a horizontal tangent and a vertical tangent at the origin. When  $y_1 = 3\sqrt[3]{2}$  we have  $x_1 = (3\sqrt[3]{2})^2/3 = 3\sqrt[3]{2}^2 = 3\sqrt[3]{4}$ .



## Exercise 3.7.48 (continued 4)

(c) Find the coordinates of the point  $A$  in Figure 3.29 where the folium has a vertical tangent.


**Solution (continued).** We see from the graph that the folium has a vertical tangent at point  $A$  (where the  $x$  coordinate is near 5).

We know that *if* the folium has a vertical tangent then it occurs at  $(x_1, y_1) = (3\sqrt[3]{4}, 3\sqrt[3]{2})$  (or at the origin  $(x_1, y_1) = (0, 0)$ ),

so it must be that point  $A$  is  $(3\sqrt[3]{4}, 3\sqrt[3]{2})$ .

We have  $3\sqrt[3]{4} \approx 4.76$ , consistent with the graph.

Also notice that the horizontal tangent is at  $(3\sqrt[3]{2}, 3\sqrt[3]{4})$  by (b), reflecting the symmetry of the folium with respect to the line  $y = x$ .  $\square$



## Exercise 3.7.48 (continued 4)

(c) Find the coordinates of the point  $A$  in Figure 3.29 where the folium has a vertical tangent.


**Solution (continued).** We see from the graph that the folium has a vertical tangent at point  $A$  (where the  $x$  coordinate is near 5).

We know that *if* the folium has a vertical tangent then it occurs at  $(x_1, y_1) = (3\sqrt[3]{4}, 3\sqrt[3]{2})$  (or at the origin  $(x_1, y_1) = (0, 0)$ ),

so it must be that point  $A$  is  $(3\sqrt[3]{4}, 3\sqrt[3]{2})$ .

We have  $3\sqrt[3]{4} \approx 4.76$ , consistent with the graph.

Also notice that the horizontal tangent is at  $(3\sqrt[3]{2}, 3\sqrt[3]{4})$  by (b), reflecting the symmetry of the folium with respect to the line  $y = x$ .  $\square$



## Exercise 3.7.50

### Exercise 3.7.50. Power Rule for Rational Exponents.

Let  $p$  and  $q$  be integers with  $q > 0$ . If  $y = x^{p/q}$ , differentiate the equivalent equation  $y^q = x^p$  implicitly and show that, for  $y \neq 0$ ,

$$\frac{d}{dx}[x^{p/q}] = \frac{p}{q}x^{(p/q)-1}.$$

**Solution.** Now  $y = x^{p/q}$  if and only if  $y^q = (x^{p/q})^q = x^p$ , so differentiating implicitly we have  $\frac{d}{dx}[y^q] = \frac{d}{dx}[x^p]$  or (since  $q > 0$  and

$$y \neq 0) \quad qy^{q-1} \left[ \frac{dy}{dx} \right] = px^{p-1} \quad \text{or} \quad \frac{dy}{dx} = \frac{px^{p-1}}{qy^{q-1}} \quad \text{or, since } y = x^{p/q},$$

$$\frac{dy}{dx} = \frac{px^{p-1}}{q(x^{p/q})^{q-1}} \quad \text{or} \quad \frac{dy}{dx} = \frac{px^{p-1}}{qx^{p-p/q}} = \frac{p}{q}x^{p-1-(p-p/q)} \quad \text{or}$$

$$\frac{dy}{dx} = \frac{p}{q}x^{(p/q)-1}, \quad \text{as claimed. } \square$$

## Exercise 3.7.50

### Exercise 3.7.50. Power Rule for Rational Exponents.

Let  $p$  and  $q$  be integers with  $q > 0$ . If  $y = x^{p/q}$ , differentiate the equivalent equation  $y^q = x^p$  implicitly and show that, for  $y \neq 0$ ,

$$\frac{d}{dx}[x^{p/q}] = \frac{p}{q}x^{(p/q)-1}.$$

**Solution.** Now  $y = x^{p/q}$  if and only if  $y^q = (x^{p/q})^q = x^p$ , so differentiating implicitly we have  $\frac{d}{dx}[y^q] = \frac{d}{dx}[x^p]$  or (since  $q > 0$  and

$$y \neq 0) \quad qy^{q-1} \left[ \frac{dy}{dx} \right] = px^{p-1} \quad \text{or} \quad \frac{dy}{dx} = \frac{px^{p-1}}{qy^{q-1}} \quad \text{or, since } y = x^{p/q},$$

$$\frac{dy}{dx} = \frac{px^{p-1}}{q(x^{p/q})^{q-1}} \quad \text{or} \quad \frac{dy}{dx} = \frac{px^{p-1}}{qx^{p-p/q}} = \frac{p}{q}x^{p-1-(p-p/q)} \quad \text{or}$$

$$\frac{dy}{dx} = \frac{p}{q}x^{(p/q)-1}, \quad \text{as claimed. } \square$$