



Chapter 7. Transcendental Functions

7.8 Hyperbolic Functions

Note. Recall that $\cos x$ and $\sin x$ are sometimes called the *circular functions*. This is because we can plot a point P on the circle $x^2 + y^2 = 1$ by letting $P = (\cos u, \sin u)$ where u is twice the area of the sector determined by $A = (1, 0)$, $O = (0, 0)$ and P :



Note. We can similarly define the *hyperbolic functions*. Consider the hyperbola $x^2 - y^2 = 1$. Choose a point P on the hyperbola and define u as twice the (signed) area determined by the sector $A = (1, 0)$, $O = (0, 0)$, and P . Now use the coordinates of P to define the hyperbolic trigonometric functions: $P = (\cosh u, \sinh u)$.



Page 545 Figure from Exercise 84

Note. We will use the exponential function to define the hyperbolic trig functions.

Definition. We define

$$\text{Hyperbolic cosine of } x: \cosh x = \frac{e^x + e^{-x}}{2}$$

$$\text{Hyperbolic sine of } x: \sinh x = \frac{e^x - e^{-x}}{2}$$


$$\text{Hyperbolic tangent } x: \tanh x = \frac{\sinh x}{\cosh x} = \frac{e^x - e^{-x}}{e^x + e^{-x}}$$

$$\text{Hyperbolic cotangent of } x: \coth x = \frac{\cosh x}{\sinh x} = \frac{e^x + e^{-x}}{e^x - e^{-x}}$$


$$\text{Hyperbolic secant of } x: \operatorname{sech} x = \frac{1}{\cosh x} = \frac{2}{e^x + e^{-x}}$$

$$\text{Hyperbolic cosecant of } x: \operatorname{csch} x = \frac{1}{\sinh x} = \frac{2}{e^x - e^{-x}}$$


The graphs are:




(a) The hyperbolic sine and its component exponentials.




(b) The hyperbolic cosine and its component exponentials.



(c) The graphs of $y = \tanh x$ and $y = \coth x = 1/\tanh x$.



(d) The graphs of $y = \cosh x$ and $y = \operatorname{sech} x = 1/\cosh x$.



(e) The graphs of $y = \sinh x$ and $y = \operatorname{csch} x = 1/\sinh x$.

Figure 6.26 from the 10th Edition

Note. We have the following identities:

$$\cosh^2 x - \sinh^2 x = 1$$

$$\tanh^2 x = 1 - \operatorname{sech}^2 x$$

$$\operatorname{coth}^2 x = 1 + \operatorname{csch}^2 x$$

$$\sinh 2x = 2 \sinh x \cosh x$$

$$\cosh^2 x = \frac{\cosh 2x + 1}{2}$$

$$\sinh^2 x = \frac{\cosh 2x - 1}{2}$$

Example. Page 542 number 2.

Theorem. (Table 7.7) We have the following differentiation properties:

$$\frac{d}{dx} [\sinh u] = \cosh u \frac{du}{dx}$$

$$\frac{d}{dx} [\cosh u] = \sinh u \frac{du}{dx}$$

$$\frac{d}{dx} [\tanh u] = \operatorname{sech}^2 u \frac{du}{dx}$$

$$\frac{d}{dx} [\operatorname{coth} u] = -\operatorname{csch}^2 u \frac{du}{dx}$$

$$\frac{d}{dx} [\operatorname{sech} u] = -\operatorname{sech} u \tanh u \frac{du}{dx}$$

$$\frac{d}{dx} [\operatorname{csch} u] = -\operatorname{csch} u \operatorname{coth} u \frac{du}{dx}$$

Example. Prove some of the results in the above theorem.

Theorem. (Table 7.8) We have the following integral properties:

$$\int \sinh u \, du = \cosh u + C$$

$$\int \cosh u \, du = \sinh u + C$$

$$\int \operatorname{sech}^2 u \, du = \tanh u + C$$

$$\int \operatorname{csch}^2 u \, du = -\operatorname{coth} u + C$$

$$\int \operatorname{sech} u \tanh u \, du = -\operatorname{sech} u + C$$

$$\int \operatorname{csch} u \operatorname{coth} u \, du = -\operatorname{csch} u + C$$

Proof. These are just the integral versions of the results in Table 6.13.

Q.E.D.

Examples. Page 543 numbers 20 and 50.

Note. Since $\frac{d}{dx}[\sinh x] = \cosh x > 0$, then $\sinh x$ is an INCreasing function and so is one-to-one. The function $\cosh x$ is not one-to-one as we can see from the graph. The function $\operatorname{sech} x = 1/\cosh x$ is also not one-to-one. Therefore, to define the inverse functions of $\cosh x$ and $\operatorname{sech} x$, we must restrict the domains.


Definition. We define some inverse hyperbolic trig functions (Table 7.9).

Define $y = \sinh^{-1} x$ if $x = \sinh y$. (The domain is then $x \in (-\infty, \infty)$.)

Define $y = \cosh^{-1} x$ if $x = \cosh y$ and $y \in [0, \infty)$. (The domain is then $x \in [1, \infty)$.)

Define $y = \operatorname{sech}^{-1} x$ if $x = \operatorname{sech} y$ and $y \in [0, \infty)$. (The domain is then $x \in (0, 1]$.)

Note. The graphs of the above defined three inverse hyperbolic trig functions are:




Page 539 Figure 7.32

Note/Definition. The hyperbolic tangent, cotangent, and cosecant are one-to-one on their domains and therefore have inverses, denoted by

$$y = \tanh^{-1} x, \quad y = \coth^{-1} x, \quad y = \operatorname{csch}^{-1} x.$$

The graphs of these functions are:



Page 539 Figure 7.33

Theorem. We can express the inverse hyperbolic trig functions in terms of the natural logarithm function as follows:

$$\sinh^{-1} x = \ln(x + \sqrt{x^2 + 1}), \quad x \in (-\infty, \infty).$$

$$\cosh^{-1} x = \ln(x + \sqrt{x^2 - 1}), \quad x \in [1, \infty).$$

$$\tanh^{-1} x = \frac{1}{2} \ln \frac{1+x}{1-x}, \quad x \in (-1, 1).$$

$$\operatorname{sech}^{-1} x = \ln \left(\frac{1 + \sqrt{1 - x^2}}{x} \right), \quad x \in (0, 1].$$

$$\operatorname{csch}^{-1} x = \ln \left(\frac{1}{x} + \frac{\sqrt{1 + x^2}}{|x|} \right), \quad x \in (-\infty, 0) \cup (0, \infty).$$

$$\coth^{-1} x = \frac{1}{2} \ln \frac{x+1}{x-1}, \quad x \in (-\infty, -1) \cup (1, \infty).$$

Note. (Table 7.9) We can verify the following identities:

$$\operatorname{sech}^{-1} x = \cosh^{-1} \frac{1}{x}$$

$$\operatorname{csch}^{-1} x = \sinh^{-1} \frac{1}{x}$$

$$\coth^{-1} x = \tanh^{-1} \frac{1}{x}$$

Theorem. (Table 7.10) The inverse hyperbolic trig functions are differentiated as follows:

$$\frac{d}{dx} [\sinh^{-1} u] = \frac{1}{\sqrt{1+u^2}} \frac{du}{dx}$$

$$\frac{d}{dx} [\cosh^{-1} u] = \frac{1}{\sqrt{u^2-1}} \frac{du}{dx}, \quad u \in (1, \infty)$$

$$\frac{d}{dx} [\tanh^{-1} u] = \frac{1}{\sqrt{1-u^2}} \frac{du}{dx}, \quad u \in (-1, 1)$$

$$\frac{d}{dx} [\coth^{-1} u] = \frac{1}{\sqrt{1-u^2}} \frac{du}{dx}, \quad u \in (-\infty, -1) \cup (1, \infty)$$

$$\frac{d}{dx} [\operatorname{sech}^{-1} u] = \frac{-1}{u\sqrt{1-u^2}} \frac{du}{dx}, \quad u \in (0, 1)$$

$$\frac{d}{dx} [\operatorname{csch}^{-1} u] = \frac{-1}{|u|\sqrt{1+u^2}} \frac{du}{dx}, \quad u \in (-\infty, 0) \cup (0, \infty)$$

Theorem. (Table 7.11) We have the following integrals involving inverse hyperbolic trig functions:

$$\int \frac{du}{\sqrt{a^2 + u^2}} = \sinh^{-1} \left(\frac{u}{a} \right) + C, \quad a > 0$$

$$\int \frac{du}{\sqrt{a^2 - u^2}} = \cosh^{-1} \left(\frac{u}{a} \right) + C, \quad u > a > 0$$

$$\int \frac{du}{a^2 - u^2} = \begin{cases} \frac{1}{a} \tanh^{-1} \left(\frac{u}{a} \right) + C & \text{if } u^2 < a^2 \\ \frac{1}{a} \coth^{-1} \left(\frac{u}{a} \right) + C & \text{if } u^2 > a^2 \end{cases}$$

$$\int \frac{du}{u\sqrt{a^2 - u^2}} = -\frac{1}{a} \operatorname{sech}^{-1} \left(\frac{u}{a} \right) + C, \quad 0 < u < a$$

$$\int \frac{du}{u\sqrt{a^2 + u^2}} = -\frac{a}{u} \operatorname{csch}^{-1} \left| \frac{u}{a} \right| + C, \quad u \neq 0$$

Examples. Page 544 numbers 70 and 79.

Corrected 1/18/2020