

# Chapter 9. Further Applications of Integration

## 9.4 Graphical Solutions of Autonomous Differential Equations

**Definition.** An equation of the form  $\frac{dy}{dx} = g(y)$  is an autonomous ordinary differential equation.

**Definition.** If  $\frac{dy}{dx} = g(y)$  is an autonomous differential equation, then the values of  $y$  for which  $\frac{dy}{dx} = 0$  are called *equilibrium values* or *rest points*.

**Example.** Page 671 number 2a.

**Note.** We make use of a *phase line* for these types of differential equations. This is a plot on the  $y$ -axis that shows the equation's equilibrium values along with the intervals where  $dy/dx$  and  $d^2y/dx^2$  are positive and negative. Then we know where the solutions are increasing and decreasing and the concavity of the solution curves.

**Example.** Page 671 number 2b.

**Note.** An equilibrium value is *stable* if we perturb the system slightly and it returns to the equilibrium value. For example, a ball at the bottom of a well is in a stable equilibrium state — perturb the ball slightly and it returns to where it was. An equilibrium value is *unstable* if a small perturbation of the system may yield solutions that do not return to the equilibrium value. For example, a ball at the top of a hill is in an unstable equilibrium — perturb the ball a little and it rolls away.

**Example.** Page 671 number 2a (continued), 2c.

**Example.** Page 670 Example 5. Consider the logistic equation

$$\frac{dP}{dt} = r(M - P)P.$$