

Introduction to Modern Algebra

Part III. Homomorphisms and Factor Groups

III.15. Factor-Group Computations and Simple Groups




Table of contents

- 1 Lemma
- 2 Example 15.6. Falsity of the Converse of the Theorem of Lagrange
- 3 Exercise 15.6
- 4 Theorem 15.8
- 5 Theorem 15.9
- 6 Exercise 15.4
- 7 Theorem 15.18
- 8 Theorem 15.20

Lemma

Lemma. If G is a finite group and N is a subgroup of G where $|N| = |G|/2$, then N is a normal subgroup of G .

Proof. Since all cosets of N must be the same size and the cosets partition G , then there are only two cosets of N , namely N and aN where $a \in G \setminus N$. Now for any $g \in G$, (1) if $g \in N$ then $gN = Ng = N$, and (2) if $g \in G \setminus N$ then $gN = Ng$ since this is the only coset of N other than N itself. So $gN = Ng$ for all $g \in G$, and N is a normal subgroup. \square

Lemma

Lemma. If G is a finite group and N is a subgroup of G where $|N| = |G|/2$, then N is a normal subgroup of G .

Proof. Since all cosets of N must be the same size and the cosets partition G , then there are only two cosets of N , namely N and aN where $a \in G \setminus N$. Now for any $g \in G$, (1) if $g \in N$ then $gN = Ng = N$, and (2) if $g \in G \setminus N$ then $gN = Ng$ since this is the only coset of N other than N itself. So $gN = Ng$ for all $g \in G$, and N is a normal subgroup. \square

Example 15.6

Example 15.6. Falsity of the Converse of the Theorem of Lagrange.

We have claimed in the past that the alternating group A_4 (of order $4!/2 = 12$) does not have a subgroup of order 6. Recall that Lagrange's Theorem states that the order of a subgroup divides the order of its group. This example shows that there may be divisors of the order of a group which may not be the order of a subgroup.

Proof. Suppose, to the contrary, that H is a subgroup of A_4 of order 6. By Lemma, H must be a normal subgroup of G . Then A_4/H has only two elements, H and σH where $\sigma \in A_n \setminus H$. Since A_4/H is a group of order 2, then it is isomorphic to \mathbb{Z}_2 and the square of each element (coset) is the identity (H). So $H \cdot H = H$ and $(\sigma H) \cdot (\sigma H) = \sigma^2 H = H$. So if $\alpha \in H$ then $\alpha^2 \in H$ and if $\beta \notin H$ (then $\beta \in \sigma H$) then $\beta^2 \in H$. So, the square of every element of A_4 is in H .

Example 15.6

Example 15.6. Falsity of the Converse of the Theorem of Lagrange.

We have claimed in the past that the alternating group A_4 (of order $4!/2 = 12$) does not have a subgroup of order 6. Recall that Lagrange's Theorem states that the order of a subgroup divides the order of its group. This example shows that there may be divisors of the order of a group which may not be the order of a subgroup.

Proof. Suppose, to the contrary, that H is a subgroup of A_4 of order 6. By Lemma, H must be a normal subgroup of G . Then A_4/H has only two elements, H and σH where $\sigma \in A_n \setminus H$. Since A_4/H is a group of order 2, then it is isomorphic to \mathbb{Z}_2 and the square of each element (coset) is the identity (H). So $H \cdot H = H$ and $(\sigma H) \cdot (\sigma H) = \sigma^2 H = H$. So if $\alpha \in H$ then $\alpha^2 \in H$ and if $\beta \notin H$ (then $\beta \in \sigma H$) then $\beta^2 \in H$. So, the square of every element of A_4 is in H .

Example 15.6 (continued)

Example 15.6. Falsity of the Converse of the Theorem of Lagrange.

We have claimed in the past that the alternating group A_4 (of order $4!/2 = 12$) does not have a subgroup of order 6. Recall that Lagrange's Theorem states that the order of a subgroup divides the order of its group. This example shows that there may be divisors of the order of a group which may not be the order of a subgroup.

Proof (continued). But in A_4 we have

$$(1, 2, 3) = (1, 3, 2)^2 \text{ and } (1, 3, 2) = (1, 2, 3)^2$$

$$(1, 2, 4) = (1, 4, 2)^2 \text{ and } (1, 4, 2) = (1, 2, 4)^2$$

$$(1, 3, 4) = (1, 4, 3)^2 \text{ and } (1, 4, 3) = (1, 3, 4)^2$$

$$(2, 3, 4) = (2, 4, 3)^2 \text{ and } (2, 4, 3) = (2, 3, 4)^2.$$

So all 8 of the above (distinct) permutations are in H . This is a contradiction, since we assumed $|H| = 6$. Therefore no such H exists. \square

Exercise 15.6

Exercise 15.6. Classify $\mathbb{Z} \times \mathbb{Z}/\langle(0, 1)\rangle$ according to the Fundamental Theorem of Finitely Generated Abelian Groups.

Solution. Notice $\langle(0, 1)\rangle = \{0\} \times \mathbb{Z} = H$ and the cosets are $(x, y) + \{0\} \times \mathbb{Z} = \{x\} \times (\mathbb{Z} + y) = \{x\} \times \mathbb{Z}$ for all $(x, y) \in \mathbb{Z} \times \mathbb{Z}$. Now $\{x\} \times \mathbb{Z}$ is isomorphic to \mathbb{Z} (define $\varphi(\{x\} \times \mathbb{Z}) = x$ and φ is an isomorphism). So $\mathbb{Z} \times \mathbb{Z}/\langle(0, 1)\rangle$ is isomorphic to \mathbb{Z} . □

Exercise 15.6

Exercise 15.6. Classify $\mathbb{Z} \times \mathbb{Z} / \langle (0, 1) \rangle$ according to the Fundamental Theorem of Finitely Generated Abelian Groups.

Solution. Notice $\langle (0, 1) \rangle = \{0\} \times \mathbb{Z} = H$ and the cosets are $(x, y) + \{0\} \times \mathbb{Z} = \{x\} \times (\mathbb{Z} + y) = \{x\} \times \mathbb{Z}$ for all $(x, y) \in \mathbb{Z} \times \mathbb{Z}$. Now $\{x\} \times \mathbb{Z}$ is isomorphic to \mathbb{Z} (define $\varphi(\{x\} \times \mathbb{Z}) = x$ and φ is an isomorphism). So $\mathbb{Z} \times \mathbb{Z} / \langle (0, 1) \rangle$ is isomorphic to \mathbb{Z} . □

Theorem 15.8

Theorem 15.8. Let $G = H \times K$ be the direct product of groups H and K . Then $H = \{(h, e) \mid h \in H\}$ is a normal subgroup of G . Also, G/\tilde{H} is isomorphic to K . Similarly G/\tilde{K} is isomorphic to H where $\tilde{K} = \{(e, k) \mid k \in K\}$.

Proof. Define $\pi_2 : H \times K \rightarrow K$ where $\pi_2(h, k) = k$. Then π_2 is a projection map (see Example 13.8) and so is a homomorphism. Now, $\text{Ker}(\pi_2) = \tilde{H}$ and so \tilde{H} is a normal subgroup of $H \times K$ by Theorem 13.15. Since π_2 is onto K then by Theorem 14.11, $(H \times K)/\tilde{H} \cong K$ (here, $H \times K$ plays the role of G , and K plays the role of $\varphi[G]$ in Theorem 14.11). \square

Theorem 15.8

Theorem 15.8. Let $G = H \times K$ be the direct product of groups H and K . Then $H = \{(h, e) \mid h \in H\}$ is a normal subgroup of G . Also, G/\tilde{H} is isomorphic to K . Similarly G/\tilde{K} is isomorphic to H where $\tilde{K} = \{(e, k) \mid k \in K\}$.

Proof. Define $\pi_2 : H \times K \rightarrow K$ where $\pi_2(h, k) = k$. Then π_2 is a projection map (see Example 13.8) and so is a homomorphism. Now, $\text{Ker}(\pi_2) = \tilde{H}$ and so \tilde{H} is a normal subgroup of $H \times K$ by Theorem 13.15. Since π_2 is onto K then by Theorem 14.11, $(H \times K)/\tilde{H} \cong K$ (here, $H \times K$ plays the role of G , and K plays the role of $\varphi[G]$ in Theorem 14.11). \square

Theorem 15.9

Theorem 15.9. A factor group of a cyclic group is cyclic.

Proof. Let G be cyclic where $\langle a \rangle = G$. Let N be a normal subgroups of G . Next, G/N consists of all cosets of N . Let $gN \in G/N$ where $g \in G$. Since a generates G , $g = a^n$ for some $n \in \mathbb{Z}$, and $gN = (aN)^n = a^nN$. Therefore $G/N = \langle aN \rangle$ and G/N is cyclic, as claimed. \square

Theorem 15.9

Theorem 15.9. A factor group of a cyclic group is cyclic.

Proof. Let G be cyclic where $\langle a \rangle = G$. Let N be a normal subgroups of G . Next, G/N consists of all cosets of N . Let $gN \in G/N$ where $g \in G$. Since a generates G , $g = a^n$ for some $n \in \mathbb{Z}$, and $gN = (aN)^n = a^nN$. Therefore $G/N = \langle aN \rangle$ and G/N is cyclic, as claimed. \square

Exercise 15.4

Exercise 15.4. Classify $(\mathbb{Z}_4 \times \mathbb{Z}_8)/\langle(1, 2)\rangle$.

Solution. Notice $\langle(1, 2)\rangle = \{(1, 2), (2, 4), (3, 6), (0, 0)\} = H$ and so $|\mathbb{Z}_4 \times \mathbb{Z}_8/\langle(1, 2)\rangle| = 4 \times 8/4 = 8$. All groups are abelian, including the factor group. The abelian groups of order 8 are $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$, $\mathbb{Z}_2 \times \mathbb{Z}_4$, and \mathbb{Z}_8 (we don't distinguish between $\mathbb{Z}_2 \times \mathbb{Z}_4$ and $\mathbb{Z}_4 \times \mathbb{Z}_2$). We try to determine which of these three choices is correct by considering orders of elements of the factor group.

Exercise 15.4

Exercise 15.4. Classify $(\mathbb{Z}_4 \times \mathbb{Z}_8)/\langle(1, 2)\rangle$.

Solution. Notice $\langle(1, 2)\rangle = \{(1, 2), (2, 4), (3, 6), (0, 0)\} = H$ and so $|\mathbb{Z}_4 \times \mathbb{Z}_8/\langle(1, 2)\rangle| = 4 \times 8/4 = 8$. All groups are abelian, including the factor group. The abelian groups of order 8 are $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$, $\mathbb{Z}_2 \times \mathbb{Z}_4$, and \mathbb{Z}_8 (we don't distinguish between $\mathbb{Z}_2 \times \mathbb{Z}_4$ and $\mathbb{Z}_4 \times \mathbb{Z}_2$). We try to determine which of these three choices is correct by considering orders of elements of the factor group. Consider the coset $(0, 1) + H$. The order of this coset is 8 since:

$$\underbrace{((0, 1) + H) + ((0, 1) + H) + \cdots + ((0, 1) + H)}_{k \text{ times}} = (0, k) + H$$

and the smallest value of k for which this yields the identity is $k = 8$. Since the only choice from the three groups which has elements of order 8 is \mathbb{Z}_8 , then the factor group must be isomorphic to \mathbb{Z}_8 . \square

Exercise 15.4

Exercise 15.4. Classify $(\mathbb{Z}_4 \times \mathbb{Z}_8)/\langle(1, 2)\rangle$.

Solution. Notice $\langle(1, 2)\rangle = \{(1, 2), (2, 4), (3, 6), (0, 0)\} = H$ and so $|\mathbb{Z}_4 \times \mathbb{Z}_8/\langle(1, 2)\rangle| = 4 \times 8/4 = 8$. All groups are abelian, including the factor group. The abelian groups of order 8 are $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$, $\mathbb{Z}_2 \times \mathbb{Z}_4$, and \mathbb{Z}_8 (we don't distinguish between $\mathbb{Z}_2 \times \mathbb{Z}_4$ and $\mathbb{Z}_4 \times \mathbb{Z}_2$). We try to determine which of these three choices is correct by considering orders of elements of the factor group. Consider the coset $(0, 1) + H$. The order of this coset is 8 since:

$$\underbrace{((0, 1) + H) + ((0, 1) + H) + \cdots + ((0, 1) + H)}_{k \text{ times}} = (0, k) + H$$

and the smallest value of k for which this yields the identity is $k = 8$. Since the only choice from the three groups which has elements of order 8 is \mathbb{Z}_8 , then the factor group must be isomorphic to \mathbb{Z}_8 . \square

Theorem 15.18

Theorem 15.18. M is a maximal normal subgroup of G if and only if G/M is simple.

Proof. Let M be a maximal normal subgroup of G . Define $\gamma : G \rightarrow G/M$ as $\gamma(g) = gM$. Then by Theorem 14.9, γ is a homomorphism with $\text{Ker}(\gamma) = M$. ASSUME G/M is *not* simple and that N' is a proper nontrivial normal subgroup of G/M . Then by Theorem 15.16, $\gamma^{-1}[N']$ is a normal subgroup of G .

Theorem 15.18

Theorem 15.18. M is a maximal normal subgroup of G if and only if G/M is simple.

Proof. Let M be a maximal normal subgroup of G . Define $\gamma : G \rightarrow G/M$ as $\gamma(g) = gM$. Then by Theorem 14.9, γ is a homomorphism with $\text{Ker}(\gamma) = M$. ASSUME G/M is *not* simple and that N' is a proper nontrivial normal subgroup of G/M . Then by Theorem 15.16, $\gamma^{-1}[N']$ is a normal subgroup of G . Since N' is nontrivial then $N' \neq \{e\} = \{M\}$ (remember M is the identity in G/M). Also, we have $\text{Ker}(\gamma) = M$, so $\text{Ker}(\gamma) \neq N'$ and $\varphi^{-1}[N'] \neq M$. But $\{e\} = \{M\} \subsetneq N'$ and so $M \subsetneq \varphi^{-1}[N']$. So $\varphi^{-1}[N']$ properly contains M .

Theorem 15.18

Theorem 15.18. M is a maximal normal subgroup of G if and only if G/M is simple.

Proof. Let M be a maximal normal subgroup of G . Define $\gamma : G \rightarrow G/M$ as $\gamma(g) = gM$. Then by Theorem 14.9, γ is a homomorphism with $\text{Ker}(\gamma) = M$. ASSUME G/M is *not* simple and that N' is a proper nontrivial normal subgroup of G/M . Then by Theorem 15.16, $\gamma^{-1}[N']$ is a normal subgroup of G . Since N' is nontrivial then $N' \neq \{e\} = \{M\}$ (remember M is the identity in G/M). Also, we have $\text{Ker}(\gamma) = M$, so $\text{Ker}(\gamma) \neq N'$ and $\gamma^{-1}[N'] \neq M$. But $\{e\} = \{M\} \subsetneq N'$ and so $M \subsetneq \gamma^{-1}[N']$. So $\gamma^{-1}[N']$ properly contains M . Since N' is a proper subgroup of G/M , then it contains some but not all of the cosets of M . Since the cosets of M partition G (Section II.10), then $\gamma^{-1}[N']$ contains some but not all elements of G . That is, $\gamma^{-1}[N']$ is a proper subgroup of G . Therefore, $\gamma^{-1}[N']$ is a nontrivial proper normal subgroup of G which properly contains M . But this CONTRADICTS the maximality of M . So no such N' exists and G/M is simple, as claimed.

Theorem 15.18

Theorem 15.18. M is a maximal normal subgroup of G if and only if G/M is simple.

Proof. Let M be a maximal normal subgroup of G . Define $\gamma : G \rightarrow G/M$ as $\gamma(g) = gM$. Then by Theorem 14.9, γ is a homomorphism with $\text{Ker}(\gamma) = M$. ASSUME G/M is *not* simple and that N' is a proper nontrivial normal subgroup of G/M . Then by Theorem 15.16, $\gamma^{-1}[N']$ is a normal subgroup of G . Since N' is nontrivial then $N' \neq \{e\} = \{M\}$ (remember M is the identity in G/M). Also, we have $\text{Ker}(\gamma) = M$, so $\text{Ker}(\gamma) \neq N'$ and $\gamma^{-1}[N'] \neq M$. But $\{e\} = \{M\} \subsetneq N'$ and so $M \subsetneq \gamma^{-1}[N']$. So $\gamma^{-1}[N']$ properly contains M . Since N' is a proper subgroup of G/M , then it contains some but not all of the cosets of M . Since the cosets of M partition G (Section II.10), then $\gamma^{-1}[N']$ contains some but not all elements of G . That is, $\gamma^{-1}[N']$ is a proper subgroup of G . Therefore, $\gamma^{-1}[N']$ is a nontrivial proper normal subgroup of G which properly contains M . But this CONTRADICTS the maximality of M . So no such N' exists and G/M is simple, as claimed.

Theorem 15.18 (continued)

Proof (continued). Now let G/M be simple. ASSUME that M is *not* a maximal normal subgroup and that there is a proper normal subgroup N of G properly containing M (i.e., $M \subsetneq N \subsetneq G$). Then by Theorem 15.16, $\gamma[N]$ is a normal subgroup of $\gamma[G] = G/M$ where γ is as defined above. Now $\text{Ker}(\gamma) = \{M\}$ (remember that M is the identity in G/M). Hence $\gamma[N]$ is a nontrivial normal subgroup of G/M . Now we show that $\gamma[N]$ is a proper subgroup of G/M . Recall that γ maps elements of G to cosets of M . So the only way that $\gamma[N] = G/M$ is if N contains an element of each coset of M . But we assumed $M \subsetneq N$, and so if N contains an element of each coset of M , then (since N is a group and so is closed under the binary operation) N must contain all cosets of M —that is, $M = G$. But this CONTRADICTS the choice of N as a proper subgroup of G .

Theorem 15.18 (continued)

Proof (continued). Now let G/M be simple. ASSUME that M is *not* a maximal normal subgroup and that there is a proper normal subgroup N of G properly containing M (i.e., $M \subsetneq N \subsetneq G$). Then by Theorem 15.16, $\gamma[N]$ is a normal subgroup of $\gamma[G] = G/M$ where γ is as defined above. Now $\text{Ker}(\gamma) = \{M\}$ (remember that M is the identity in G/M). Hence $\gamma[N]$ is a nontrivial normal subgroup of G/M . Now we show that $\gamma[N]$ is a proper subgroup of G/M . Recall that γ maps elements of G to cosets of M . So the only way that $\gamma[N] = G/M$ is if N contains an element of each coset of M . But we assumed $M \subsetneq N$, and so if N contains an element of each coset of M , then (since N is a group and so is closed under the binary operation) N must contain all cosets of M —that is, $M = G$. But this **CONTRADICTS** the choice of N as a proper subgroup of G . Hence N does not contain an element from each coset of M and $\gamma[N] \neq G/M$. That is, $\gamma[N] \subsetneq G/M$. Therefore, $\gamma[N]$ is a proper nontrivial subgroup of G/M . But this contradicts the fact that G/M is simple. Therefore, no such N exists and M is a maximal normal subgroup of G , as claimed. \square

Theorem 15.18 (continued)

Proof (continued). Now let G/M be simple. ASSUME that M is *not* a maximal normal subgroup and that there is a proper normal subgroup N of G properly containing M (i.e., $M \subsetneq N \subsetneq G$). Then by Theorem 15.16, $\gamma[N]$ is a normal subgroup of $\gamma[G] = G/M$ where γ is as defined above. Now $\text{Ker}(\gamma) = \{M\}$ (remember that M is the identity in G/M). Hence $\gamma[N]$ is a nontrivial normal subgroup of G/M . Now we show that $\gamma[N]$ is a proper subgroup of G/M . Recall that γ maps elements of G to cosets of M . So the only way that $\gamma[N] = G/M$ is if N contains an element of each coset of M . But we assumed $M \subsetneq N$, and so if N contains an element of each coset of M , then (since N is a group and so is closed under the binary operation) N must contain all cosets of M —that is, $M = G$. But this CONTRADICTS the choice of N as a proper subgroup of G . Hence N does not contain an element from each coset of M and $\gamma[N] \neq G/M$. That is, $\gamma[N] \subsetneq G/M$. Therefore, $\gamma[N]$ is a proper nontrivial subgroup of G/M . But this contradicts the fact that G/M is simple. Therefore, no such N exists and M is a maximal normal subgroup of G , as claimed. \square

Theorem 15.20

Theorem 15.20. Let G be a group. Then the set $C = \{aba^{-1}b^{-1} \mid a, b \in G\}$ is a subgroup of G . Additionally, C is a normal subgroup of G then G/N is abelian if and only if $C \leq N$.

Proof. Since C is a subset of G , associativity of the binary operation is inherited from G and G_1 holds. Taking $a = e$, we see that $aba^{-1}be \in C$, we see that $aba^{-1}b^{-1} = e \in C$ and G_2 holds. If $x \in C$, then $x = aba^{-1}b^{-1}$ and $x^{-1} = bab^{-1}a^{-1} \in C$ and G_3 holds. Hence C is a group, as claimed.

Theorem 15.20

Theorem 15.20. Let G be a group. Then the set $C = \{aba^{-1}b^{-1} \mid a, b \in G\}$ is a subgroup of G . Additionally, C is a normal subgroup of G then G/N is abelian if and only if $C \leq N$.

Proof. Since C is a subset of G , associativity of the binary operation is inherited from G and G_1 holds. Taking $a = e$, we see that $aba^{-1}be \in C$, we see that $aba^{-1}b^{-1} = e \in C$ and G_2 holds. If $x \in C$, then $x = aba^{-1}b^{-1}$ and $x^{-1} = bab^{-1}a^{-1} \in C$ and G_3 holds. Hence C is a group, as claimed.

For normality, let $x \in C$ and $g \in G$. Then $x = cdc^{-1}d^{-1}$ for some $c, d \in G$. Hence

$$\begin{aligned} g^{-1}xg &= g^{-1}(cdc^{-1}d^{-1})g = g^{-1}(cdc^{-1}ed^{-1})g \\ &= g^{-1}cdc^{-1}(gd^1dg^{-1})d^{-1}g \text{ since } e = gd^{-1}dg^{-1} \\ &= \left[(g^{-1}c) d (g^{-1}c)^{-1} d^{-1} \right] [dg^{-1}d^{-1}g] \dots \end{aligned}$$

Theorem 15.20

Theorem 15.20. Let G be a group. Then the set $C = \{aba^{-1}b^{-1} \mid a, b \in G\}$ is a subgroup of G . Additionally, C is a normal subgroup of G then G/N is abelian if and only if $C \leq N$.

Proof. Since C is a subset of G , associativity of the binary operation is inherited from G and G_1 holds. Taking $a = e$, we see that $aba^{-1}be \in C$, we see that $aba^{-1}b^{-1} = e \in C$ and G_2 holds. If $x \in C$, then $x = aba^{-1}b^{-1}$ and $x^{-1} = bab^{-1}a^{-1} \in C$ and G_3 holds. Hence C is a group, as claimed.

For normality, let $x \in C$ and $g \in G$. Then $x = cdc^{-1}d^{-1}$ for some $c, d \in G$. Hence

$$\begin{aligned} g^{-1}xg &= g^{-1}(cdc^{-1}d^{-1})g = g^{-1}(cdc^{-1}ed^{-1})g \\ &= g^{-1}cdc^{-1}(gd^1dg^{-1})d^{-1}g \text{ since } e = gd^{-1}dg^{-1} \\ &= \left[(g^{-1}c) d (g^{-1}c)^{-1} d^{-1} \right] [dg^{-1}d^{-1}g] \dots \end{aligned}$$

Theorem 15.20 (continued).

Proof (continued). ...

$$g^{-1}xg = \left[(g^{-1}c) d (g^{-1}c)^{-1} d^{-1} \right] [dg^{-1}d^{-1}g] \in C$$

since $(g^{-1}c) d (g^{-1}c)^{-1} d^{-1}$,

so that $dg^{-1}d^{-1}g \in C$. So $g^{-1}xg \in C$ and C is a normal subgroup of G by Theorem 14.13, as claimed.

Now let N be a normal subgroup of G ($N \triangleleft G$). Suppose G/N is abelian.

Then for all $a, b \in G$

$$(a^{-1}b^{-1})N = (a^{-1}N)(b^{-1}N) = (b^{-1}N)(a^{-1}N) = b^{-1}a^{-1}N \text{ or}$$

$$(aba^{-1}b^{-1})N = N \text{ and so } aba^{-1}b^{-1} \in N \text{ and } C \leq N.$$

Theorem 15.20 (continued).

Proof (continued). ...

$$g^{-1}xg = \left[(g^{-1}c) d (g^{-1}c)^{-1} d^{-1} \right] [dg^{-1}d^{-1}g] \in C$$

since $(g^{-1}c) d (g^{-1}c)^{-1} d^{-1}$,

so that $dg^{-1}d^{-1}g \in C$. So $g^{-1}xg \in C$ and C is a normal subgroup of G by Theorem 14.13, as claimed.

Now let N be a normal subgroup of G ($N \triangleleft G$). Suppose G/N is abelian.

Then for all $a, b \in G$

$$(a^{-1}b^{-1})N = (a^{-1}N)(b^{-1}N) = (b^{-1}N)(a^{-1}N) = b^{-1}a^{-1}N \text{ or}$$

$$(aba^{-1}b^{-1})N = N \text{ and so } aba^{-1}b^{-1} \in N \text{ and } C \leq N.$$

Let N be a normal subgroup of G . Suppose $C \leq N$. Then for all $a, b \in G$

$$(aN)(bN) = (ab)N = ab(b^{-1}a^{-1}ba)N, \text{ since}$$

$$b^{-1}a^{-1}ba \in C \subseteq N = baN = (bN)(aN), \text{ and so } G/N \text{ is abelian, as}$$

claimed. □

Theorem 15.20 (continued).

Proof (continued). ...

$$g^{-1}xg = \left[(g^{-1}c) d (g^{-1}c)^{-1} d^{-1} \right] [dg^{-1}d^{-1}g] \in C$$

since $(g^{-1}c) d (g^{-1}c)^{-1} d^{-1}$,

so that $dg^{-1}d^{-1}g \in C$. So $g^{-1}xg \in C$ and C is a normal subgroup of G by Theorem 14.13, as claimed.

Now let N be a normal subgroup of G ($N \triangleleft G$). Suppose G/N is abelian.

Then for all $a, b \in G$

$$(a^{-1}b^{-1})N = (a^{-1}N)(b^{-1}N) = (b^{-1}N)(a^{-1}N) = b^{-1}a^{-1}N \text{ or}$$

$$(aba^{-1}b^{-1})N = N \text{ and so } aba^{-1}b^{-1} \in N \text{ and } C \leq N.$$

Let N be a normal subgroup of G . Suppose $C \leq N$. Then for all $a, b \in G$

$$(aN)(bN) = (ab)N = ab(b^{-1}a^{-1}ba)N, \text{ since}$$

$$b^{-1}a^{-1}ba \in C \subseteq N = baN = (bN)(aN), \text{ and so } G/N \text{ is abelian, as}$$

claimed. □