


# Real Analysis

## Chapter 20. The Construction of Particular Measures

### 20.4. Carathéodory Outer Measure and Hausdorff Measures on a Metric Space—Proofs of Theorems



# Table of contents

- 1 Proposition 20.27
- 2 Proposition 20.28
- 3 Proposition 20.29
- 4 Proposition 20.30
- 5 Theorem 20.4.A

# Proposition 20.27

**Proposition 20.27.** Let  $\varphi$  be a real-valued function on a set  $X$  and  $\mu^* : 2^X \rightarrow [0, \infty]$  an outer measure with the property that whenever two subsets  $A$  and  $B$  of  $X$  are separated by  $\varphi$ , then

$$\mu^*(A \cup B) = \mu^*(A) + \mu^*(B).$$

Then  $\varphi$  is measurable with respect to the measure induced by  $\mu^*$ .

**Proof.** Let  $a \in \mathbb{R}$ . We show that  $E = \{x \in X \mid \varphi(x) > a\}$  is  $\mu^*$ -measurable, implying the measurability of function  $\varphi$ . By definition, an outer measure is countably monotone (see Section 17.3) so

$$\mu^*(A) = \mu^*((A \cap E) \cup (A \cap E^c)) \leq \mu^*(A \cap E) + \mu^*(A \cap E^c).$$

So we need only show that for  $A$  a set of finite measure and for any  $\varepsilon > 0$ ,

$$\mu^*(A) + \varepsilon > \mu^*(A \cap E) + \mu^*(A \cap E^c). \quad (32)$$

Notice that (32) trivially holds if  $\mu^*(A) = \infty$ , so we can assume  $\mu^*(A) < \infty$ . Define  $B = A \cap E$  and  $C = A \cap E^c$ .

# Proposition 20.27

**Proposition 20.27.** Let  $\varphi$  be a real-valued function on a set  $X$  and  $\mu^* : 2^X \rightarrow [0, \infty]$  an outer measure with the property that whenever two subsets  $A$  and  $B$  of  $X$  are separated by  $\varphi$ , then

$$\mu^*(A \cup B) = \mu^*(A) + \mu^*(B).$$

Then  $\varphi$  is measurable with respect to the measure induced by  $\mu^*$ .

**Proof.** Let  $a \in \mathbb{R}$ . We show that  $E = \{x \in X \mid \varphi(x) > a\}$  is  $\mu^*$ -measurable, implying the measurability of function  $\varphi$ . By definition, an outer measure is countably monotone (see Section 17.3) so

$$\mu^*(A) = \mu^*((A \cap E) \cup (A \cap E^c)) \leq \mu^*(A \cap E) + \mu^*(A \cap E^c).$$

So we need only show that for  $A$  a set of finite measure and for any  $\varepsilon > 0$ ,

$$\mu^*(A) + \varepsilon > \mu^*(A \cap E) + \mu^*(A \cap E^c). \quad (32)$$

Notice that (32) trivially holds if  $\mu^*(A) = \infty$ , so we can assume  $\mu^*(A) < \infty$ . Define  $B = A \cap E$  and  $C = A \cap E^c$ .

# Proposition 20.27 (continued 1)

**Proof (continued).** For each  $n \in \mathbb{N}$ , define

$$B_n = \{x \in I \mid \varphi(x) > a + 1/n\} \text{ and } R_n = B_n \setminus B_{n-1}.$$

Notice that  $B_1 \subset B_2 \subset \dots$ ,  $\bigcup_{k=1}^{\infty} B_k = B$ , and

$$B = b_n \cup \left( \bigcup_{k=n+1}^{\infty} (B_k \setminus B_{k-1}) \right) = B_n \cup \left( \bigcup_{k=n+1}^{\infty} R_k \right).$$

Now on  $B_{n-2}$  (by definition of  $B_n$ ) we have  $\varphi > a + 1/(n-2)$ , which on  $R_n = B_n \setminus B_{n-1}$  we have  $a + 1/n < \varphi \leq a + 1/(n-1)$ . Thus  $\varphi$  separates  $R_n$  and  $B_{n-2}$  and hence separates  $R_{2k}$  and  $\bigcup_{j=1}^{k-1} R_{2j}$  since

$\bigcup_{j=1}^{k-1} R_{2j} \subset B_{2k-2}$ . So by hypothesis,

$$\mu^* \left( \bigcup_{j=1}^k R_{2j} \right) = \mu^*(R_{2k}) + \mu^* \left( \bigcup_{j=1}^{k-1} R_{2j} \right).$$

So by induction on  $k$ ,

$$\mu^* \left( \bigcup_{j=1}^k R_{2j} \right) = \mu^*(R_{2k}) + \mu^* \left( \bigcup_{j=1}^{k-1} R_{2j} \right) = \sum_{j=1}^k \mu^*(R_{2j}).$$

# Proposition 20.27 (continued 1)

**Proof (continued).** For each  $n \in \mathbb{N}$ , define

$$B_n = \{x \in I \mid \varphi(x) > a + 1/n\} \text{ and } R_n = B_n \setminus B_{n-1}.$$

Notice that  $B_1 \subset B_2 \subset \cdots$ ,  $\bigcup_{k=1}^{\infty} B_k = B$ , and

$$B = b_n \cup \left( \bigcup_{k=n+1}^{\infty} (B_k \setminus B_{k-1}) \right) = B_n \cup \left( \bigcup_{k=n+1}^{\infty} R_k \right).$$

Now on  $B_{n-2}$  (by definition of  $B_n$ ) we have  $\varphi > a + 1/(n-2)$ , which on  $R_n = B_n \setminus B_{n-1}$  we have  $a + 1/n < \varphi \leq a + 1/(n-1)$ . Thus  $\varphi$  separates  $R_n$  and  $B_{n-2}$  and hence separates  $R_{2k}$  and  $\bigcup_{j=1}^{k-1} R_{2j}$  since

$\bigcup_{j=1}^{k-1} R_{2j} \subset B_{2k-2}$ . So by hypothesis,

$$\mu^* \left( \bigcup_{j=1}^k R_{2j} \right) = \mu^*(R_{2k}) + \mu^* \left( \bigcup_{j=1}^{k-1} R_{2j} \right).$$

So by induction on  $k$ ,

$$\mu^* \left( \bigcup_{j=1}^k R_{2j} \right) = \mu^*(R_{2k}) + \mu^* \left( \bigcup_{j=1}^{k-1} R_{2j} \right) = \sum_{j=1}^k \mu^*(R_{2j}).$$

## Proposition 20.27 (continued 2)

**Proof (continued).** Since  $\cup_{j=1}^k R_{2j} \subset B_{2k} \subset B \subset A$  we have by monotonicity that

$$\mu^* \left( \cup_{j=1}^k R_{2j} \right) \leq \mu^*(A) \text{ or } \sum_{j=1}^k \mu^*(R_{2j}) \leq \mu^*(A).$$

Since we have  $\mu^*(A) < \infty$  then the series  $\sum_{j=1}^{\infty} \mu^*(R_{2j})$  converges (absolutely). Similarly, the series  $\sum_{j=1}^{\infty} \mu^*(R_k)$  converges. So there is  $n \in \mathbb{N}$  such that  $\sum_{k=n+1}^{\infty} \mu^*(R_k) < \varepsilon$ . Since  $B = B_n \cup (\cup_{k=n+1}^{\infty} R_k)$  then by the countable monotonicity of  $\mu^*$ ,

$$\mu^*(B) \leq \mu^*(B_n) + \sum_{k=n+1}^{\infty} \mu^*(R_k) < \mu^*(B_n) + \varepsilon$$

or  $\mu^*(B_n) > \mu^*(B) - \varepsilon$ . Now  $C = A \cap E^c$  by definition so  $C \subset A$ , and  $B = A \cap E$  by definition so  $B \subset A$ . So by monotonicity of  $\mu^*$ ,  $\mu^*(A) \geq \mu^*(B_n \cup C)$ .

## Proposition 20.27 (continued 2)

**Proof (continued).** Since  $\cup_{j=1}^k R_{2j} \subset B_{2k} \subset B \subset A$  we have by monotonicity that

$$\mu^* \left( \cup_{j=1}^k R_{2j} \right) \leq \mu^*(A) \text{ or } \sum_{j=1}^k \mu^*(R_{2j}) \leq \mu^*(A).$$

Since we have  $\mu^*(A) < \infty$  then the series  $\sum_{j=1}^{\infty} \mu^*(R_{2j})$  converges (absolutely). Similarly, the series  $\sum_{j=1}^{\infty} \mu^*(R_k)$  converges. So there is  $n \in \mathbb{N}$  such that  $\sum_{k=n+1}^{\infty} \mu^*(R_k) < \varepsilon$ . Since  $B = B_n \cup (\cup_{k=n+1}^{\infty} R_k)$  then by the countable monotonicity of  $\mu^*$ ,

$$\mu^*(B) \leq \mu^*(B_n) + \sum_{k=n+1}^{\infty} \mu^*(R_k) < \mu^*(B_n) + \varepsilon$$

or  $\mu^*(B_n) > \mu^*(B) - \varepsilon$ . Now  $C = A \cap E^c$  by definition so  $C \subset A$ , and  $B = A \cap E$  by definition so  $B \subset A$ . So by monotonicity of  $\mu^*$ ,  $\mu^*(A) \geq \mu^*(B_n \cup C)$ .



# Proposition 20.27 (continued 3)

**Proposition 20.27.** Let  $\varphi$  be a real-valued function on a set  $X$  and  $\mu^* : 2^X \rightarrow [0, \infty]$  an outer measure with the property that whenever two subsets  $A$  and  $B$  of  $X$  are separated by  $\varphi$ , then

$$\mu^*(A \cup B) = \mu^*(A) + \mu^*(B).$$

Then  $\varphi$  is measurable with respect to the measure induced by  $\mu^*$ .

**Proof (continued).** Now  $\varphi > a + 1/n$  on  $B_n$  and  $\varphi \leq a$  on  $E^c$  (by the definition of  $E$ ) and so  $\varphi$  separates  $B_n$  and  $C$ . So by hypothesis,  $\mu^*(A) \geq \mu^*(B_n \cup C) = \mu^*(B_n) + \mu^*(C)$ . Since  $\mu^*(B_n) > \mu^*(B) - \varepsilon$  then  $\mu^*(A) > \mu^*(B) + \mu^*(C)$ , or  $\mu^*(A) + \varepsilon > \mu^*(A \cap E) + \mu^*(A \cap E^c)$ , and hence  $\mu^*(A) = \mu^*(A \cap E) + \mu^*(A \cap E^c)$  for all  $A \subset X$  where  $A$  is of finite outer measure. That is,  $E = \{x \in X \mid \varphi(x) > a\}$  is measurable. Since  $a \in \mathbb{R}$  is arbitrary, then function  $\varphi$  is measurable. □

## Proposition 20.28

**Theorem 20.28.** Let  $\mu^*$  be a Carathéodory outer measure on matrix space  $(X, \rho)$ . Then every Borel subset of  $X$  is measurable with respect to  $\mu^*$ .

**Proof.** The collection of Borel sets is the smallest  $\sigma$ -algebra containing the closed sets, and the measurable sets are a  $\sigma$ -algebra. So if we show that each closed set is measurable, then the result follows. For closed set  $F$ , define function  $f(x) = \rho(F, \{x\})$ . In Exercise 20.4.A it is to be shown that  $f$  is continuous on  $X$  and that  $f^{-1}(\{0\}) = F$ .

# Proposition 20.28

**Theorem 20.28.** Let  $\mu^*$  be a Carathéodory outer measure on matrix space  $(X, \rho)$ . Then every Borel subset of  $X$  is measurable with respect to  $\mu^*$ .

**Proof.** The collection of Borel sets is the smallest  $\sigma$ -algebra containing the closed sets, and the measurable sets are a  $\sigma$ -algebra. So if we show that each closed set is measurable, then the result follows. For closed set  $F$ , define function  $f(x) = \rho(F, \{x\})$ . In Exercise 20.4.A it is to be shown that  $f$  is continuous on  $X$  and that  $f^{-1}(\{0\}) = F$ . So if we show  $f$  is measurable then this implies  $f^{-1}(\{0\}) = \{x \in X \mid f(x) \geq 0\} \cap \{x \in X \mid f(x) \leq 0\}$  is measurable (that is, arbitrary closed set  $F$  is measurable) and the result then follows. We use Proposition 20.27 to show that every continuous real valued function on  $X$  is measurable. First, if a continuous function separates no sets, then the hypothesis of Proposition 20.27 are vacuously satisfied and that function is measurable.

# Proposition 20.28

**Theorem 20.28.** Let  $\mu^*$  be a Carathéodory outer measure on matrix space  $(X, \rho)$ . Then every Borel subset of  $X$  is measurable with respect to  $\mu^*$ .

**Proof.** The collection of Borel sets is the smallest  $\sigma$ -algebra containing the closed sets, and the measurable sets are a  $\sigma$ -algebra. So if we show that each closed set is measurable, then the result follows. For closed set  $F$ , define function  $f(x) = \rho(F, \{x\})$ . In Exercise 20.4.A it is to be shown that  $f$  is continuous on  $X$  and that  $f^{-1}(\{0\}) = F$ . So if we show  $f$  is measurable then this implies  $f^{-1}(\{0\}) = \{x \in X \mid f(x) \geq 0\} \cap \{x \in X \mid f(x) \leq 0\}$  is measurable (that is, arbitrary closed set  $F$  is measurable) and the result then follows. We use Proposition 20.27 to show that every continuous real valued function on  $X$  is measurable. First, if a continuous function separates no sets, then the hypothesis of Proposition 20.27 are vacuously satisfied and that function is measurable.

## Proposition 20.28 (continued)

**Proof.** Second, let  $A$  and  $B$  be subsets of  $X$  which are separated by some continuous function  $f$ , say  $f \leq a$  on  $A$  and  $f \geq b$  on  $B$  where  $a < b$ . ASSUME that  $\rho(A, B) = 0$ . Then there are sequences  $\{u_k\} \subset A$  and  $\{v_k\} \subset B$  such that  $\lim_{k \rightarrow \infty} \rho(u_k, v_k) = 0$ . Since  $f$  is continuous then  $\lim_{k \rightarrow \infty} |f(u_k) - f(v_k)| = 0$ . But then there is some  $u_N \in A$  and  $v_N \in B$  with  $|f(u_N) - f(v_N)| < (b - a)/2$ , a CONTRADICTION. So the assumption that  $\rho(A, B) = 0$  is false and in fact  $\rho(A, B) > 0$ .

# Proposition 20.28 (continued)

**Proof.** Second, let  $A$  and  $B$  be subsets of  $X$  which are separated by some continuous function  $f$ , say  $f \leq a$  on  $A$  and  $f \geq b$  on  $B$  where  $a < b$ . ASSUME that  $\rho(A, B) = 0$ . Then there are sequences  $\{u_k\} \subset A$  and  $\{v_k\} \subset B$  such that  $\lim_{k \rightarrow \infty} \rho(u_k, v_k) = 0$ . Since  $f$  is continuous then  $\lim_{k \rightarrow \infty} |f(u_k) - f(v_k)| = 0$ . But then there is some  $u_N \in A$  and  $v_N \in B$  with  $|f(u_N) - f(v_N)| < (b - a)/2$ , a CONTRADICTION. So the assumption that  $\rho(A, B) = 0$  is false and in fact  $\rho(A, B) > 0$ . Since  $\mu^*$  is a Carathéodory outer measure then (by definition)

$\mu^*(A \cup B) = \mu^*(A) + \mu^*(B)$ . So for any  $A, B \subset X$  separated by some continuous function  $f$ , we have by Proposition 20.27 that continuous  $f$  is a measurable function with respect to  $\mu^*$ . Since this holds for arbitrary  $A$  and  $B$ , all continuous functions are measurable with respect to  $\mu^*$ . As discussed above, this establishes that all Borel sets are  $\mu_n^*$ -measurable.  $\square$

# Proposition 20.28 (continued)

**Proof.** Second, let  $A$  and  $B$  be subsets of  $X$  which are separated by some continuous function  $f$ , say  $f \leq a$  on  $A$  and  $f \geq b$  on  $B$  where  $a < b$ . ASSUME that  $\rho(A, B) = 0$ . Then there are sequences  $\{u_k\} \subset A$  and  $\{v_k\} \subset B$  such that  $\lim_{k \rightarrow \infty} \rho(u_k, v_k) = 0$ . Since  $f$  is continuous then  $\lim_{k \rightarrow \infty} |f(u_k) - f(v_k)| = 0$ . But then there is some  $u_N \in A$  and  $v_N \in B$  with  $|f(u_N) - f(v_N)| < (b - a)/2$ , a CONTRADICTION. So the assumption that  $\rho(A, B) = 0$  is false and in fact  $\rho(A, B) > 0$ . Since  $\mu^*$  is a Carathéodory outer measure then (by definition)

$\mu^*(A \cup B) = \mu^*(A) + \mu^*(B)$ . So for *any*  $A, B \subset X$  separated by some continuous function  $f$ , we have by Proposition 20.27 that continuous  $f$  is a measurable function with respect to  $\mu^*$ . Since this holds for arbitrary  $A$  and  $B$ , all continuous functions are measurable with respect to  $\mu^*$ . As discussed above, this establishes that all Borel sets are  $\mu_n^*$ -measurable.  $\square$

# Proposition 20.29

**Proposition 20.29.** Let  $(X, \rho)$  be a metric space and  $\alpha$  a positive real number. Then  $H_\alpha^* : 2^X \rightarrow [0, \infty]$  is a Carathéodory outer measure.

**Proof.** First, for any  $\varepsilon > 0$ ,  $H_\alpha^{(\varepsilon)}(\emptyset) = 0$  and so  $H_\alpha^*(\emptyset) = 0$ . For countable monotonicity, let  $\{E_k\}_{k=1}^\infty$  be a countable cover of  $E$ . For any coverings  $\{A_i^k\}_{i=1}^\infty$  of  $E_k$  (for  $k = 1, 2, \dots$ ) we have  $\{A_i^k\}_{i,k=1}^\infty$  is a countable cover of  $E \subset \bigcup_{k=1}^\infty E_k$ .



# Proposition 20.29

**Proposition 20.29.** Let  $(X, \rho)$  be a metric space and  $\alpha$  a positive real number. Then  $H_\alpha^* : 2^X \rightarrow [0, \infty]$  is a Carathéodory outer measure.

**Proof.** First, for any  $\varepsilon > 0$ ,  $H_\alpha^{(\varepsilon)}(\emptyset) = 0$  and so  $H_\alpha^*(\emptyset) = 0$ . For countable monotonicity, let  $\{E_k\}_{k=1}^\infty$  be a countable cover of  $E$ . For any coverings  $\{A_i^k\}_{i=1}^\infty$  of  $E_k$  (for  $k = 1, 2, \dots$ ) we have  $\{A_i^k\}_{i,k=1}^\infty$  is a countable cover of  $E \subset \bigcup_{k=1}^\infty E_k$ . Now

$$\sum_{i,k=1}^\infty (\text{diam}(A_i^k))^\alpha = \sum_{k=1}^\infty \left( \sum_{i=1}^\infty (\text{diam}(A_i^k))^\alpha \right)$$

and taking an infimum over all such  $A_i^k$  we get

$$\inf \sum_{i,k=1}^\infty (\text{diam}(A_i^k))^\alpha = \inf \left( \sum_{k=1}^\infty \sum_{i=1}^\infty (\text{diam}(A_i^k))^\alpha \right)$$

# Proposition 20.29

**Proposition 20.29.** Let  $(X, \rho)$  be a metric space and  $\alpha$  a positive real number. Then  $H_\alpha^* : 2^X \rightarrow [0, \infty]$  is a Carathéodory outer measure.

**Proof.** First, for any  $\varepsilon > 0$ ,  $H_\alpha^{(\varepsilon)}(\emptyset) = 0$  and so  $H_\alpha^*(\emptyset) = 0$ . For countable monotonicity, let  $\{E_k\}_{k=1}^\infty$  be a countable cover of  $E$ . For any coverings  $\{A_i^k\}_{i=1}^\infty$  of  $E_k$  (for  $k = 1, 2, \dots$ ) we have  $\{A_i^k\}_{i,k=1}^\infty$  is a countable cover of  $E \subset \bigcup_{k=1}^\infty E_k$ . Now

$$\sum_{i,k=1}^\infty (\text{diam}(A_i^k))^\alpha = \sum_{k=1}^\infty \left( \sum_{i=1}^\infty (\text{diam}(A_i^k))^\alpha \right)$$

and taking an infimum over all such  $A_i^k$  we get

$$\inf \sum_{i,k=1}^\infty (\text{diam}(A_i^k))^\alpha = \inf \left( \sum_{k=1}^\infty \sum_{i=1}^\infty (\text{diam}(A_i^k))^\alpha \right)$$

## Proposition 20.29 (continued 1)

**Proof (continued).**

$$\begin{aligned}
 &= \sum_{k=1}^{\infty} \inf_k \sum_{i=1}^{\infty} \left( \text{diam}(A_i^k) \right)^\alpha \text{ where } \inf_k \text{ denotes an infimum} \\
 &\quad \text{over all } \{A_i^k\}_{i=1}^{\infty} \text{ coverings of } E_k \\
 &= \sum_{k=1}^{\infty} H_\alpha^{(\varepsilon)}(E_k).
 \end{aligned}$$

Now since  $\{A_i^k\}_{i,k=1}^{\infty}$  is *some* covering of  $E$  (namely, one based on a union of coverings of the  $E_k$ ) then when an infimum is taken over all coverings of  $E$  we have  $H_\alpha^{(\varepsilon)}(E) \leq \inf \sum_{i,k=1}^{\infty} \left( \text{diam}(A_i^k) \right)^\alpha$  and hence

$H_\alpha^{(\varepsilon)}(E) \leq \sum_{k=1}^{\infty} H_\alpha^{(\varepsilon)}(E_k)$ . a limit as  $\varepsilon \rightarrow 0$  we have

$H_\alpha^*(E) \leq \sum_{k=1}^{\infty} H_\alpha^*(E_k)$  so that  $H_\alpha^*$  is countably monotone. So  $H_\alpha^*$  is (by definition) an outer measure on  $2^X$ .

## Proposition 20.29 (continued 1)

**Proof (continued).**

$$\begin{aligned}
 &= \sum_{k=1}^{\infty} \inf_k \sum_{i=1}^{\infty} \left( \text{diam}(A_i^k) \right)^\alpha \text{ where } \inf_k \text{ denotes an infimum} \\
 &\quad \text{over all } \{A_i^k\}_{i=1}^{\infty} \text{ coverings of } E_k \\
 &= \sum_{k=1}^{\infty} H_\alpha^{(\varepsilon)}(E_k).
 \end{aligned}$$

Now since  $\{A_i^k\}_{i,k=1}^{\infty}$  is *some* covering of  $E$  (namely, one based on a union of coverings of the  $E_k$ ) then when an infimum is taken over all coverings of  $E$  we have  $H_\alpha^{(\varepsilon)}(E) \leq \inf \sum_{i,k=1}^{\infty} \left( \text{diam}(A_i^k) \right)^\alpha$  and hence

$H_\alpha^{(\varepsilon)}(E) \leq \sum_{k=1}^{\infty} H_\alpha^{(\varepsilon)}(E_k)$ . a limit as  $\varepsilon \rightarrow 0$  we have

$H_\alpha^*(E) \leq \sum_{k=1}^{\infty} H_\alpha^*(E_k)$  so that  $H_\alpha^*$  is countably monotone. So  $H_\alpha^*$  is (by definition) an outer measure on  $2^X$ .

## Proposition 20.29 (continued 2)

**Proof (continued).** To establish that  $H_\alpha^*$  is a Carathéodory outer measure, let  $E, F \subset X$  for which  $\rho(E, F) > \delta > 0$ . Let  $\varepsilon > 0$  be such that  $\varepsilon < \delta$ . If  $\{A_k\}_{k=1}^\infty$  is a cover of  $E \cup F$  then since  $\varepsilon < \delta$  then each  $A_k$  can intersect at most one of  $E$  and  $F$ . So any such cover of  $E \cup F$  yields a cover of  $E$  and a cover of  $F$ . Taking an infimum over all such coverings of  $E \cup F$  we have

$$\begin{aligned} H_\alpha^{(\varepsilon)}(E \cup F) &= \inf \sum_{k=1}^{\infty} (\text{diam}(A_k))^\alpha \\ &\geq \inf_E \sum_{k=1}^{\infty} (\text{diam}(A_k^E))^\alpha + \inf_F \sum_{k=1}^{\infty} (\text{diam}(A_k^E))^\alpha = H_\alpha^{(\varepsilon)}(E) + H_\alpha^{(\varepsilon)}(F) \end{aligned}$$

where  $\inf_E$  is an infimum over all coverings  $\{A_k^E\}_{k=1}^\infty$  of  $E$  (and similarly for  $\inf_F$ ); the inequality is introduced since (for  $\varepsilon < \delta$ ) every covering  $E \cup F$  implies a covering of  $E$  and a covering of  $F$  but  $\inf_E$  and  $\inf_F$  involves *more* potential coverings of  $E$  and  $F$ .

## Proposition 20.29 (continued 2)

**Proof (continued).** To establish that  $H_\alpha^*$  is a Carathéodory outer measure, let  $E, F \subset X$  for which  $\rho(E, F) > \delta > 0$ . Let  $\varepsilon > 0$  be such that  $\varepsilon < \delta$ . If  $\{A_k\}_{k=1}^\infty$  is a cover of  $E \cup F$  then since  $\varepsilon < \delta$  then each  $A_k$  can intersect at most one of  $E$  and  $F$ . So any such cover of  $E \cup F$  yields a cover of  $E$  and a cover of  $F$ . Taking an infimum over all such coverings of  $E \cup F$  we have

$$\begin{aligned} H_\alpha^{(\varepsilon)}(E \cup F) &= \inf \sum_{k=1}^{\infty} (\text{diam}(A_k))^\alpha \\ &\geq \inf_E \sum_{k=1}^{\infty} (\text{diam}(A_k^E))^\alpha + \inf_F \sum_{k=1}^{\infty} (\text{diam}(A_k^E))^\alpha = H_\alpha^{(\varepsilon)}(E) + H_\alpha^{(\varepsilon)}(F) \end{aligned}$$

where  $\inf_E$  is an infimum over all coverings  $\{A_k^E\}_{k=1}^\infty$  of  $E$  (and similarly for  $\inf_F$ ); the inequality is introduced since (for  $\varepsilon < \delta$ ) every covering  $E \cup F$  implies a covering of  $E$  and a covering of  $F$  but  $\inf_E$  and  $\inf_F$  involves *more* potential coverings of  $E$  and  $F$ .

## Proposition 20.29 (continued 3)

**Proposition 20.29.** Let  $(X, \rho)$  be a metric space and  $\alpha$  a positive real number. Then  $H_\alpha^* : 2^X \rightarrow [0, \infty]$  is a Carathéodory outer measure.

**Proof (continued).** Taking a limit as  $\varepsilon \rightarrow 0$  we get  $H_\alpha^*(E \cup F) \geq H_\alpha^*(E) + H_\alpha^*(F)$ . We showed above that  $H_\alpha^*$  is countably monotone and so  $H_\alpha^*(E \cup F) \leq H_\alpha^*(E) + H_\alpha^*(F)$  so that  $H_\alpha^*$  is (by definition) a Carathéodory outer measure. □

# Proposition 20.30

**Proposition 20.30.** Let  $(X, \rho)$  be a metric space. Let  $A$  be a Borel subset of  $X$ , and let  $\alpha, \beta$  be positive real numbers for which  $\alpha < \beta$ . If  $H_\alpha(A) < \infty$  then  $H_\beta(A) = 0$ .

**Proof.** Let  $\varepsilon > 0$ . Choose  $\{A_k\}_{k=1}^\infty$  as a covering of  $A$  by sets of diameter less than or equal to  $\varepsilon$  for which

$$\sum_{k=1}^{\infty} (\text{diam}(A_k))^\alpha \leq H_\alpha^*(A) + 1 = H_\alpha(A) + 1$$

(which can be done, by the definition of “infimum”).



# Proposition 20.30

**Proposition 20.30.** Let  $(X, \rho)$  be a metric space. Let  $A$  be a Borel subset of  $X$ , and let  $\alpha, \beta$  be positive real numbers for which  $\alpha < \beta$ . If  $H_\alpha(A) < \infty$  then  $H_\beta(A) = 0$ .

**Proof.** Let  $\varepsilon > 0$ . Choose  $\{A_k\}_{k=1}^\infty$  as a covering of  $A$  by sets of diameter less than or equal to  $\varepsilon$  for which

$$\sum_{k=1}^{\infty} (\text{diam}(A_k))^\alpha \leq H_\alpha^*(A) + 1 = H_\alpha(A) + 1$$

(which can be done, by the definition of “infimum”). Then

$$\begin{aligned} H_\beta^{(\varepsilon)}(A) &\leq \sum_{k=1}^{\infty} (\text{diam}(A_k))^\beta \text{ by the infimum definition of } H_\beta^{(\varepsilon)}(A) \\ &= \varepsilon^\beta \sum_{k=1}^{\infty} \left( \frac{\text{diam}(A_k)}{\varepsilon} \right)^\alpha \end{aligned}$$

# Proposition 20.30

**Proposition 20.30.** Let  $(X, \rho)$  be a metric space. Let  $A$  be a Borel subset of  $X$ , and let  $\alpha, \beta$  be positive real numbers for which  $\alpha < \beta$ . If  $H_\alpha(A) < \infty$  then  $H_\beta(A) = 0$ .

**Proof.** Let  $\varepsilon > 0$ . Choose  $\{A_k\}_{k=1}^\infty$  as a covering of  $A$  by sets of diameter less than or equal to  $\varepsilon$  for which

$$\sum_{k=1}^{\infty} (\text{diam}(A_k))^\alpha \leq H_\alpha^*(A) + 1 = H_\alpha(A) + 1$$

(which can be done, by the definition of “infimum”). Then

$$\begin{aligned} H_\beta^{(\varepsilon)}(A) &\leq \sum_{k=1}^{\infty} (\text{diam}(A_k))^\beta \text{ by the infimum definition of } H_\beta^{(\varepsilon)}(A) \\ &= \varepsilon^\beta \sum_{k=1}^{\infty} \left( \frac{\text{diam}(A_k)}{\varepsilon} \right)^\alpha \end{aligned}$$

## Proposition 20.30 (continued 1)

**Proof (continued).**

$$\begin{aligned} &\leq \varepsilon^\beta \sum_{k=1}^{\infty} \left( \frac{\text{diam}(A_k)}{\varepsilon} \right)^\alpha \text{ since } \text{diam}(A_k) \leq \varepsilon, \frac{\text{diam}(A_k)}{\varepsilon} \leq 1 \text{ and so} \\ &\quad \left( \frac{\text{diam}(A_k)}{\varepsilon} \right)^\beta \leq \left( \frac{\text{diam}(A_k)}{\varepsilon} \right)^\alpha \text{ because } \alpha < \beta \\ &= \varepsilon^{\beta-\alpha} \sum_{k=1}^{\infty} (\text{diam}(A_k))^\alpha \leq \varepsilon^{\beta-\alpha} (H_\alpha(A) + 1). \end{aligned}$$

Taking a limit as  $\varepsilon \rightarrow 0$  we get

$$H_\beta^*(A) = \lim_{\varepsilon \rightarrow 0} H_\beta^{(\varepsilon)}(A) \leq \lim_{\varepsilon \rightarrow 0} \varepsilon^{\beta-\alpha} (H_\alpha(A) + 1) = 0$$

since  $\beta - \alpha > 0$  and  $H_\alpha(A) + 1$  is finite. Since  $H_\beta^*(A) = 0$  (here,  $H_\beta^*$  is a Carathéodory outer measure) then the induced Hausdorff  $\beta$ -dimensional measure satisfies  $H_\beta(A) = 0$ . □

## Proposition 20.30 (continued 1)

**Proof (continued).**

$$\begin{aligned} &\leq \varepsilon^\beta \sum_{k=1}^{\infty} \left( \frac{\text{diam}(A_k)}{\varepsilon} \right)^\alpha \text{ since } \text{diam}(A_k) \leq \varepsilon, \frac{\text{diam}(A_k)}{\varepsilon} \leq 1 \text{ and so} \\ &\quad \left( \frac{\text{diam}(A_k)}{\varepsilon} \right)^\beta \leq \left( \frac{\text{diam}(A_k)}{\varepsilon} \right)^\alpha \text{ because } \alpha < \beta \\ &= \varepsilon^{\beta-\alpha} \sum_{k=1}^{\infty} (\text{diam}(A_k))^\alpha \leq \varepsilon^{\beta-\alpha} (H_\alpha(A) + 1). \end{aligned}$$

Taking a limit as  $\varepsilon \rightarrow 0$  we get

$$H_\beta^*(A) = \lim_{\varepsilon \rightarrow 0} H_\beta^{(\varepsilon)}(A) \leq \lim_{\varepsilon \rightarrow 0} \varepsilon^{\beta-\alpha} (H_\alpha(A) + 1) = 0$$

since  $\beta - \alpha > 0$  and  $H_\alpha(A) + 1$  is finite. Since  $H_\beta^*(A) = 0$  (here,  $H_\beta^*$  is a Carathéodory outer measure) then the induced Hausdorff  $\beta$ -dimensional measure satisfies  $H_\beta(A) = 0$ . □

# Theorem 20.4.A

**Theorem 20.4.A.** The Hausdorff 1-dimensional measure,  $H_1$ , is the same as Lebesgue measure on the  $\sigma$ -algebra of Lebesgue measurable sets of real numbers.

**Proof.** Let  $I \subset \mathbb{R}$  be an interval. Given  $\varepsilon > 0$ ,  $I$  can be expressed as the disjoint union of subintervals of length less than  $\varepsilon$  and the diameter of each subinterval is its length (we make no restriction on any of the intervals in terms of open/closed). So

$$H_1^{(\varepsilon)}(I) = \inf \sum_{k=1}^{\infty} \ell(I_k) = m^*(I) = m(I) = \ell(I)$$

and so  $H_1(I) = \lim_{\varepsilon \rightarrow 0} H_1^{(\varepsilon)}(I) = m^*(I)$ .

# Theorem 20.4.A

**Theorem 20.4.A.** The Hausdorff 1-dimensional measure,  $H_1$ , is the same as Lebesgue measure on the  $\sigma$ -algebra of Lebesgue measurable sets of real numbers.

**Proof.** Let  $I \subset \mathbb{R}$  be an interval. Given  $\varepsilon > 0$ ,  $I$  can be expressed as the disjoint union of subintervals of length less than  $\varepsilon$  and the diameter of each subinterval is its length (we make no restriction on any of the intervals in terms of open/closed). So

$$H_1^{(\varepsilon)}(I) = \inf \sum_{k=1}^{\infty} \ell(I_k) = m^*(I) = m(I) = \ell(I)$$

and so  $H_1(I) = \lim_{\varepsilon \rightarrow 0} H_1^{(\varepsilon)}(I) = m^*(I)$ . Thus  $H_1$  and Lebesgue measure agree on the semiring of intervals of real numbers. Since  $H_1$  and Lebesgue measure are extensions of the same premeasure on the semiring of intervals, then by the uniqueness claim of the Carathéodory-Hahn Theorem, Lebesgue measure and  $H_1$  are equal on the  $\sigma$ -algebra of measurable sets. □

# Theorem 20.4.A

**Theorem 20.4.A.** The Hausdorff 1-dimensional measure,  $H_1$ , is the same as Lebesgue measure on the  $\sigma$ -algebra of Lebesgue measurable sets of real numbers.

**Proof.** Let  $I \subset \mathbb{R}$  be an interval. Given  $\varepsilon > 0$ ,  $I$  can be expressed as the disjoint union of subintervals of length less than  $\varepsilon$  and the diameter of each subinterval is its length (we make no restriction on any of the intervals in terms of open/closed). So

$$H_1^{(\varepsilon)}(I) = \inf \sum_{k=1}^{\infty} \ell(I_k) = m^*(I) = m(I) = \ell(I)$$

and so  $H_1(I) = \lim_{\varepsilon \rightarrow 0} H_1^{(\varepsilon)}(I) = m^*(I)$ . Thus  $H_1$  and Lebesgue measure agree on the semiring of intervals of real numbers. Since  $H_1$  and Lebesgue measure are extensions of the same premeasure on the semiring of intervals, then by the uniqueness claim of the Carathéodory-Hahn Theorem, Lebesgue measure and  $H_1$  are equal on the  $\sigma$ -algebra of measurable sets. □