


# Introduction to Topology

## Chapter 2. Topological Spaces and Continuous Functions

### Section 18. Continuous Functions—Proofs of Theorems



# Table of contents

- 1 Lemma 18.A
- 2 Theorem 18.1
- 3 Theorem 18.2(a,b,c)
- 4 Theorem 18.2(d,e,f)
- 5 Theorem 18.3
- 6 Theorem 18.4

## Lemma 18.A

**Lemma 18.A.** Let  $f : X \rightarrow Y$ , let  $\mathcal{B}$  be a basis for the topology on  $Y$ , and let  $\mathcal{S}$  be a subbasis for the topology on  $Y$ .

- (1)  $f$  is continuous if  $f^{-1}(B)$  is open in  $X$  for each  $B \in \mathcal{B}$ .
- (2)  $f$  is continuous if  $f^{-1}(S)$  is open in  $X$  for each  $S \in \mathcal{S}$ .

**Proof.** (1) Let  $V \subset Y$  be open. Then (by definition of basis) there are  $B_\alpha \in \mathcal{B}$  for  $\alpha \in J$  such that  $V = \cup_{\alpha \in J} B_\alpha$ .

## Lemma 18.A

**Lemma 18.A.** Let  $f : X \rightarrow Y$ , let  $\mathcal{B}$  be a basis for the topology on  $Y$ , and let  $\mathcal{S}$  be a subbasis for the topology on  $Y$ .

- (1)  $f$  is continuous if  $f^{-1}(B)$  is open in  $X$  for each  $B \in \mathcal{B}$ .
- (2)  $f$  is continuous if  $f^{-1}(S)$  is open in  $X$  for each  $S \in \mathcal{S}$ .

**Proof.** (1) Let  $V \subset Y$  be open. Then (by definition of basis) there are  $B_\alpha \in \mathcal{B}$  for  $\alpha \in J$  such that  $V = \bigcup_{\alpha \in J} B_\alpha$ . Then  $f^{-1}(V) = f^{-1}(\bigcup_{\alpha \in J} B_\alpha) = \bigcup_{\alpha \in J} f^{-1}(B_\alpha)$  is an open set in  $X$  by hypothesis. So each  $f^{-1}(B_\alpha)$  is open in  $X$  and  $f^{-1}(V)$  is open in  $X$ . Hence  $f$  is continuous.

## Lemma 18.A

**Lemma 18.A.** Let  $f : X \rightarrow Y$ , let  $\mathcal{B}$  be a basis for the topology on  $Y$ , and let  $\mathcal{S}$  be a subbasis for the topology on  $Y$ .

- (1)  $f$  is continuous if  $f^{-1}(B)$  is open in  $X$  for each  $B \in \mathcal{B}$ .
- (2)  $f$  is continuous if  $f^{-1}(S)$  is open in  $X$  for each  $S \in \mathcal{S}$ .

**Proof.** (1) Let  $V \subset Y$  be open. Then (by definition of basis) there are  $B_\alpha \in \mathcal{B}$  for  $\alpha \in J$  such that  $V = \bigcup_{\alpha \in J} B_\alpha$ . Then  $f^{-1}(V) = f^{-1}(\bigcup_{\alpha \in J} B_\alpha) = \bigcup_{\alpha \in J} f^{-1}(B_\alpha)$  is an open set in  $X$  by hypothesis. So each  $f^{-1}(B_\alpha)$  is open in  $X$  and  $f^{-1}(V)$  is open in  $X$ . Hence  $f$  is continuous.

## Lemma 18.A (continued)

**Lemma 18.A.** Let  $f : X \rightarrow Y$ , let  $\mathcal{B}$  be a basis for the topology on  $Y$ , and let  $\mathcal{S}$  be a subbasis for the topology on  $Y$ .

- (1)  $f$  is continuous if  $f^{-1}(B)$  is open in  $X$  for each  $B \in \mathcal{B}$ .
- (2)  $f$  is continuous if  $f^{-1}(S)$  is open in  $X$  for each  $S \in \mathcal{S}$ .

**Proof (continued).** (2) Let  $V \subset Y$  be open. Then (by the definition of subbasis) there are  $S_\alpha^i$  for  $\alpha \in J$ ,  $i \in \mathbb{N}$  such that  $V = \bigcup_{\alpha \in J} (S_\alpha^1 \cap S_\alpha^2 \cap \cdots \cap S_\alpha^{n_\alpha})$ .

## Lemma 18.A (continued)

**Lemma 18.A.** Let  $f : X \rightarrow Y$ , let  $\mathcal{B}$  be a basis for the topology on  $Y$ , and let  $\mathcal{S}$  be a subbasis for the topology on  $Y$ .

- (1)  $f$  is continuous if  $f^{-1}(B)$  is open in  $X$  for each  $B \in \mathcal{B}$ .
- (2)  $f$  is continuous if  $f^{-1}(S)$  is open in  $X$  for each  $S \in \mathcal{S}$ .

**Proof (continued).** (2) Let  $V \subset Y$  be open. Then (by the definition of subbasis) there are  $S_\alpha^i$  for  $\alpha \in J$ ,  $i \in \mathbb{N}$  such that  $V = \cup_{\alpha \in J} (S_\alpha^1 \cap S_\alpha^2 \cap \cdots \cap S_\alpha^{n_\alpha})$ . Then

$$\begin{aligned} f^{-1}(V) &= f^{-1}(\cup_{\alpha \in J} (S_\alpha^1 \cap S_\alpha^2 \cap \cdots \cap S_\alpha^{n_\alpha})) = \cup_{\alpha \in J} f^{-1}(S_\alpha^1 \cap S_\alpha^2 \cap \cdots \cap S_\alpha^{n_\alpha}) \\ &= \cup_{\alpha \in J} (f^{-1}(S_\alpha^1) \cap f^{-1}(S_\alpha^2) \cap \cdots \cap f^{-1}(S_\alpha^{n_\alpha})) \end{aligned}$$

is open in  $X$  since each  $f^{-1}(S_\alpha^i)$  is open in  $X$  by hypothesis and so  $f^{-1}(S_\alpha^1) \cap f^{-1}(S_\alpha^2) \cap \cdots \cap f^{-1}(S_\alpha^{n_\alpha})$  is open for each  $\alpha \in J$ , and hence the union is open. So  $f^{-1}(V)$  is open and  $f$  is continuous.  $\square$

# Lemma 18.A (continued)

**Lemma 18.A.** Let  $f : X \rightarrow Y$ , let  $\mathcal{B}$  be a basis for the topology on  $Y$ , and let  $\mathcal{S}$  be a subbasis for the topology on  $Y$ .

- (1)  $f$  is continuous if  $f^{-1}(B)$  is open in  $X$  for each  $B \in \mathcal{B}$ .
- (2)  $f$  is continuous if  $f^{-1}(S)$  is open in  $X$  for each  $S \in \mathcal{S}$ .

**Proof (continued).** (2) Let  $V \subset Y$  be open. Then (by the definition of subbasis) there are  $S_\alpha^i$  for  $\alpha \in J$ ,  $i \in \mathbb{N}$  such that  $V = \cup_{\alpha \in J} (S_\alpha^1 \cap S_\alpha^2 \cap \dots \cap S_\alpha^{n_\alpha})$ . Then

$$\begin{aligned} f^{-1}(V) &= f^{-1}(\cup_{\alpha \in J} (S_\alpha^1 \cap S_\alpha^2 \cap \dots \cap S_\alpha^{n_\alpha})) = \cup_{\alpha \in J} f^{-1}(S_\alpha^1 \cap S_\alpha^2 \cap \dots \cap S_\alpha^{n_\alpha}) \\ &= \cup_{\alpha \in J} (f^{-1}(S_\alpha^1) \cap f^{-1}(S_\alpha^2) \cap \dots \cap f^{-1}(S_\alpha^{n_\alpha})) \end{aligned}$$

is open in  $X$  since each  $f^{-1}(S_\alpha^i)$  is open in  $X$  by hypothesis and so  $f^{-1}(S_\alpha^1) \cap f^{-1}(S_\alpha^2) \cap \dots \cap f^{-1}(S_\alpha^{n_\alpha})$  is open for each  $\alpha \in J$ , and hence the union is open. So  $f^{-1}(V)$  is open and  $f$  is continuous.  $\square$



# Theorem 18.1

**Theorem 18.1.** Let  $X$  and  $Y$  be topological spaces. Let  $f : X \rightarrow Y$ . Then the following are equivalent:

- (1)  $f$  is continuous.
- (2) For every subset  $Z$  of  $X$ , one has  $f(\overline{Z}) \subset \overline{f(Z)}$ .
- (3) For every closed subset  $B$  of  $Y$ , the set  $f^{-1}(B)$  is closed in  $X$ .
- (4) For each  $x \in X$  and each neighborhood  $V$  of  $f(x)$ , there is a neighborhood  $U$  of  $x$  such that  $f(U) \subset V$ .

**Proof.** (1) $\Rightarrow$ (2) Suppose  $f$  is continuous. Let  $A \subset X$  and  $x \in \overline{A}$ . If  $x \in A$  then  $f(x) \in f(A) \subset \overline{f(A)}$ .

# Theorem 18.1

**Theorem 18.1.** Let  $X$  and  $Y$  be topological spaces. Let  $f : X \rightarrow Y$ . Then the following are equivalent:

- (1)  $f$  is continuous.
- (2) For every subset  $Z$  of  $X$ , one has  $f(\overline{A}) \subset \overline{f(A)}$ .
- (3) For every closed subset  $B$  of  $Y$ , the set  $f^{-1}(B)$  is closed in  $X$ .
- (4) For each  $x \in X$  and each neighborhood  $V$  of  $f(x)$ , there is a neighborhood  $U$  of  $x$  such that  $f(U) \subset V$ .

**Proof.** (1) $\Rightarrow$ (2) Suppose  $f$  is continuous. Let  $A \subset X$  and  $x \in \overline{A}$ . If  $x \in A$  then  $f(x) \in f(A) \subset \overline{f(A)}$ . If  $x \notin A$  then let  $V$  be a neighborhood of  $f(x)$ . Then  $f^{-1}(V)$  is open in  $X$  and  $x \in f^{-1}(V)$ . By definition of  $\overline{A}$ ,  $f^{-1}(V)$  intersects  $A$  at some point  $y \neq x$ .

# Theorem 18.1

**Theorem 18.1.** Let  $X$  and  $Y$  be topological spaces. Let  $f : X \rightarrow Y$ . Then the following are equivalent:

- (1)  $f$  is continuous.
- (2) For every subset  $Z$  of  $X$ , one has  $f(\overline{A}) \subset \overline{f(A)}$ .
- (3) For every closed subset  $B$  of  $Y$ , the set  $f^{-1}(B)$  is closed in  $X$ .
- (4) For each  $x \in X$  and each neighborhood  $V$  of  $f(x)$ , there is a neighborhood  $U$  of  $x$  such that  $f(U) \subset V$ .

**Proof.** (1) $\Rightarrow$ (2) Suppose  $f$  is continuous. Let  $A \subset X$  and  $x \in \overline{A}$ . If  $x \in A$  then  $f(x) \in f(A) \subset \overline{f(A)}$ . If  $x \notin A$  then let  $V$  be a neighborhood of  $f(x)$ . Then  $f^{-1}(V)$  is open in  $X$  and  $x \in f^{-1}(V)$ . By definition of  $\overline{A}$ ,  $f^{-1}(V)$  intersects  $A$  at some point  $y \neq x$ . So  $f(y) \in V \cap f(A)$  (notice that  $f(y) \neq f(x)$  since  $f(x) \notin f(A)$ ). So  $f(x) \in \overline{f(A)}$ . So  $f(x) \in \overline{f(A)}$  for any  $x \in \overline{A}$  and hence  $f(\overline{A}) \subset \overline{f(A)}$ .

# Theorem 18.1

**Theorem 18.1.** Let  $X$  and  $Y$  be topological spaces. Let  $f : X \rightarrow Y$ . Then the following are equivalent:

- (1)  $f$  is continuous.
- (2) For every subset  $Z$  of  $X$ , one has  $f(\overline{A}) \subset \overline{f(A)}$ .
- (3) For every closed subset  $B$  of  $Y$ , the set  $f^{-1}(B)$  is closed in  $X$ .
- (4) For each  $x \in X$  and each neighborhood  $V$  of  $f(x)$ , there is a neighborhood  $U$  of  $x$  such that  $f(U) \subset V$ .

**Proof.** (1) $\Rightarrow$ (2) Suppose  $f$  is continuous. Let  $A \subset X$  and  $x \in \overline{A}$ . If  $x \in A$  then  $f(x) \in f(A) \subset \overline{f(A)}$ . If  $x \notin A$  then let  $V$  be a neighborhood of  $f(x)$ . Then  $f^{-1}(V)$  is open in  $X$  and  $x \in f^{-1}(V)$ . By definition of  $\overline{A}$ ,  $f^{-1}(V)$  intersects  $A$  at some point  $y \neq x$ . So  $f(y) \in V \cap f(A)$  (notice that  $f(y) \neq f(x)$  since  $f(x) \notin f(A)$ ). So  $f(x) \in \overline{f(A)}$ . So  $f(x) \in \overline{f(A)}$  for any  $x \in \overline{A}$  and hence  $f(\overline{A}) \subset \overline{f(A)}$ .

# Theorem 18.1 (continued 1)

**Theorem 18.1.** Let  $X$  and  $Y$  be topological spaces. Let  $f : X \rightarrow Y$ . Then the following are equivalent:

- (1)  $f$  is continuous.
- (2) For every subset  $Z$  of  $X$ , one has  $f(\overline{A}) \subset \overline{f(A)}$ .
- (3) For every closed subset  $B$  of  $Y$ , the set  $f^{-1}(B)$  is closed in  $X$ .
- (4) For each  $x \in X$  and each neighborhood  $V$  of  $f(x)$ , there is a neighborhood  $U$  of  $x$  such that  $f(U) \subset V$ .

**Proof (continued).** (2) $\Rightarrow$ (3) Let  $B$  be closed in  $Y$  and let  $A = f^{-1}(B)$ . Then  $f(A) \subset B$  ( $f$  may not be onto  $B$  and so we may not have  $f(A) = B$ ). So if  $x \in \overline{A}$  then  $f(x) \in \overline{f(A)} \subset \overline{f(A)}$  by hypothesis (2) and  $\overline{f(A)} \subset \overline{B} = B$  since  $f(A) \subset B$  and  $B$  is closed.

# Theorem 18.1 (continued 1)

**Theorem 18.1.** Let  $X$  and  $Y$  be topological spaces. Let  $f : X \rightarrow Y$ . Then the following are equivalent:

- (1)  $f$  is continuous.
- (2) For every subset  $Z$  of  $X$ , one has  $f(\overline{A}) \subset \overline{f(A)}$ .
- (3) For every closed subset  $B$  of  $Y$ , the set  $f^{-1}(B)$  is closed in  $X$ .
- (4) For each  $x \in X$  and each neighborhood  $V$  of  $f(x)$ , there is a neighborhood  $U$  of  $x$  such that  $f(U) \subset V$ .

**Proof (continued).** (2) $\Rightarrow$ (3) Let  $B$  be closed in  $Y$  and let  $A = f^{-1}(B)$ . Then  $f(A) \subset B$  ( $f$  may not be onto  $B$  and so we may not have  $f(A) = B$ ). So if  $x \in \overline{A}$  then  $f(x) \in f(\overline{A}) \subset \overline{f(A)}$  by hypothesis (2) and  $\overline{f(A)} \subset \overline{B} = B$  since  $f(A) \subset B$  and  $B$  is closed. Hence  $f(x) \in B$  and  $x \in f^{-1}(B) = A$ . So  $\overline{A} \subset A$  and (since  $A \subset \overline{A}$ ) we have  $A = \overline{A}$  so that  $A = f^{-1}(B)$  is closed (by Lemma 17.A), as claimed.

# Theorem 18.1 (continued 1)

**Theorem 18.1.** Let  $X$  and  $Y$  be topological spaces. Let  $f : X \rightarrow Y$ . Then the following are equivalent:

- (1)  $f$  is continuous.
- (2) For every subset  $Z$  of  $X$ , one has  $f(\overline{A}) \subset \overline{f(A)}$ .
- (3) For every closed subset  $B$  of  $Y$ , the set  $f^{-1}(B)$  is closed in  $X$ .
- (4) For each  $x \in X$  and each neighborhood  $V$  of  $f(x)$ , there is a neighborhood  $U$  of  $x$  such that  $f(U) \subset V$ .

**Proof (continued).** (2) $\Rightarrow$ (3) Let  $B$  be closed in  $Y$  and let  $A = f^{-1}(B)$ . Then  $f(A) \subset B$  ( $f$  may not be onto  $B$  and so we may not have  $f(A) = B$ ). So if  $x \in \overline{A}$  then  $f(x) \in f(\overline{A}) \subset \overline{f(A)}$  by hypothesis (2) and  $\overline{f(A)} \subset \overline{B} = B$  since  $f(A) \subset B$  and  $B$  is closed. Hence  $f(x) \in B$  and  $x \in f^{-1}(B) = A$ . So  $\overline{A} \subset A$  and (since  $A \subset \overline{A}$ ) we have  $A = \overline{A}$  so that  $A = f^{-1}(B)$  is closed (by Lemma 17.A), as claimed.

## Theorem 18.1 (continued 2)

**Theorem 18.1.** Let  $X$  and  $Y$  be topological spaces. Let  $f : X \rightarrow Y$ . Then the following are equivalent:

- (1)  $f$  is continuous.
- (3) For every closed subset  $B$  of  $Y$ , the set  $f^{-1}(B)$  is closed in  $X$ .

**Proof (continued).** (3) $\Rightarrow$ (1) Let  $V$  be an open set in  $Y$ . Set  $B = Y \setminus V$ . Then

$$\begin{aligned} f^{-1}(B) &= f^{-1}(Y \setminus V) = f^{-1} \setminus f^{-1}(V) \text{ by Exercise 2.2(d)} \\ &= X \setminus f^{-1}(V) \text{ since } X \text{ is the domain of } f. \end{aligned}$$



## Theorem 18.1 (continued 2)

**Theorem 18.1.** Let  $X$  and  $Y$  be topological spaces. Let  $f : X \rightarrow Y$ . Then the following are equivalent:

- (1)  $f$  is continuous.
- (3) For every closed subset  $B$  of  $Y$ , the set  $f^{-1}(B)$  is closed in  $X$ .

**Proof (continued).** (3) $\Rightarrow$ (1) Let  $V$  be an open set in  $Y$ . Set  $B = Y \setminus V$ . Then

$$\begin{aligned} f^{-1}(B) &= f^{-1}(Y \setminus V) = f^{-1} \setminus f^{-1}(V) \text{ by Exercise 2.2(d)} \\ &= X \setminus f^{-1}(V) \text{ since } X \text{ is the domain of } f. \end{aligned}$$

Since  $V$  is open,  $B$  is closed in  $Y$  and so by hypothesis (3),  $f^{-1}(B) = X \setminus f^{-1}(V)$  is closed in  $X$  and so  $f^{-1}(V)$  is open. Therefore, by the definition of continuous function,  $f$  is continuous.

## Theorem 18.1 (continued 2)

**Theorem 18.1.** Let  $X$  and  $Y$  be topological spaces. Let  $f : X \rightarrow Y$ . Then the following are equivalent:

- (1)  $f$  is continuous.
- (3) For every closed subset  $B$  of  $Y$ , the set  $f^{-1}(B)$  is closed in  $X$ .

**Proof (continued).** (3) $\Rightarrow$ (1) Let  $V$  be an open set in  $Y$ . Set  $B = Y \setminus V$ . Then

$$\begin{aligned} f^{-1}(B) &= f^{-1}(Y \setminus V) = f^{-1} \setminus f^{-1}(V) \text{ by Exercise 2.2(d)} \\ &= X \setminus f^{-1}(V) \text{ since } X \text{ is the domain of } f. \end{aligned}$$

Since  $V$  is open,  $B$  is closed in  $Y$  and so by hypothesis (3),  $f^{-1}(B) = X \setminus f^{-1}(V)$  is closed in  $X$  and so  $f^{-1}(V)$  is open. Therefore, by the definition of continuous function,  $f$  is continuous.

## Theorem 18.1 (continued 3)

**Theorem 18.1.** Let  $X$  and  $Y$  be topological spaces. Let  $f : X \rightarrow Y$ . Then the following are equivalent:

- (1)  $f$  is continuous.
- (4) For each  $x \in X$  and each neighborhood  $V$  of  $f(x)$ , there is a neighborhood  $U$  of  $x$  such that  $f(U) \subset V$ .

**Proof (continued).** (1) $\Rightarrow$ (4) Let  $x \in X$  and let  $V$  be a neighborhood of  $f(x)$ . Then  $U = f^{-1}(V)$  is open since  $f$  is continuous and  $x \in U$ . That is,  $f(U) \subset V$ , as claimed.

## Theorem 18.1 (continued 3)

**Theorem 18.1.** Let  $X$  and  $Y$  be topological spaces. Let  $f : X \rightarrow Y$ . Then the following are equivalent:

- (1)  $f$  is continuous.
- (4) For each  $x \in X$  and each neighborhood  $V$  of  $f(x)$ , there is a neighborhood  $U$  of  $x$  such that  $f(U) \subset V$ .

**Proof (continued).** (1) $\Rightarrow$ (4) Let  $x \in X$  and let  $V$  be a neighborhood of  $f(x)$ . Then  $U = f^{-1}(V)$  is open since  $f$  is continuous and  $x \in U$ . That is,  $f(U) \subset V$ , as claimed.

(4) $\Rightarrow$ (1) Let  $V$  be an open set of  $Y$ . Let  $x \in f^{-1}(V)$ . Then  $f(x) \in V$  and so by hypothesis (4) there is open  $U_x$  in  $X$  with  $x \in U_x$  and  $f(U_x) \subset V$ . Then  $U_x \subset f^{-1}(V)$ .

## Theorem 18.1 (continued 3)

**Theorem 18.1.** Let  $X$  and  $Y$  be topological spaces. Let  $f : X \rightarrow Y$ . Then the following are equivalent:

- (1)  $f$  is continuous.
- (4) For each  $x \in X$  and each neighborhood  $V$  of  $f(x)$ , there is a neighborhood  $U$  of  $x$  such that  $f(U) \subset V$ .

**Proof (continued).** (1) $\Rightarrow$ (4) Let  $x \in X$  and let  $V$  be a neighborhood of  $f(x)$ . Then  $U = f^{-1}(V)$  is open since  $f$  is continuous and  $x \in U$ . That is,  $f(U) \subset V$ , as claimed.

(4) $\Rightarrow$ (1) Let  $V$  be an open set of  $Y$ . Let  $x \in f^{-1}(V)$ . Then  $f(x) \in V$  and so by hypothesis (4) there is open  $U_x$  in  $X$  with  $x \in U_x$  and  $f(U_x) \subset V$ . Then  $U_x \subset f^{-1}(V)$ . Then with such open  $U_x$  chosen for each  $x \in f^{-1}(V)$  we have  $f^{-1}(V) = \cup_{x \in f^{-1}(V)} U_x$  and hence  $f^{-1}(V)$  is open. Therefore, by the definition of continuous function,  $f$  is continuous and (1) follows.  $\square$

## Theorem 18.1 (continued 3)

**Theorem 18.1.** Let  $X$  and  $Y$  be topological spaces. Let  $f : X \rightarrow Y$ . Then the following are equivalent:

- (1)  $f$  is continuous.
- (4) For each  $x \in X$  and each neighborhood  $V$  of  $f(x)$ , there is a neighborhood  $U$  of  $x$  such that  $f(U) \subset V$ .

**Proof (continued).** (1) $\Rightarrow$ (4) Let  $x \in X$  and let  $V$  be a neighborhood of  $f(x)$ . Then  $U = f^{-1}(V)$  is open since  $f$  is continuous and  $x \in U$ . That is,  $f(U) \subset V$ , as claimed.

(4) $\Rightarrow$ (1) Let  $V$  be an open set of  $Y$ . Let  $x \in f^{-1}(V)$ . Then  $f(x) \in V$  and so by hypothesis (4) there is open  $U_x$  in  $X$  with  $x \in U_x$  and  $f(U_x) \subset V$ . Then  $U_x \subset f^{-1}(V)$ . Then with such open  $U_x$  chosen for each  $x \in f^{-1}(V)$  we have  $f^{-1}(V) = \bigcup_{x \in f^{-1}(V)} U_x$  and hence  $f^{-1}(V)$  is open. Therefore, by the definition of continuous function,  $f$  is continuous and (1) follows.  $\square$

# Theorem 18.2(a,b,c)

## Theorem 18.2. Rules for Constructing Continuous Functions.

Let  $X$ ,  $Y$ , and  $Z$  be topological spaces.

- (a) (Constant Function) If  $f : X \rightarrow Y$  maps all of  $X$  into a single point  $y_0 \in Y$ , then  $f$  is continuous.
- (b) (Inclusion) if  $A$  is a subspace of  $X$ , the inclusion function  $j : A \rightarrow X$  is continuous.
- (c) (Composites) If  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are continuous, then the map  $g \circ f : X \rightarrow Z$  is continuous.

**Proof.** (a) Let  $f(x) = y_0$  for every  $x \in X$ . Let  $V$  be open in  $Y$ . Then  $f^{-1}(V) = X$  if  $y_0 \in V$  and  $f^{-1}(V) = \emptyset$  if  $y_0 \notin V$ . In either case,  $f^{-1}(V)$  is open and so  $f$  is continuous.

# Theorem 18.2(a,b,c)

## Theorem 18.2. Rules for Constructing Continuous Functions.

Let  $X$ ,  $Y$ , and  $Z$  be topological spaces.

- (a) (Constant Function) If  $f : X \rightarrow Y$  maps all of  $X$  into a single point  $y_0 \in Y$ , then  $f$  is continuous.
- (b) (Inclusion) if  $A$  is a subspace of  $X$ , the inclusion function  $j : A \rightarrow X$  is continuous.
- (c) (Composites) If  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are continuous, then the map  $g \circ f : X \rightarrow Z$  is continuous.

**Proof.** (a) Let  $f(x) = y_0$  for every  $x \in X$ . Let  $V$  be open in  $Y$ . Then  $f^{-1}(V) = X$  if  $y_0 \in V$  and  $f^{-1}(V) = \emptyset$  if  $y_0 \notin V$ . In either case,  $f^{-1}(V)$  is open and so  $f$  is continuous. (b) If  $U$  is open in  $X$ , then  $j^{-1}(U) = U \cap A$  which is open in  $A$  (by definition of the subspace topology).



# Theorem 18.2(a,b,c)

## Theorem 18.2. Rules for Constructing Continuous Functions.

Let  $X$ ,  $Y$ , and  $Z$  be topological spaces.

- (a) (Constant Function) If  $f : X \rightarrow Y$  maps all of  $X$  into a single point  $y_0 \in Y$ , then  $f$  is continuous.
- (b) (Inclusion) if  $A$  is a subspace of  $X$ , the inclusion function  $j : A \rightarrow X$  is continuous.
- (c) (Composites) If  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are continuous, then the map  $g \circ f : X \rightarrow Z$  is continuous.

**Proof.** (a) Let  $f(x) = y_0$  for every  $x \in X$ . Let  $V$  be open in  $Y$ . Then  $f^{-1}(V) = X$  if  $y_0 \in V$  and  $f^{-1}(V) = \emptyset$  if  $y_0 \notin V$ . In either case,  $f^{-1}(V)$  is open and so  $f$  is continuous. (b) If  $U$  is open in  $X$ , then  $j^{-1}(U) = U \cap A$  which is open in  $A$  (by definition of the subspace topology). (c) If  $U$  is open in  $Z$  then  $g^{-1}(U)$  is open in  $Y$  since  $g$  is continuous and  $f^{-1}(g^{-1}(U))$  is open in  $X$  since  $f$  is continuous. Now  $(g \circ f)^{-1}(U) = f^{-1} \circ g^{-1}(U) = f^{-1}(g^{-1}(U))$  and so  $g \circ f$  is continuous. □

# Theorem 18.2(a,b,c)

## Theorem 18.2. Rules for Constructing Continuous Functions.

Let  $X$ ,  $Y$ , and  $Z$  be topological spaces.

- (a) (Constant Function) If  $f : X \rightarrow Y$  maps all of  $X$  into a single point  $y_0 \in Y$ , then  $f$  is continuous.
- (b) (Inclusion) if  $A$  is a subspace of  $X$ , the inclusion function  $j : A \rightarrow X$  is continuous.
- (c) (Composites) If  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are continuous, then the map  $g \circ f : X \rightarrow Z$  is continuous.

**Proof.** (a) Let  $f(x) = y_0$  for every  $x \in X$ . Let  $V$  be open in  $Y$ . Then  $f^{-1}(V) = X$  if  $y_0 \in V$  and  $f^{-1}(V) = \emptyset$  if  $y_0 \notin V$ . In either case,  $f^{-1}(V)$  is open and so  $f$  is continuous. (b) If  $U$  is open in  $X$ , then  $j^{-1}(U) = U \cap A$  which is open in  $A$  (by definition of the subspace topology). (c) If  $U$  is open in  $Z$  then  $g^{-1}(U)$  is open in  $Y$  since  $g$  is continuous and  $f^{-1}(g^{-1}(U))$  is open in  $X$  since  $f$  is continuous. Now  $(g \circ f)^{-1}(U) = f^{-1} \circ g^{-1}(U) = f^{-1}(g^{-1}(U))$  and so  $g \circ f$  is continuous. □

## Theorem 18.2

### Theorem 18.2. Rules for Constructing Continuous Functions.

Let  $X$ ,  $Y$ , and  $Z$  be topological spaces.

- (d) (Restricting the Domain) If  $f : X \rightarrow Y$  is continuous and if  $A$  is a subspace of  $X$ , then the restricted function  $f|_A : A \rightarrow Y$  is continuous.
- (e) (Restricting or Expanding the Range) let  $f : X \rightarrow Y$  be continuous. If  $X$  is a subspace of  $Y$  containing the image set  $f(X)$ , then the function  $g : X \rightarrow Z$  obtained by restricting the range of  $f$  is continuous. If  $Z$  is a space having  $Y$  as a subspace, then the functions  $h : X \rightarrow Z$  obtained by expanding the range of  $f$  is continuous.
- (e) (Local Formulation of Continuity) The map  $f : X \rightarrow Y$  is continuous if  $X$  can be written as the union of open sets  $U_\alpha$  such that  $f|_{U_\alpha}$  is continuous for each  $\alpha$ .

## Theorem 18.2(d, e, f) (continued 1)

**Proof.** (d) The function  $f|_A$  equals the composition of the inclusion map  $j : A \rightarrow Y$  (which is continuous by part (b)) and  $f : X \rightarrow Y$  (which is continuous by hypothesis). So by part (c),  $f|_A$  is continuous.

(e) Let  $f : X \rightarrow Y$  be continuous and  $f(X) \subset Z \subset Y$ . Let  $B$  be open in  $Z$ . Then (by definition)  $B = Z \cap U$  for some open  $U$  in  $Y$ .

## Theorem 18.2(d, e, f) (continued 1)

**Proof.** (d) The function  $f|_A$  equals the composition of the inclusion map  $j : A \rightarrow Y$  (which is continuous by part (b)) and  $f : X \rightarrow Y$  (which is continuous by hypothesis). So by part (c),  $f|_A$  is continuous.

(e) Let  $f : X \rightarrow Y$  be continuous and  $f(X) \subset Z \subset Y$ . Let  $B$  be open in  $Z$ . Then (by definition)  $B = Z \cap U$  for some open  $U$  in  $Y$ . Then

$$\begin{aligned} g^{-1}(B) &= g^{-1}(Z \cap U) = g^{-1}(Z) \cap g^{-1}(U) \\ &= X \cap g^{-1}(U) \text{ since } f(X) = g(X) \subset Z \\ &= g^{-1}(U) \\ &= f^{-1}(U) \text{ since } f(x) \in Y \text{ for some } x \in X \text{ implies } g(x) = f(x) \in \end{aligned}$$

Since  $f$  is continuous,  $f^{-1}(U)$  is open in  $X$  and so  $g^{-1}(U)$  is open in  $X$ . Therefore,  $g$  is continuous.

## Theorem 18.2(d, e, f) (continued 1)

**Proof.** (d) The function  $f|_A$  equals the composition of the inclusion map  $j : A \rightarrow Y$  (which is continuous by part (b)) and  $f : X \rightarrow Y$  (which is continuous by hypothesis). So by part (c),  $f|_A$  is continuous.

(e) Let  $f : X \rightarrow Y$  be continuous and  $f(X) \subset Z \subset Y$ . Let  $B$  be open in  $Z$ . Then (by definition)  $B = Z \cap U$  for some open  $U$  in  $Y$ . Then

$$\begin{aligned} g^{-1}(B) &= g^{-1}(Z \cap U) = g^{-1}(Z) \cap g^{-1}(U) \\ &= X \cap g^{-1}(U) \text{ since } f(X) = g(X) \subset Z \\ &= g^{-1}(U) \\ &= f^{-1}(U) \text{ since } f(x) \in Y \text{ for some } x \in X \text{ implies } g(x) = f(x) \in \end{aligned}$$

Since  $f$  is continuous,  $f^{-1}(U)$  is open in  $X$  and so  $g^{-1}(U)$  is open in  $X$ . Therefore,  $g$  is continuous.

Now let  $h : X \rightarrow Z \supset Y$  be as described. Then  $h$  is the composition of  $f : X \times Y$  (which is continuous by hypothesis) and the inclusion map  $j : Y \rightarrow Z$  (which is continuous by part (b)). So, by part (c),  $h$  is continuous.

## Theorem 18.2(d, e, f) (continued 1)

**Proof.** (d) The function  $f|_A$  equals the composition of the inclusion map  $j : A \rightarrow Y$  (which is continuous by part (b)) and  $f : X \rightarrow Y$  (which is continuous by hypothesis). So by part (c),  $f|_A$  is continuous.

(e) Let  $f : X \rightarrow Y$  be continuous and  $f(X) \subset Z \subset Y$ . Let  $B$  be open in  $Z$ . Then (by definition)  $B = Z \cap U$  for some open  $U$  in  $Y$ . Then

$$\begin{aligned} g^{-1}(B) &= g^{-1}(Z \cap U) = g^{-1}(Z) \cap g^{-1}(U) \\ &= X \cap g^{-1}(U) \text{ since } f(X) = g(X) \subset Z \\ &= g^{-1}(U) \\ &= f^{-1}(U) \text{ since } f(x) \in Y \text{ for some } x \in X \text{ implies } g(x) = f(x) \in \end{aligned}$$

Since  $f$  is continuous,  $f^{-1}(U)$  is open in  $X$  and so  $g^{-1}(U)$  is open in  $X$ . Therefore,  $g$  is continuous.

Now let  $h : X \rightarrow Z \supset Y$  be as described. Then  $h$  is the composition of  $f : X \times Y$  (which is continuous by hypothesis) and the inclusion map  $j : Y \rightarrow Z$  (which is continuous by part (b)). So, by part (c),  $h$  is continuous.

## Theorem 18.2(d, e, f) (continued 2)

**Proof.** (f) Suppose  $X = \cup_{\alpha \in J} U_\alpha$  for open  $U_\alpha$  in  $X$  where  $f|_{U_\alpha}$  is continuous for each  $\alpha \in J$ . Let  $V$  be an open set in  $Y$ . Since  $f^{-1}(V) \cap U_\alpha$  consists of  $x \in X \cap U_\alpha = U_\alpha$  such that  $f(x) \in V$  and  $(f|_{U_\alpha})^{-1}(V)$  consists of  $x \in U_\alpha$  such that  $f(x) \in V$ , then  $f^{-1}(V) \cap U_\alpha = (f|_{U_\alpha})^{-1}(V)$  for all  $\alpha \in J$ . Since  $f|_{U_\alpha}$  is continuous by hypothesis, then this set is open in  $U_\alpha$  and since  $U_\alpha$  is open then (by Lemma 16.2) this set is open in  $X$ .



## Theorem 18.2(d, e, f) (continued 2)

**Proof.** (f) Suppose  $X = \cup_{\alpha \in J} U_{\alpha}$  for open  $U_{\alpha}$  in  $X$  where  $f|_{U_{\alpha}}$  is continuous for each  $\alpha \in J$ . Let  $V$  be an open set in  $Y$ . Since  $f^{-1}(V) \cap U_{\alpha}$  consists of  $x \in X \cap U_{\alpha} = U_{\alpha}$  such that  $f(x) \in V$  and  $(f|_{U_{\alpha}})^{-1}(V)$  consists of  $x \in U_{\alpha}$  such that  $f(x) \in U_{\alpha}$ , then  $f^{-1}(V) \cap U_{\alpha} = (f|_{U_{\alpha}})^{-1}(V)$  for all  $\alpha \in J$ . Since  $f|_{U_{\alpha}}$  is continuous by hypothesis, then this set is open in  $U_{\alpha}$  and since  $U_{\alpha}$  is open then (by Lemma 16.2) this set is open in  $X$ . Since  $X = \cup_{\alpha \in J} U_{\alpha}$  then

$$f^{-1}(V) = f^{-1}(V) \cap X = f^{-1}(V) \cap (\cup_{\alpha \in J} U_{\alpha}) = \cup_{\alpha \in J} (f^{-1}(V) \cap U_{\alpha})$$

is open in  $X$  since each set in the union is open. Therefore (by definition)  $f$  is continuous.  $\square$

## Theorem 18.2(d, e, f) (continued 2)

**Proof.** (f) Suppose  $X = \cup_{\alpha \in J} U_\alpha$  for open  $U_\alpha$  in  $X$  where  $f|_{U_\alpha}$  is continuous for each  $\alpha \in J$ . Let  $V$  be an open set in  $Y$ . Since  $f^{-1}(V) \cap U_\alpha$  consists of  $x \in X \cap U_\alpha = U_\alpha$  such that  $f(x) \in V$  and  $(f|_{U_\alpha})^{-1}(V)$  consists of  $x \in U_\alpha$  such that  $f(x) \in U_\alpha$ , then  $f^{-1}(V) \cap U_\alpha = (f|_{U_\alpha})^{-1}(V)$  for all  $\alpha \in J$ . Since  $f|_{U_\alpha}$  is continuous by hypothesis, then this set is open in  $U_\alpha$  and since  $U_\alpha$  is open then (by Lemma 16.2) this set is open in  $X$ . Since  $X = \cup_{\alpha \in J} U_\alpha$  then

$$f^{-1}(V) = f^{-1}(V) \cap X = f^{-1}(V) \cap (\cup_{\alpha \in J} U_\alpha) = \cup_{\alpha \in J} (f^{-1}(V) \cap U_\alpha)$$

is open in  $X$  since each set in the union is open. Therefore (by definition)  $f$  is continuous. □

## Theorem 18.3

### Theorem 18.3. The Pasting Lemma for Closed Sets.

Let  $X = A \cup B$  where  $A$  and  $B$  are closed in  $X$ . Let  $f : A \rightarrow Y$  and  $g : B \rightarrow Y$  be continuous. If  $f(x) = g(x)$  for all  $x \in A \cap B$ , then  $f$  and  $g$  combine (or “paste”) to give a continuous function  $h : X \rightarrow Y$  defined by setting  $h(x) = f(x)$  if  $x \in A$  and  $h(x) = g(x)$  if  $x \in B$ .

**Proof.** Let  $C$  be closed in  $Y$ . Then  $h^{-1}(C) = f^{-1}(C) \cup g^{-1}(C)$ .

## Theorem 18.3

### Theorem 18.3. The Pasting Lemma for Closed Sets.

Let  $X = A \cup B$  where  $A$  and  $B$  are closed in  $X$ . Let  $f : A \rightarrow Y$  and  $g : B \rightarrow Y$  be continuous. If  $f(x) = g(x)$  for all  $x \in A \cup B$ , then  $f$  and  $g$  combine (or “paste”) to give a continuous function  $h : X \rightarrow Y$  defined by setting  $h(x) = f(x)$  if  $x \in A$  and  $h(x) = g(x)$  if  $x \in B$ .

**Proof.** Let  $C$  be closed in  $Y$ . Then  $h^{-1}(C) = f^{-1}(C) \cup g^{-1}(C)$ . Since  $f$  is continuous by hypothesis then  $f^{-1}(C)$  is closed in  $A$ , by Theorem 18.1 (the (1) $\Rightarrow$ (3) part), and so  $f^{-1}(C)$  is closed in  $X$  since  $A$  is closed (that is,  $f^{-1}(C) = A \cap D$  for closed  $D$  in  $X$ , so  $f^{-1}(C)$  is closed in  $X$ ). Similarly,  $g^{-1}(C)$  is closed in  $B$  and in  $X$ .

# Theorem 18.3

## Theorem 18.3. The Pasting Lemma for Closed Sets.

Let  $X = A \cup B$  where  $A$  and  $B$  are closed in  $X$ . Let  $f : A \rightarrow Y$  and  $g : B \rightarrow Y$  be continuous. If  $f(x) = g(x)$  for all  $x \in A \cup B$ , then  $f$  and  $g$  combine (or “paste”) to give a continuous function  $h : X \rightarrow Y$  defined by setting  $h(x) = f(x)$  if  $x \in A$  and  $h(x) = g(x)$  if  $x \in B$ .

**Proof.** Let  $C$  be closed in  $Y$ . Then  $h^{-1}(C) = f^{-1}(C) \cup g^{-1}(C)$ . Since  $f$  is continuous by hypothesis then  $f^{-1}(C)$  is closed in  $A$ , by Theorem 18.1 (the (1) $\Rightarrow$ (3) part), and so  $f^{-1}(C)$  is closed in  $X$  since  $A$  is closed (that is,  $f^{-1}(C) = A \cap D$  for closed  $D$  in  $X$ , so  $f^{-1}(C)$  is closed in  $X$ ). Similarly,  $g^{-1}(C)$  is closed in  $B$  and in  $X$ . Therefore  $h^{-1}(C)$  is closed in  $X$  and so by Theorem 18.2 (the (3) $\Rightarrow$ (1) part)  $h$  is continuous.  $\square$

## Theorem 18.3

**Theorem 18.3. The Pasting Lemma for Closed Sets.**

Let  $X = A \cup B$  where  $A$  and  $B$  are closed in  $X$ . Let  $f : A \rightarrow Y$  and  $g : B \rightarrow Y$  be continuous. If  $f(x) = g(x)$  for all  $x \in A \cup B$ , then  $f$  and  $g$  combine (or “paste”) to give a continuous function  $h : X \rightarrow Y$  defined by setting  $h(x) = f(x)$  if  $x \in A$  and  $h(x) = g(x)$  if  $x \in B$ .

**Proof.** Let  $C$  be closed in  $Y$ . Then  $h^{-1}(C) = f^{-1}(C) \cup g^{-1}(C)$ . Since  $f$  is continuous by hypothesis then  $f^{-1}(C)$  is closed in  $A$ , by Theorem 18.1 (the (1) $\Rightarrow$ (3) part), and so  $f^{-1}(C)$  is closed in  $X$  since  $A$  is closed (that is,  $f^{-1}(C) = A \cap D$  for closed  $D$  in  $X$ , so  $f^{-1}(C)$  is closed in  $X$ ). Similarly,  $g^{-1}(C)$  is closed in  $B$  and in  $X$ . Therefore  $h^{-1}(C)$  is closed in  $X$  and so by Theorem 18.2 (the (3) $\Rightarrow$ (1) part)  $h$  is continuous.  $\square$

## Theorem 18.4

### Theorem 18.4. Maps into Products.

Let  $f : A \rightarrow X \times Y$  be given by the equation  $f(a) = (f_1(a), f_2(a))$  where  $f_1 : A \rightarrow X$  and  $f_2 : A \rightarrow Y$ . Then  $f$  is continuous if and only if the functions  $f_1$  and  $f_2$  are continuous.

**Proof.** Let  $\pi_1 : X \times Y \rightarrow X$  and  $\pi_2 : X \times Y \rightarrow Y$ . Then for  $U$  open in  $X$  and  $V$  open in  $Y$ , we have  $\pi_1^{-1}(U) = U \times T$  and  $\pi_2^{-1}(V) = X \times V$  open in  $X \times Y$  (by the definition of product topology; these are basis elements for the product topology on  $X \times Y$ ). So  $\pi_1$  and  $\pi_2$  are continuous.

# Theorem 18.4

## Theorem 18.4. Maps into Products.

Let  $f : A \rightarrow X \times Y$  be given by the equation  $f(a) = (f_1(a), f_2(a))$  where  $f_1 : A \rightarrow X$  and  $f_2 : A \rightarrow Y$ . Then  $f$  is continuous if and only if the functions  $f_1$  and  $f_2$  are continuous.

**Proof.** Let  $\pi_1 : X \times Y \rightarrow X$  and  $\pi_2 : X \times Y \rightarrow Y$ . Then for  $U$  open in  $X$  and  $V$  open in  $Y$ , we have  $\pi_1^{-1}(U) = U \times T$  and  $\pi_2^{-1}(V) = X \times V$  open in  $X \times Y$  (by the definition of product topology; these are basis elements for the product topology on  $X \times Y$ ). So  $\pi_1$  and  $\pi_2$  are continuous. Note that for each  $a \in A$ ,  $\pi_1(f(a)) = \pi_1((f_1(a), f_2(a))) = f_1(a)$  and  $\pi_2(f(a)) = \pi_2((f_1(a), f_2(a))) = f_2(a)$ . So  $f_1 = \pi_1 \circ f$  and  $f_2 = \pi_2 \circ f$ .



# Theorem 18.4

## Theorem 18.4. Maps into Products.

Let  $f : A \rightarrow X \times Y$  be given by the equation  $f(a) = (f_1(a), f_2(a))$  where  $f_1 : A \rightarrow X$  and  $f_2 : A \rightarrow Y$ . Then  $f$  is continuous if and only if the functions  $f_1$  and  $f_2$  are continuous.

**Proof.** Let  $\pi_1 : X \times Y \rightarrow X$  and  $\pi_2 : X \times Y \rightarrow Y$ . Then for  $U$  open in  $X$  and  $V$  open in  $Y$ , we have  $\pi_1^{-1}(U) = U \times T$  and  $\pi_2^{-1}(V) = X \times V$  open in  $X \times Y$  (by the definition of product topology; these are basis elements for the product topology on  $X \times Y$ ). So  $\pi_1$  and  $\pi_2$  are continuous. Note that for each  $a \in A$ ,  $\pi_1(f(a)) = \pi_1((f_1(a), f_2(a))) = f_1(a)$  and  $\pi_2(f(a)) = \pi_2((f_1(a), f_2(a))) = f_2(a)$ . So  $f_1 = \pi_1 \circ f$  and  $f_2 = \pi_2 \circ f$ . Suppose  $f$  is continuous. Then, by Theorem 18.2 part (c),  $f_1$  and  $f_2$  are continuous.

# Theorem 18.4

## Theorem 18.4. Maps into Products.

Let  $f : A \rightarrow X \times Y$  be given by the equation  $f(a) = (f_1(a), f_2(a))$  where  $f_1 : A \rightarrow X$  and  $f_2 : A \rightarrow Y$ . Then  $f$  is continuous if and only if the functions  $f_1$  and  $f_2$  are continuous.

**Proof.** Let  $\pi_1 : X \times Y \rightarrow X$  and  $\pi_2 : X \times Y \rightarrow Y$ . Then for  $U$  open in  $X$  and  $V$  open in  $Y$ , we have  $\pi_1^{-1}(U) = U \times T$  and  $\pi_2^{-1}(V) = X \times V$  open in  $X \times Y$  (by the definition of product topology; these are basis elements for the product topology on  $X \times Y$ ). So  $\pi_1$  and  $\pi_2$  are continuous. Note that for each  $a \in A$ ,  $\pi_1(f(a)) = \pi_1((f_1(a), f_2(a))) = f_1(a)$  and  $\pi_2(f(a)) = \pi_2((f_1(a), f_2(a))) = f_2(a)$ . So  $f_1 = \pi_1 \circ f$  and  $f_2 = \pi_2 \circ f$ . Suppose  $f$  is continuous. Then, by Theorem 18.2 part (c),  $f_1$  and  $f_2$  are continuous.

## Theorem 18.4 (continued)

### Theorem 18.4. Maps into Products.

Let  $f : A \rightarrow X \times Y$  be given by the equation  $f(a) = (f_1(a), f_2(a))$  where  $f_1 : A \rightarrow X$  and  $f_2 : A \rightarrow Y$ . Then  $f$  is continuous if and only if the functions  $f_1$  and  $f_2$  are continuous.

**Proof (continued).** Suppose  $f_1$  and  $f_2$  are continuous. Let  $U \times V$  be a basis element for the product topology of  $X \times Y$  (so  $U$  is open in  $X$  and  $V$  is open in  $Y$ ).

## Theorem 18.4 (continued)

### Theorem 18.4. Maps into Products.

Let  $f : A \rightarrow X \times Y$  be given by the equation  $f(a) = (f_1(a), f_2(a))$  where  $f_1 : A \rightarrow X$  and  $f_2 : A \rightarrow Y$ . Then  $f$  is continuous if and only if the functions  $f_1$  and  $f_2$  are continuous.

**Proof (continued).** Suppose  $f_1$  and  $f_2$  are continuous. Let  $U \times V$  be a basis element for the product topology of  $X \times Y$  (so  $U$  is open in  $X$  and  $V$  is open in  $Y$ ). Now  $a \in f^{-1}(U \times V)$  if and only if  $f(a) \in U \times V$ , or if and only if  $f_1(a) \in U$  and  $f_2(a) \in V$ , or if and only if  $a \in f_1^{-1}(U) \cap f_2^{-1}(V)$ . That is,  $f^{-1}(U \times V) = f_1^{-1}(U) \cap f_2^{-1}(V)$ . Since  $f_1$  and  $f_2$  are continuous then  $f_1^{-1}(U)$  and  $f_2^{-1}(V)$  are open in  $X$  and so  $f^{-1}(U \times V)$  is open in  $X$ .

## Theorem 18.4 (continued)

### Theorem 18.4. Maps into Products.

Let  $f : A \rightarrow X \times Y$  be given by the equation  $f(a) = (f_1(a), f_2(a))$  where  $f_1 : A \rightarrow X$  and  $f_2 : A \rightarrow Y$ . Then  $f$  is continuous if and only if the functions  $f_1$  and  $f_2$  are continuous.

**Proof (continued).** Suppose  $f_1$  and  $f_2$  are continuous. Let  $U \times V$  be a basis element for the product topology of  $X \times Y$  (so  $U$  is open in  $X$  and  $V$  is open in  $Y$ ). Now  $a \in f^{-1}(U \times V)$  if and only if  $f(a) \in U \times V$ , or if and only if  $f_1(a) \in U$  and  $f_2(a) \in V$ , or if and only if  $a \in f_1^{-1}(U) \cap f_2^{-1}(V)$ . That is,  $f^{-1}(U \times V) = f_1^{-1}(U) \cap f_2^{-1}(V)$ . Since  $f_1$  and  $f_2$  are continuous then  $f_1^{-1}(U)$  and  $f_2^{-1}(V)$  are open in  $A$  and so  $f^{-1}(U \times V)$  is open in  $A$ . Since every open set in  $X \times Y$  can be written as a union of basis elements by Lemma 13.1, say  $\bigcup_{\alpha \in J} U_\alpha \times V_\alpha$ , and  $f^{-1}(\bigcup_{\alpha \in J} U_\alpha \times V_\alpha) = \bigcup_{\alpha \in J} f^{-1}(U_\alpha \times V_\alpha)$ , then the inverse image of any open set in  $X \times Y$  is open in  $A$ . That is,  $f$  is continuous.

□

## Theorem 18.4 (continued)

### Theorem 18.4. Maps into Products.

Let  $f : A \rightarrow X \times Y$  be given by the equation  $f(a) = (f_1(a), f_2(a))$  where  $f_1 : A \rightarrow X$  and  $f_2 : A \rightarrow Y$ . Then  $f$  is continuous if and only if the functions  $f_1$  and  $f_2$  are continuous.

**Proof (continued).** Suppose  $f_1$  and  $f_2$  are continuous. Let  $U \times V$  be a basis element for the product topology of  $X \times Y$  (so  $U$  is open in  $X$  and  $V$  is open in  $Y$ ). Now  $a \in f^{-1}(U \times V)$  if and only if  $f(a) \in U \times V$ , or if and only if  $f_1(a) \in U$  and  $f_2(a) \in V$ , or if and only if  $a \in f_1^{-1}(U) \cap f_2^{-1}(V)$ . That is,  $f^{-1}(U \times V) = f_1^{-1}(U) \cap f_2^{-1}(V)$ . Since  $f_1$  and  $f_2$  are continuous then  $f_1^{-1}(U)$  and  $f_2^{-1}(V)$  are open in  $A$  and so  $f^{-1}(U \times V)$  is open in  $A$ . Since every open set in  $X \times Y$  can be written as a union of basis elements by Lemma 13.1, say  $\bigcup_{\alpha \in J} U_\alpha \times V_\alpha$ , and  $f^{-1}(\bigcup_{\alpha \in J} U_\alpha \times V_\alpha) = \bigcup_{\alpha \in J} f^{-1}(U_\alpha \times V_\alpha)$ , then the inverse image of any open set in  $X \times Y$  is open in  $A$ . That is,  $f$  is continuous.

