


# Introduction to Topology

## Chapter 7. Complete Metric Spaces and Function Spaces

### Section 43. Complete Metric Spaces—Proofs of Theorems



# Table of contents

- 1 Lemma 43.1
- 2 Theorem 43.2
- 3 Lemma 43.3
- 4 Theorem 43.4
- 5 Theorem 43.5
- 6 Theorem 43.6
- 7 Theorem 43.7

# Lemma 43.1

**Lemma 43.1.** A metric space  $(X, d)$  is complete if every Cauchy sequence in  $X$  has a convergent subsequence.

**Proof.** Let  $(x_n)$  be a Cauchy sequence in  $(X, d)$ . Let  $(x_{n_i})$  be a subsequence of  $(x_n)$  that converges to some  $x \in X$ . Let  $\varepsilon > 0$ .

# Lemma 43.1

**Lemma 43.1.** A metric space  $(X, d)$  is complete if every Cauchy sequence in  $X$  has a convergent subsequence.

**Proof.** Let  $(x_n)$  be a Cauchy sequence in  $(X, d)$ . Let  $(x_{n_i})$  be a subsequence of  $(x_n)$  that converges to some  $x \in X$ . Let  $\varepsilon > 0$ . Then there exists  $N_1 \in \mathbb{N}$  such that  $d(x_n, x_m) < \varepsilon/2$  for all  $n, m \geq N_1$ . Since  $(x_{n_i}) \rightarrow x$ , let  $N_2 \in \mathbb{N}$  such that for  $n_i \geq N_2$  we have  $d(x_{n_i}, x) < \varepsilon$ .

# Lemma 43.1

**Lemma 43.1.** A metric space  $(X, d)$  is complete if every Cauchy sequence in  $X$  has a convergent subsequence.

**Proof.** Let  $(x_n)$  be a Cauchy sequence in  $(X, d)$ . Let  $(x_{n_i})$  be a subsequence of  $(x_n)$  that converges to some  $x \in X$ . Let  $\varepsilon > 0$ . Then there exists  $N_1 \in \mathbb{N}$  such that  $d(x_n, x_m) < \varepsilon/2$  for all  $n, m \geq N_1$ . Since  $(x_{n_i}) \rightarrow x$ , let  $N_2 \in \mathbb{N}$  such that for  $n_i \geq N_2$  we have  $d(x_{n_i}, x) < \varepsilon$ . So with  $n \geq N_1$  and  $n_i \geq N_2$  we have

$$d(x_n, x) \leq d(x_n, x_{n_i}) + d(x_{n_i}, x) < \varepsilon/2 + \varepsilon/2 = \varepsilon.$$

So arbitrary Cauchy sequence  $(x_n)$  converges (to  $x$ ) and  $(X, d)$  is complete. □

# Lemma 43.1

**Lemma 43.1.** A metric space  $(X, d)$  is complete if every Cauchy sequence in  $X$  has a convergent subsequence.

**Proof.** Let  $(x_n)$  be a Cauchy sequence in  $(X, d)$ . Let  $(x_{n_i})$  be a subsequence of  $(x_n)$  that converges to some  $x \in X$ . Let  $\varepsilon > 0$ . Then there exists  $N_1 \in \mathbb{N}$  such that  $d(x_n, x_m) < \varepsilon/2$  for all  $n, m \geq N_1$ . Since  $(x_{n_i}) \rightarrow x$ , let  $N_2 \in \mathbb{N}$  such that for  $n_i \geq N_2$  we have  $d(x_{n_i}, x) < \varepsilon/2$ . So with  $n \geq N_1$  and  $n_i \geq N_2$  we have

$$d(x_n, x) \leq d(x_n, x_{n_i}) + d(x_{n_i}, x) < \varepsilon/2 + \varepsilon/2 = \varepsilon.$$

So arbitrary Cauchy sequence  $(x_n)$  converges (to  $x$ ) and  $(X, d)$  is complete. □

## Theorem 43.2

**Theorem 43.2.** Euclidean space  $\mathbb{R}^k$  (where  $k \in \mathbb{N}$ ) is complete in either of its usual metrics, the Euclidean metric  $d$  or the square metric  $\rho$ .

**Proof.** Let  $(x_n)$  be a Cauchy sequence in  $(\mathbb{R}^k, \rho)$ . Notice that with  $\varepsilon = 1$  there is  $N \in \mathbb{N}$  such that for all  $m, n \geq N$  we have  $\rho(x_n, x_m) \leq 1$ , so

$$M = \max\{\rho(x_1, \mathbf{0}), \rho(x_2, \mathbf{0}), \dots, \rho(x_{N-1}, \mathbf{0}), \rho(x_N, \mathbf{0}) + 1\}$$

is an upper bound for  $\rho(x_n, \mathbf{0})$  for all  $n \in \mathbb{N}$  (that is,  $(x_n)$  is a bounded sequence). So  $(x_n) \subset [-M, M]^k$ .

## Theorem 43.2

**Theorem 43.2.** Euclidean space  $\mathbb{R}^k$  (where  $k \in \mathbb{N}$ ) is complete in either of its usual metrics, the Euclidean metric  $d$  or the square metric  $\rho$ .

**Proof.** Let  $(x_n)$  be a Cauchy sequence in  $(\mathbb{R}^k, \rho)$ . Notice that with  $\varepsilon = 1$  there is  $N \in \mathbb{N}$  such that for all  $m, n \geq N$  we have  $\rho(x_n, x_m) \leq 1$ , so

$$M = \max\{\rho(x_1, \mathbf{0}), \rho(x_2, \mathbf{0}), \dots, \rho(x_{N-1}, \mathbf{0}), \rho(x_N, \mathbf{0}) + 1\}$$

is an upper bound for  $\rho(x_n, \mathbf{0})$  for all  $n \in \mathbb{N}$  (that is,  $(x_n)$  is a bounded sequence). So  $(x_n) \subset [-M, M]^k$ . Now the cube  $[-M, M]^k$  is closed and bounded and so is compact by Theorem 27.3 (The Heine-Borel Theorem) in both  $(\mathbb{R}^k, \rho)$  and  $(\mathbb{R}^k, d)$ . By Theorem 28.2,  $[-M, M]^k$  is sequentially compact and so (by the definition of sequentially compact)  $(x_n)$  has a convergent subsequence; so by Lemma 43.1,  $(\mathbb{R}^k, \rho)$  is complete.



## Theorem 43.2

**Theorem 43.2.** Euclidean space  $\mathbb{R}^k$  (where  $k \in \mathbb{N}$ ) is complete in either of its usual metrics, the Euclidean metric  $d$  or the square metric  $\rho$ .

**Proof.** Let  $(x_n)$  be a Cauchy sequence in  $(\mathbb{R}^k, \rho)$ . Notice that with  $\varepsilon = 1$  there is  $N \in \mathbb{N}$  such that for all  $m, n \geq N$  we have  $\rho(x_n, x_m) \leq 1$ , so

$$M = \max\{\rho(x_1, \mathbf{0}), \rho(x_2, \mathbf{0}), \dots, \rho(x_{N-1}, \mathbf{0}), \rho(x_N, \mathbf{0}) + 1\}$$

is an upper bound for  $\rho(x_n, \mathbf{0})$  for all  $n \in \mathbb{N}$  (that is,  $(x_n)$  is a bounded sequence). So  $(x_n) \subset [-M, M]^k$ . Now the cube  $[-M, M]^k$  is closed and bounded and so is compact by Theorem 27.3 (The Heine-Borel Theorem) in both  $(\mathbb{R}^k, \rho)$  and  $(\mathbb{R}^k, d)$ . By Theorem 28.2,  $[-M, M]^k$  is sequentially compact and so (by the definition of sequentially compact)  $(x_n)$  has a convergent subsequence; so by Lemma 43.1,  $(\mathbb{R}^k, \rho)$  is complete.

By Theorem 20.3,  $\rho$  and  $d$  induce the same topology on  $\mathbb{R}^k$  (namely, the product topology) so a sequence is Cauchy (or convergent) relative to  $\rho$  if and only if it is Cauchy (or convergent, respectively) relative to  $d$ . So the same argument given above shows that  $(\mathbb{R}^k, d)$  is also complete.  $\square$

## Theorem 43.2

**Theorem 43.2.** Euclidean space  $\mathbb{R}^k$  (where  $k \in \mathbb{N}$ ) is complete in either of its usual metrics, the Euclidean metric  $d$  or the square metric  $\rho$ .

**Proof.** Let  $(x_n)$  be a Cauchy sequence in  $(\mathbb{R}^k, \rho)$ . Notice that with  $\varepsilon = 1$  there is  $N \in \mathbb{N}$  such that for all  $m, n \geq N$  we have  $\rho(x_n, x_m) \leq 1$ , so

$$M = \max\{\rho(x_1, \mathbf{0}), \rho(x_2, \mathbf{0}), \dots, \rho(x_{N-1}, \mathbf{0}), \rho(x_N, \mathbf{0}) + 1\}$$

is an upper bound for  $\rho(x_n, \mathbf{0})$  for all  $n \in \mathbb{N}$  (that is,  $(x_n)$  is a bounded sequence). So  $(x_n) \subset [-M, M]^k$ . Now the cube  $[-M, M]^k$  is closed and bounded and so is compact by Theorem 27.3 (The Heine-Borel Theorem) in both  $(\mathbb{R}^k, \rho)$  and  $(\mathbb{R}^k, d)$ . By Theorem 28.2,  $[-M, M]^k$  is sequentially compact and so (by the definition of sequentially compact)  $(x_n)$  has a convergent subsequence; so by Lemma 43.1,  $(\mathbb{R}^k, \rho)$  is complete.

By Theorem 20.3,  $\rho$  and  $d$  induce the same topology on  $\mathbb{R}^k$  (namely, the product topology) so a sequence is Cauchy (or convergent) relative to  $\rho$  if and only if it is Cauchy (or convergent, respectively) relative to  $d$ . So the same argument given above shows that  $(\mathbb{R}^k, d)$  is also complete.  $\square$

## Lemma 43.3

**Lemma 43.3.** Let  $X$  be the product space  $X = \prod_{\alpha \in J} X_\alpha$  (under the product topology) and let  $(\mathbf{x}_n)$  be a sequence of points in  $X$ . Then  $\mathbf{x}_n \rightarrow \mathbf{x}$  if and only if  $\pi_\alpha(\mathbf{x}_n) \rightarrow \pi_\alpha(\mathbf{x})$  for all  $\alpha \in J$ .

**Proof.** Suppose  $\mathbf{x}_n \rightarrow \mathbf{x}$ . Each projection  $\pi_\alpha$  is continuous (see the proof of Theorem 19.6), so for all  $\alpha \in J$ ,

$$\begin{aligned}\lim_{n \rightarrow \infty} \pi_\alpha(\mathbf{x}_n) &= \pi_\alpha \left( \lim_{n \rightarrow \infty} \mathbf{x}_n \right) \text{ by Theorem 21.3} \\ &= \pi_\alpha(\mathbf{x}).\end{aligned}$$

## Lemma 43.3

**Lemma 43.3.** Let  $X$  be the product space  $X = \prod_{\alpha \in J} X_\alpha$  (under the product topology) and let  $(\mathbf{x}_n)$  be a sequence of points in  $X$ . Then  $\mathbf{x}_n \rightarrow \mathbf{x}$  if and only if  $\pi_\alpha(\mathbf{x}_n) \rightarrow \pi_\alpha(\mathbf{x})$  for all  $\alpha \in J$ .

**Proof.** Suppose  $\mathbf{x}_n \rightarrow \mathbf{x}$ . Each projection  $\pi_\alpha$  is continuous (see the proof of Theorem 19.6), so for all  $\alpha \in J$ ,

$$\begin{aligned} \lim_{n \rightarrow \infty} \pi_\alpha(\mathbf{x}_n) &= \pi_\alpha \left( \lim_{n \rightarrow \infty} \mathbf{x}_n \right) \text{ by Theorem 21.3} \\ &= \pi_\alpha(\mathbf{x}). \end{aligned}$$

Suppose  $\pi_\alpha(\mathbf{x}_n) \rightarrow \pi_\alpha(\mathbf{x})$  for all  $\alpha \in J$ . Let  $U = \prod_{\alpha \in J} U_\alpha$  be a basis element for  $X$  in the product topology which contains  $\mathbf{x}$  (so by Theorem 19.1,  $U_\alpha = X_\alpha$  for all but finitely many  $\alpha \in J$ ).

## Lemma 43.3

**Lemma 43.3.** Let  $X$  be the product space  $X = \prod_{\alpha \in J} X_\alpha$  (under the product topology) and let  $(\mathbf{x}_n)$  be a sequence of points in  $X$ . Then  $\mathbf{x}_n \rightarrow \mathbf{x}$  if and only if  $\pi_\alpha(\mathbf{x}_n) \rightarrow \pi_\alpha(\mathbf{x})$  for all  $\alpha \in J$ .

**Proof.** Suppose  $\mathbf{x}_n \rightarrow \mathbf{x}$ . Each projection  $\pi_\alpha$  is continuous (see the proof of Theorem 19.6), so for all  $\alpha \in J$ ,

$$\begin{aligned} \lim_{n \rightarrow \infty} \pi_\alpha(\mathbf{x}_n) &= \pi_\alpha \left( \lim_{n \rightarrow \infty} \mathbf{x}_n \right) \text{ by Theorem 21.3} \\ &= \pi_\alpha(\mathbf{x}). \end{aligned}$$

Suppose  $\pi_\alpha(\mathbf{x}_n) \rightarrow \pi_\alpha(\mathbf{x})$  for all  $\alpha \in J$ . Let  $U = \prod_{\alpha \in J} U_\alpha$  be a basis element for  $X$  in the product topology which contains  $\mathbf{x}$  (so by Theorem 19.1,  $U_\alpha = X_\alpha$  for all but finitely many  $\alpha \in J$ ). For each  $\alpha \in J$  for which  $U_\alpha \neq X_\alpha$ , choose  $N_\alpha \in \mathbb{N}$  such that  $\pi_\alpha(\mathbf{x}_n) \in U_\alpha$  for all  $n \geq N_\alpha$  (such  $N_\alpha \in \mathbb{N}$  exists since  $\pi_\alpha(\mathbf{x}_n) \rightarrow \pi_\alpha(\mathbf{x})$ ). Let  $N$  be the largest of the  $N_\alpha$  (since there are only finitely many  $N_\alpha$ , there is a largest).

## Lemma 43.3

**Lemma 43.3.** Let  $X$  be the product space  $X = \prod_{\alpha \in J} X_\alpha$  (under the product topology) and let  $(\mathbf{x}_n)$  be a sequence of points in  $X$ . Then  $\mathbf{x}_n \rightarrow \mathbf{x}$  if and only if  $\pi_\alpha(\mathbf{x}_n) \rightarrow \pi_\alpha(\mathbf{x})$  for all  $\alpha \in J$ .

**Proof.** Suppose  $\mathbf{x}_n \rightarrow \mathbf{x}$ . Each projection  $\pi_\alpha$  is continuous (see the proof of Theorem 19.6), so for all  $\alpha \in J$ ,

$$\begin{aligned} \lim_{n \rightarrow \infty} \pi_\alpha(\mathbf{x}_n) &= \pi_\alpha \left( \lim_{n \rightarrow \infty} \mathbf{x}_n \right) \text{ by Theorem 21.3} \\ &= \pi_\alpha(\mathbf{x}). \end{aligned}$$

Suppose  $\pi_\alpha(\mathbf{x}_n) \rightarrow \pi_\alpha(\mathbf{x})$  for all  $\alpha \in J$ . Let  $U = \prod_{\alpha \in J} U_\alpha$  be a basis element for  $X$  in the product topology which contains  $\mathbf{x}$  (so by Theorem 19.1,  $U_\alpha = X_\alpha$  for all but finitely many  $\alpha \in J$ ). For each  $\alpha \in J$  for which  $U_\alpha \neq X_\alpha$ , choose  $N_\alpha \in \mathbb{N}$  such that  $\pi_\alpha(\mathbf{x}_n) \in U_\alpha$  for all  $n \geq N_\alpha$  (such  $N_\alpha \in \mathbb{N}$  exists since  $\pi_\alpha(\mathbf{x}_n) \rightarrow \pi_\alpha(\mathbf{x})$ ). Let  $N$  be the largest of the  $N_\alpha$  (since there are only finitely many  $N_\alpha$ , there is a largest). Then for all  $n \geq N$ ,  $\mathbf{x}_n \in U$ . Therefore, since  $U$  is an arbitrary basis element, then  $\mathbf{x}_n \rightarrow \mathbf{x}$ . □

# Lemma 43.3

**Lemma 43.3.** Let  $X$  be the product space  $X = \prod_{\alpha \in J} X_\alpha$  (under the product topology) and let  $(\mathbf{x}_n)$  be a sequence of points in  $X$ . Then  $\mathbf{x}_n \rightarrow \mathbf{x}$  if and only if  $\pi_\alpha(\mathbf{x}_n) \rightarrow \pi_\alpha(\mathbf{x})$  for all  $\alpha \in J$ .

**Proof.** Suppose  $\mathbf{x}_n \rightarrow \mathbf{x}$ . Each projection  $\pi_\alpha$  is continuous (see the proof of Theorem 19.6), so for all  $\alpha \in J$ ,

$$\begin{aligned} \lim_{n \rightarrow \infty} \pi_\alpha(\mathbf{x}_n) &= \pi_\alpha \left( \lim_{n \rightarrow \infty} \mathbf{x}_n \right) \text{ by Theorem 21.3} \\ &= \pi_\alpha(\mathbf{x}). \end{aligned}$$

Suppose  $\pi_\alpha(\mathbf{x}_n) \rightarrow \pi_\alpha(\mathbf{x})$  for all  $\alpha \in J$ . Let  $U = \prod_{\alpha \in J} U_\alpha$  be a basis element for  $X$  in the product topology which contains  $\mathbf{x}$  (so by Theorem 19.1,  $U_\alpha = X_\alpha$  for all but finitely many  $\alpha \in J$ ). For each  $\alpha \in J$  for which  $U_\alpha \neq X_\alpha$ , choose  $N_\alpha \in \mathbb{N}$  such that  $\pi_\alpha(\mathbf{x}_n) \in U_\alpha$  for all  $n \geq N_\alpha$  (such  $N_\alpha \in \mathbb{N}$  exists since  $\pi_\alpha(\mathbf{x}_n) \rightarrow \pi_\alpha(\mathbf{x})$ ). Let  $N$  be the largest of the  $N_\alpha$  (since there are only finitely many  $N_\alpha$ , there is a largest). Then for all  $n \geq N$ ,  $\mathbf{x}_n \in U$ . Therefore, since  $U$  is an arbitrary basis element, then  $\mathbf{x}_n \rightarrow \mathbf{x}$ . □

## Theorem 43.4

**Theorem 43.4.** There is a metric for the product space  $\mathbb{R}^\omega$  relative to which  $\mathbb{R}^\omega$  is complete.

**Proof.** Consider  $D(\mathbf{x}, \mathbf{y}) = \sup_{i \in \mathbb{N}} \{\bar{d}(x_i, y_i)/i\}$  where  $\bar{d}$  is the standard bounded metric on  $\mathbb{R}$ . Then  $D$  is a metric on  $\mathbb{R}^\omega$  (see Section 20) and  $D$  induces the product topology by Theorem 20.5.



## Theorem 43.4

**Theorem 43.4.** There is a metric for the product space  $\mathbb{R}^\omega$  relative to which  $\mathbb{R}^\omega$  is complete.

**Proof.** Consider  $D(\mathbf{x}, \mathbf{y}) = \sup_{i \in \mathbb{N}} \{\bar{d}(x_i, y_i)/i\}$  where  $\bar{d}$  is the standard bounded metric on  $\mathbb{R}$ . Then  $D$  is a metric on  $\mathbb{R}^\omega$  (see Section 20) and  $D$  induces the product topology by Theorem 20.5. Now let  $(\mathbf{x}_n)$  be a Cauchy sequence in  $(\mathbb{R}^\omega, D)$ . For all  $i \in \mathbb{N}$ ,

$$\bar{d}(x_i, y_i)/i = \bar{d}(\pi_i(\mathbf{x}), \pi_i(\mathbf{y}))/i \leq \sup_{i \in \mathbb{N}} \{\bar{d}(x_i, y_i)/i\} = D(\mathbf{x}, \mathbf{y}),$$

so for all  $i \in \mathbb{N}$  we have  $\bar{d}(\pi_i(\mathbf{x}), \pi_i(\mathbf{y})) \leq iD(\mathbf{x}, \mathbf{y})$ .

# Theorem 43.4

**Theorem 43.4.** There is a metric for the product space  $\mathbb{R}^\omega$  relative to which  $\mathbb{R}^\omega$  is complete.

**Proof.** Consider  $D(\mathbf{x}, \mathbf{y}) = \sup_{i \in \mathbb{N}} \{\bar{d}(x_i, y_i)/i\}$  where  $\bar{d}$  is the standard bounded metric on  $\mathbb{R}$ . Then  $D$  is a metric on  $\mathbb{R}^\omega$  (see Section 20) and  $D$  induces the product topology by Theorem 20.5. Now let  $(\mathbf{x}_n)$  be a Cauchy sequence in  $(\mathbb{R}^\omega, D)$ . For all  $i \in \mathbb{N}$ ,

$$\bar{d}(x_i, y_i)/i = \bar{d}(\pi_i(\mathbf{x}), \pi_i(\mathbf{y}))/i \leq \sup_{i \in \mathbb{N}} \{\bar{d}(x_i, y_i)/i\} = D(\mathbf{x}, \mathbf{y}),$$

so for all  $i \in \mathbb{N}$  we have  $\bar{d}(\pi_i(\mathbf{x}), \pi_i(\mathbf{y})) \leq iD(\mathbf{x}, \mathbf{y})$ . So for a fixed  $i$ , since  $(\mathbf{x}_n)$  is Cauchy, we have for all  $\varepsilon > 0$  that there exists  $N \in \mathbb{N}$  such that for  $m, n \geq N$  we have  $\bar{d}(\pi_i(\mathbf{x}_n), \pi_i(\mathbf{x}_m)) \leq iD(\mathbf{x}_n, \mathbf{x}_m) < i(\varepsilon/i) = \varepsilon$ , and so  $(\pi_i(\mathbf{x}_n))$  is a Cauchy sequence in  $\mathbb{R}$ . So  $\pi_i(\mathbf{x}_n) \rightarrow a_i$ .

# Theorem 43.4

**Theorem 43.4.** There is a metric for the product space  $\mathbb{R}^\omega$  relative to which  $\mathbb{R}^\omega$  is complete.

**Proof.** Consider  $D(\mathbf{x}, \mathbf{y}) = \sup_{i \in \mathbb{N}} \{\bar{d}(x_i, y_i)/i\}$  where  $\bar{d}$  is the standard bounded metric on  $\mathbb{R}$ . Then  $D$  is a metric on  $\mathbb{R}^\omega$  (see Section 20) and  $D$  induces the product topology by Theorem 20.5. Now let  $(\mathbf{x}_n)$  be a Cauchy sequence in  $(\mathbb{R}^\omega, D)$ . For all  $i \in \mathbb{N}$ ,

$$\bar{d}(x_i, y_i)/i = \bar{d}(\pi_i(\mathbf{x}), \pi_i(\mathbf{y}))/i \leq \sup_{i \in \mathbb{N}} \{\bar{d}(x_i, y_i)/i\} = D(\mathbf{x}, \mathbf{y}),$$

so for all  $i \in \mathbb{N}$  we have  $\bar{d}(\pi_i(\mathbf{x}_n), \pi_i(\mathbf{x}_m)) \leq iD(\mathbf{x}_n, \mathbf{x}_m)$ . So for a fixed  $i$ , since  $(\mathbf{x}_n)$  is Cauchy, we have for all  $\varepsilon > 0$  that there exists  $N \in \mathbb{N}$  such that for  $m, n \geq N$  we have  $\bar{d}(\pi_i(\mathbf{x}_n), \pi_i(\mathbf{x}_m)) \leq iD(\mathbf{x}_n, \mathbf{x}_m) < i(\varepsilon/i) = \varepsilon$ , and so  $(\pi_i(\mathbf{x}_n))$  is a Cauchy sequence in  $\mathbb{R}$ . So  $\pi_i(\mathbf{x}_n) \rightarrow a_i$ . Consider  $\mathbf{a} = (a_1, a_2, \dots) \in \mathbb{R}^\omega$ . Then by Lemma 43.3,  $\mathbf{x}_n \rightarrow \mathbf{a}$  and so  $(\mathbb{R}^\omega, D)$  is complete. □

# Theorem 43.4

**Theorem 43.4.** There is a metric for the product space  $\mathbb{R}^\omega$  relative to which  $\mathbb{R}^\omega$  is complete.

**Proof.** Consider  $D(\mathbf{x}, \mathbf{y}) = \sup_{i \in \mathbb{N}} \{\bar{d}(x_i, y_i)/i\}$  where  $\bar{d}$  is the standard bounded metric on  $\mathbb{R}$ . Then  $D$  is a metric on  $\mathbb{R}^\omega$  (see Section 20) and  $D$  induces the product topology by Theorem 20.5. Now let  $(\mathbf{x}_n)$  be a Cauchy sequence in  $(\mathbb{R}^\omega, D)$ . For all  $i \in \mathbb{N}$ ,

$$\bar{d}(x_i, y_i)/i = \bar{d}(\pi_i(\mathbf{x}), \pi_i(\mathbf{y}))/i \leq \sup_{i \in \mathbb{N}} \{\bar{d}(x_i, y_i)/i\} = D(\mathbf{x}, \mathbf{y}),$$

so for all  $i \in \mathbb{N}$  we have  $\bar{d}(\pi_i(\mathbf{x}), \pi_i(\mathbf{y})) \leq iD(\mathbf{x}, \mathbf{y})$ . So for a fixed  $i$ , since  $(\mathbf{x}_n)$  is Cauchy, we have for all  $\varepsilon > 0$  that there exists  $N \in \mathbb{N}$  such that for  $m, n \geq N$  we have  $\bar{d}(\pi_i(\mathbf{x}_n), \pi_i(\mathbf{x}_m)) \leq iD(\mathbf{x}_n, \mathbf{x}_m) < i(\varepsilon/i) = \varepsilon$ , and so  $(\pi_i(\mathbf{x}_n))$  is a Cauchy sequence in  $\mathbb{R}$ . So  $\pi_i(\mathbf{x}_n) \rightarrow a_i$ . Consider  $\mathbf{a} = (a_1, a_2, \dots) \in \mathbb{R}^\omega$ . Then by Lemma 43.3,  $\mathbf{x}_n \rightarrow \mathbf{a}$  and so  $(\mathbb{R}^\omega, D)$  is complete. □

# Theorem 43.5

**Theorem 43.5.** If the space  $Y$  is complete in the metric  $d$ , then the space  $Y^J$  is complete in the uniform metric  $\bar{\rho}$  corresponding to  $d$ .

**Proof.** If  $(Y, d)$  is complete then  $(Y, \bar{d})$  is complete (see the Note in the class notes before Lemma 43.1). Let  $(f_n)$  be a Cauchy sequence in  $(Y, \bar{\rho})$ .

# Theorem 43.5

**Theorem 43.5.** If the space  $Y$  is complete in the metric  $d$ , then the space  $Y^J$  is complete in the uniform metric  $\bar{\rho}$  corresponding to  $d$ .

**Proof.** If  $(Y, d)$  is complete then  $(Y, \bar{d})$  is complete (see the Note in the class notes before Lemma 43.1). Let  $(f_n)$  be a Cauchy sequence in  $(Y, \bar{\rho})$ . For any  $\alpha \in J$  we have

$$\bar{d}(f_n(\alpha), f_m(\alpha)) \leq \sup\{\bar{d}(f_n(\alpha), f_m(\alpha)) \mid \alpha \in J\} = \bar{\rho}(f_n, f_m), \quad (*)$$

so  $(f_n(\alpha))$  is a Cauchy sequence in  $(Y, \bar{d})$ .

## Theorem 43.5

**Theorem 43.5.** If the space  $Y$  is complete in the metric  $d$ , then the space  $Y^J$  is complete in the uniform metric  $\bar{\rho}$  corresponding to  $d$ .

**Proof.** If  $(Y, d)$  is complete then  $(Y, \bar{d})$  is complete (see the Note in the class notes before Lemma 43.1). Let  $(f_n)$  be a Cauchy sequence in  $(Y, \bar{\rho})$ . For any  $\alpha \in J$  we have

$$\bar{d}(f_n(\alpha), f_m(\alpha)) \leq \sup\{\bar{d}(f_n(\alpha), f_m(\alpha)) \mid \alpha \in J\} = \bar{\rho}(f_n, f_m), \quad (*)$$

so  $(f_n(\alpha))$  is a Cauchy sequence in  $(Y, \bar{d})$ . Since  $(Y, \bar{d})$  is complete, then there is  $y_\alpha \in Y$  such that  $f_n(\alpha) \rightarrow y_\alpha$  with respect to  $\bar{d}$ . Define  $f : J \rightarrow Y$  as  $f(\alpha) = y_\alpha$ ; so  $f \in Y^J$ . We next show that  $f_n \rightarrow f$  with respect to  $\bar{\rho}$ .

## Theorem 43.5

**Theorem 43.5.** If the space  $Y$  is complete in the metric  $d$ , then the space  $Y^J$  is complete in the uniform metric  $\bar{\rho}$  corresponding to  $d$ .

**Proof.** If  $(Y, d)$  is complete then  $(Y, \bar{d})$  is complete (see the Note in the class notes before Lemma 43.1). Let  $(f_n)$  be a Cauchy sequence in  $(Y, \bar{\rho})$ . For any  $\alpha \in J$  we have

$$\bar{d}(f_n(\alpha), f_m(\alpha)) \leq \sup\{\bar{d}(f_n(\alpha), f_m(\alpha)) \mid \alpha \in J\} = \bar{\rho}(f_n, f_m), \quad (*)$$

so  $(f_n(\alpha))$  is a Cauchy sequence in  $(Y, \bar{d})$ . Since  $(Y, \bar{d})$  is complete, then there is  $y_\alpha \in Y$  such that  $f_n(\alpha) \rightarrow y_\alpha$  with respect to  $\bar{d}$ . Define  $f : J \rightarrow Y$  as  $f(\alpha) = y_\alpha$ ; so  $f \in Y^J$ . We next show that  $f_n \rightarrow f$  with respect to  $\bar{\rho}$ .



## Theorem 43.5 (continued)

**Theorem 43.5.** If the space  $Y$  is complete in the metric  $d$ , then the space  $Y^J$  is complete in the uniform metric  $\bar{\rho}$  corresponding to  $d$ .

**Proof (continued).** Let  $\varepsilon > 0$ . There is  $N_1 \in \mathbb{N}$  such that for all  $m, n \geq N_1$  we have  $\bar{\rho}(f_n, f_m) < \varepsilon/2$  since  $(f_n)$  is Cauchy with respect to  $\bar{\rho}$ . So by (\*), for all  $\alpha \in J$ ,  $\bar{d}(f_n(\alpha), f_m(\alpha)) < \varepsilon/4$  whenever  $m, n \geq N_1$ . Since  $f_m(\alpha) \rightarrow y_\alpha = f(\alpha)$  with respect to  $\bar{d}$  then there is  $N_2 \in \mathbb{N}$  such that for all  $m \geq N_2$  we have  $\bar{d}(f_m(\alpha), f(\alpha)) < \varepsilon/4$ .

## Theorem 43.5 (continued)

**Theorem 43.5.** If the space  $Y$  is complete in the metric  $d$ , then the space  $Y^J$  is complete in the uniform metric  $\bar{\rho}$  corresponding to  $d$ .

**Proof (continued).** Let  $\varepsilon > 0$ . There is  $N_1 \in \mathbb{N}$  such that for all  $m, n \geq N_1$  we have  $\bar{\rho}(f_n, f_m) < \varepsilon/2$  since  $(f_n)$  is Cauchy with respect to  $\bar{\rho}$ . So by (\*), for all  $\alpha \in J$ ,  $\bar{d}(f_n(\alpha), f_m(\alpha)) < \varepsilon/4$  whenever  $m, n \geq N_1$ . Since  $f_m(\alpha) \rightarrow y_\alpha = f(\alpha)$  with respect to  $\bar{d}$  then there is  $N_2 \in \mathbb{N}$  such that for all  $m \geq N_2$  we have  $\bar{d}(f_m(\alpha), f(\alpha)) < \varepsilon/4$ . So for given  $\alpha \in J$ , with  $n \geq N_1$  and  $m \geq N_2$  we have

$$\bar{d}(f_n(\alpha), f(\alpha)) \leq \bar{d}(f_n(\alpha), f_m(\alpha)) + \bar{d}(f_m(\alpha), f(\alpha)) < \varepsilon/4 + \varepsilon/4 = \varepsilon/2.$$

That is, for all  $\alpha \in J$ , if  $n \geq N_1$  then  $\bar{d}(f_n(\alpha), f(\alpha)) < \varepsilon/2$ .

## Theorem 43.5 (continued)

**Theorem 43.5.** If the space  $Y$  is complete in the metric  $d$ , then the space  $Y^J$  is complete in the uniform metric  $\bar{\rho}$  corresponding to  $d$ .

**Proof (continued).** Let  $\varepsilon > 0$ . There is  $N_1 \in \mathbb{N}$  such that for all  $m, n \geq N_1$  we have  $\bar{\rho}(f_n, f_m) < \varepsilon/2$  since  $(f_n)$  is Cauchy with respect to  $\bar{\rho}$ . So by (\*), for all  $\alpha \in J$ ,  $\bar{d}(f_n(\alpha), f_m(\alpha)) < \varepsilon/4$  whenever  $m, n \geq N_1$ . Since  $f_m(\alpha) \rightarrow y_\alpha = f(\alpha)$  with respect to  $\bar{d}$  then there is  $N_2 \in \mathbb{N}$  such that for all  $m \geq N_2$  we have  $\bar{d}(f_m(\alpha), f(\alpha)) < \varepsilon/4$ . So for given  $\alpha \in J$ , with  $n \geq N_1$  and  $m \geq N_2$  we have

$$\bar{d}(f_n(\alpha), f(\alpha)) \leq \bar{d}(f_n(\alpha), f_m(\alpha)) + \bar{d}(f_m(\alpha), f(\alpha)) < \varepsilon/4 + \varepsilon/4 = \varepsilon/2.$$

That is, for all  $\alpha \in J$ , if  $n \geq N_1$  then  $\bar{d}(f_n(\alpha), f(\alpha)) < \varepsilon/2$ . Hence, for all  $n \geq N_1$  we have  $\bar{\rho}(f_n, f) = \sup\{\bar{d}(f_n(\alpha), f(\alpha)) \mid \alpha \in J\} \leq \varepsilon/2$ , and so  $(f_n) \rightarrow f$  with respect to  $\bar{\rho}$ . Since  $(f_n)$  is an arbitrary Cauchy sequence in  $(Y^J, \bar{\rho})$ , the  $(Y^J, \bar{\rho})$  is complete.  $\square$

## Theorem 43.5 (continued)

**Theorem 43.5.** If the space  $Y$  is complete in the metric  $d$ , then the space  $Y^J$  is complete in the uniform metric  $\bar{\rho}$  corresponding to  $d$ .

**Proof (continued).** Let  $\varepsilon > 0$ . There is  $N_1 \in \mathbb{N}$  such that for all  $m, n \geq N_1$  we have  $\bar{\rho}(f_n, f_m) < \varepsilon/2$  since  $(f_n)$  is Cauchy with respect to  $\bar{\rho}$ . So by (\*), for all  $\alpha \in J$ ,  $\bar{d}(f_n(\alpha), f_m(\alpha)) < \varepsilon/4$  whenever  $m, n \geq N_1$ . Since  $f_m(\alpha) \rightarrow y_\alpha = f(\alpha)$  with respect to  $\bar{d}$  then there is  $N_2 \in \mathbb{N}$  such that for all  $m \geq N_2$  we have  $\bar{d}(f_m(\alpha), f(\alpha)) < \varepsilon/4$ . So for given  $\alpha \in J$ , with  $n \geq N_1$  and  $m \geq N_2$  we have

$$\bar{d}(f_n(\alpha), f(\alpha)) \leq \bar{d}(f_n(\alpha), f_m(\alpha)) + \bar{d}(f_m(\alpha), f(\alpha)) < \varepsilon/4 + \varepsilon/4 = \varepsilon/2.$$

That is, for all  $\alpha \in J$ , if  $n \geq N_1$  then  $\bar{d}(f_n(\alpha), f(\alpha)) < \varepsilon/2$ . Hence, for all  $n \geq N_1$  we have  $\bar{\rho}(f_n, f) = \sup\{\bar{d}(f_n(\alpha), f(\alpha)) \mid \alpha \in J\} \leq \varepsilon/2$ , and so  $(f_n) \rightarrow f$  with respect to  $\bar{\rho}$ . Since  $(f_n)$  is an arbitrary Cauchy sequence in  $(Y^J, \bar{\rho})$ , the  $(Y^J, \bar{\rho})$  is complete.  $\square$

## Theorem 43.6

**Theorem 43.6.** Let  $X$  be a topological space and let  $(Y, d)$  be a metric space. The set  $\mathcal{C}(X, Y)$  of continuous functions is closed in  $Y^X$  under the uniform metric. So is the set  $\mathcal{B}(X, Y)$  of bounded functions. Therefore, if  $Y$  is a complete metric space, then both  $\mathcal{C}(X, Y)$  and  $\mathcal{B}(X, Y)$  are complete metric spaces under the uniform metric.

**Proof.** Let  $(f_n) \rightarrow f$  in  $Y^X$  relative to  $\bar{\rho}$ . Let  $\varepsilon > 0$ . Then there exists  $N \in \mathbb{N}$  such that for all  $n \geq N$  we have  $\bar{\rho}(f_n, f) < \varepsilon$ .

## Theorem 43.6

**Theorem 43.6.** Let  $X$  be a topological space and let  $(Y, d)$  be a metric space. The set  $\mathcal{C}(X, Y)$  of continuous functions is closed in  $Y^X$  under the uniform metric. So is the set  $\mathcal{B}(X, Y)$  of bounded functions. Therefore, if  $Y$  is a complete metric space, then both  $\mathcal{C}(X, Y)$  and  $\mathcal{B}(X, Y)$  are complete metric spaces under the uniform metric.

**Proof.** Let  $(f_n) \rightarrow f$  in  $Y^X$  relative to  $\bar{\rho}$ . Let  $\varepsilon > 0$ . Then there exists  $N \in \mathbb{N}$  such that for all  $n \geq N$  we have  $\bar{\rho}(f_n, f) < \varepsilon$ . So for all  $x \in X$  and for all  $n \geq N$  we have

$$\bar{d}(f_n(x), f(x)) \leq \sup\{\bar{d}(f_n(x), f(x)) \mid x \in X\} = \bar{\rho}(f_n, f) < \varepsilon.$$

Therefore  $(f_n)$  converges uniformly to  $f$ .

## Theorem 43.6

**Theorem 43.6.** Let  $X$  be a topological space and let  $(Y, d)$  be a metric space. The set  $\mathcal{C}(X, Y)$  of continuous functions is closed in  $Y^X$  under the uniform metric. So is the set  $\mathcal{B}(X, Y)$  of bounded functions. Therefore, if  $Y$  is a complete metric space, then both  $\mathcal{C}(X, Y)$  and  $\mathcal{B}(X, Y)$  are complete metric spaces under the uniform metric.

**Proof.** Let  $(f_n) \rightarrow f$  in  $Y^X$  relative to  $\bar{\rho}$ . Let  $\varepsilon > 0$ . Then there exists  $N \in \mathbb{N}$  such that for all  $n \geq N$  we have  $\bar{\rho}(f_n, f) < \varepsilon$ . So for all  $x \in X$  and for all  $n \geq N$  we have

$$\bar{d}(f_n(x), f(x)) \leq \sup\{\bar{d}(f_n(x), f(x)) \mid x \in X\} = \bar{\rho}(f_n, f) < \varepsilon.$$

Therefore  $(f_n)$  converges uniformly to  $f$ .

Now we show that  $\mathcal{C}(X, Y)$  is closed in  $Y^X$  relative to  $\bar{\rho}$ . Let  $f \in Y^X$  where  $f$  is a limit point of  $\mathcal{C}(X, Y)$ .

## Theorem 43.6

**Theorem 43.6.** Let  $X$  be a topological space and let  $(Y, d)$  be a metric space. The set  $\mathcal{C}(X, Y)$  of continuous functions is closed in  $Y^X$  under the uniform metric. So is the set  $\mathcal{B}(X, Y)$  of bounded functions. Therefore, if  $Y$  is a complete metric space, then both  $\mathcal{C}(X, Y)$  and  $\mathcal{B}(X, Y)$  are complete metric spaces under the uniform metric.

**Proof.** Let  $(f_n) \rightarrow f$  in  $Y^X$  relative to  $\bar{\rho}$ . Let  $\varepsilon > 0$ . Then there exists  $N \in \mathbb{N}$  such that for all  $n \geq N$  we have  $\bar{\rho}(f_n, f) < \varepsilon$ . So for all  $x \in X$  and for all  $n \geq N$  we have

$$\bar{d}(f_n(x), f(x)) \leq \sup\{\bar{d}(f_n(x), f(x)) \mid x \in X\} = \bar{\rho}(f_n, f) < \varepsilon.$$

Therefore  $(f_n)$  converges uniformly to  $f$ .

Now we show that  $\mathcal{C}(X, Y)$  is closed in  $Y^X$  relative to  $\bar{\rho}$ . Let  $f \in Y^X$  where  $f$  is a limit point of  $\mathcal{C}(X, Y)$ . By Theorem 17.6,  $f$  is a point of the closure of  $\mathcal{C}(X, Y)$  and so by The Sequence Lemma (Lemma 21.2) there is a sequence  $(f_n)$  of elements of  $\mathcal{C}(X, Y)$  which converges to  $f$  relative to  $\bar{\rho}$ .



## Theorem 43.6

**Theorem 43.6.** Let  $X$  be a topological space and let  $(Y, d)$  be a metric space. The set  $\mathcal{C}(X, Y)$  of continuous functions is closed in  $Y^X$  under the uniform metric. So is the set  $\mathcal{B}(X, Y)$  of bounded functions. Therefore, if  $Y$  is a complete metric space, then both  $\mathcal{C}(X, Y)$  and  $\mathcal{B}(X, Y)$  are complete metric spaces under the uniform metric.

**Proof.** Let  $(f_n) \rightarrow f$  in  $Y^X$  relative to  $\bar{\rho}$ . Let  $\varepsilon > 0$ . Then there exists  $N \in \mathbb{N}$  such that for all  $n \geq N$  we have  $\bar{\rho}(f_n, f) < \varepsilon$ . So for all  $x \in X$  and for all  $n \geq N$  we have

$$\bar{d}(f_n(x), f(x)) \leq \sup\{\bar{d}(f_n(x), f(x)) \mid x \in X\} = \bar{\rho}(f_n, f) < \varepsilon.$$

Therefore  $(f_n)$  converges uniformly to  $f$ .

Now we show that  $\mathcal{C}(X, Y)$  is closed in  $Y^X$  relative to  $\bar{\rho}$ . Let  $f \in Y^X$  where  $f$  is a limit point of  $\mathcal{C}(X, Y)$ . By Theorem 17.6,  $f$  is a point of the closure of  $\mathcal{C}(X, Y)$  and so by The Sequence Lemma (Lemma 21.2) there is a sequence  $(f_n)$  of elements of  $\mathcal{C}(X, Y)$  which converges to  $f$  relative to  $\bar{\rho}$ .

## Theorem 43.6 (continued 1)

**Proof (continued).** But as shown above, this means that  $f$  is the uniform limit of  $(f_n)$  and so by the Uniform Limit Theorem (Theorem 21.6)  $f$  is continuous. That is,  $f \in \mathcal{C}(X, Y)$  and since  $f$  is an arbitrary limit point of  $\mathcal{C}(X, Y)$ , then  $\mathcal{C}(X, Y)$  contains all of its limit points and so is closed by Theorem 17.6, as claimed.

Now to show that  $\mathcal{B}(X, Y)$  is closed in  $Y^X$  relative to  $\bar{\rho}$ . Let  $f$  be a limit point of  $\mathcal{B}(X, Y)$ . As above, there is a sequence  $(f_n)$  of elements of  $\mathcal{B}(X, Y)$  where  $(f_n) \rightarrow f$  relative to  $\bar{\rho}$ . So there exists  $N \in \mathbb{N}$  such that  $\bar{\rho}(f_N, f) < 1/2$ .

## Theorem 43.6 (continued 1)

**Proof (continued).** But as shown above, this means that  $f$  is the uniform limit of  $(f_n)$  and so by the Uniform Limit Theorem (Theorem 21.6)  $f$  is continuous. That is,  $f \in \mathcal{C}(X, Y)$  and since  $f$  is an arbitrary limit point of  $\mathcal{C}(X, Y)$ , then  $\mathcal{C}(X, Y)$  contains all of its limit points and so is closed by Theorem 17.6, as claimed.

Now to show that  $\mathcal{B}(X, Y)$  is closed in  $Y^X$  relative to  $\bar{\rho}$ . Let  $f$  be a limit point of  $\mathcal{B}(X, Y)$ . As above, there is a sequence  $(f_n)$  of elements of  $\mathcal{B}(X, Y)$  where  $(f_n) \rightarrow f$  relative to  $\bar{\rho}$ . So there exists  $N \in \mathbb{N}$  such that  $\bar{\rho}(f_N, f) < 1/2$ . Then for all  $x \in X$  we have

$$\bar{d}(f_N(x), f(x)) \leq \sup\{\bar{d}(f_N(x), f(x)) \mid x \in X\} = \bar{\rho}(f_N, f) < 1/2.$$

Since  $\bar{d}(f_N(x), f(x)) = \min\{d(f_N(x), f(x)), 1\}$  this means that for all  $x \in X$  we have  $d(f_N(x), f(x)) < 1/2$ . Let  $M$  be the diameter of the set  $f_N(x)$  (which exists as a finite number since  $f_N$  is bounded) then by the Triangle Inequality for  $d$ , the diameter of  $f(X)$  is at most  $M + 1$ . Hence  $f \in \mathcal{B}(X, Y)$  and, as above,  $\mathcal{B}(X, Y)$  is closed, as claimed.

## Theorem 43.6 (continued 1)

**Proof (continued).** But as shown above, this means that  $f$  is the uniform limit of  $(f_n)$  and so by the Uniform Limit Theorem (Theorem 21.6)  $f$  is continuous. That is,  $f \in \mathcal{C}(X, Y)$  and since  $f$  is an arbitrary limit point of  $\mathcal{C}(X, Y)$ , then  $\mathcal{C}(X, Y)$  contains all of its limit points and so is closed by Theorem 17.6, as claimed.

Now to show that  $\mathcal{B}(X, Y)$  is closed in  $Y^X$  relative to  $\bar{\rho}$ . Let  $f$  be a limit point of  $\mathcal{B}(X, Y)$ . As above, there is a sequence  $(f_n)$  of elements of  $\mathcal{B}(X, Y)$  where  $(f_n) \rightarrow f$  relative to  $\bar{\rho}$ . So there exists  $N \in \mathbb{N}$  such that  $\bar{\rho}(f_N, f) < 1/2$ . Then for all  $x \in X$  we have

$$\bar{d}(f_N(x), f(x)) \leq \sup\{\bar{d}(f_N(x), f(x)) \mid x \in X\} = \bar{\rho}(f_N, f) < 1/2.$$

Since  $\bar{d}(f_N(x), f(x)) = \min\{d(f_N(x), f(x)), 1\}$  this means that for all  $x \in X$  we have  $d(f_N(x), f(x)) < 1/2$ . Let  $M$  be the diameter of the set  $f_N(x)$  (which exists as a finite number since  $f_N$  is bounded) then by the Triangle Inequality for  $d$ , the diameter of  $f(X)$  is at most  $M + 1$ . Hence  $f \in \mathcal{B}(X, Y)$  and, as above,  $\mathcal{B}(X, Y)$  is closed, as claimed.

## Theorem 43.6 (continued 2)

**Theorem 43.6.** Let  $X$  be a topological space and let  $(Y, d)$  be a metric space. The set  $\mathcal{C}(X, Y)$  of continuous functions is closed in  $Y^X$  under the uniform metric. So is the set  $\mathcal{B}(X, Y)$  of bounded functions. Therefore, if  $Y$  is a complete metric space, then both  $\mathcal{C}(X, Y)$  and  $\mathcal{B}(X, Y)$  are complete metric spaces under the uniform metric.

**Proof (continued).** Suppose  $(Y, d)$  is complete. By Theorem 43.5, we know that if  $Y$  is complete in the metric  $d$  then  $Y^X$  is complete in the uniform metric  $\bar{\rho}$ . So if  $(f_n)$  is any Cauchy sequence in either  $\mathcal{C}(X, Y)$  or  $\mathcal{B}(X, Y)$ , then  $(f_n)$  is a Cauchy sequence in  $Y^X$  (since  $\mathcal{C}(X, Y) \subset Y^X$  and  $\mathcal{B}(X, Y) \subset Y^X$ ) and since  $Y^X$  is complete then  $(f_n) \rightarrow f$  for some  $f \in Y^X$ . Then  $f$  is a limit point of both  $\mathcal{C}(X, Y)$  and  $\mathcal{B}(X, Y)$  by The Sequence Lemma (Lemma 21.2), and as shown above,  $f \in \mathcal{C}(X, Y)$  and  $f \in \mathcal{B}(X, Y)$ . Therefore, if  $(Y, d)$  is complete then both  $\mathcal{C}(X, Y)$  and  $\mathcal{B}(X, Y)$  are complete with respect to  $\bar{\rho}$ . □

## Theorem 43.6 (continued 2)

**Theorem 43.6.** Let  $X$  be a topological space and let  $(Y, d)$  be a metric space. The set  $\mathcal{C}(X, Y)$  of continuous functions is closed in  $Y^X$  under the uniform metric. So is the set  $\mathcal{B}(X, Y)$  of bounded functions. Therefore, if  $Y$  is a complete metric space, then both  $\mathcal{C}(X, Y)$  and  $\mathcal{B}(X, Y)$  are complete metric spaces under the uniform metric.

**Proof (continued).** Suppose  $(Y, d)$  is complete. By Theorem 43.5, we know that if  $Y$  is complete in the metric  $d$  then  $Y^X$  is complete in the uniform metric  $\bar{\rho}$ . So if  $(f_n)$  is any Cauchy sequence in either  $\mathcal{C}(X, Y)$  or  $\mathcal{B}(X, Y)$ , then  $(f_n)$  is a Cauchy sequence in  $Y^X$  (since  $\mathcal{C}(X, Y) \subset Y^X$  and  $\mathcal{B}(X, Y) \subset Y^X$ ) and since  $Y^X$  is complete then  $(f_n) \rightarrow f$  for some  $f \in Y^X$ . Then  $f$  is a limit point of both  $\mathcal{C}(X, Y)$  and  $\mathcal{B}(X, Y)$  by The Sequence Lemma (Lemma 21.2), and as shown above,  $f \in \mathcal{C}(X, Y)$  and  $f \in \mathcal{B}(X, Y)$ . Therefore, if  $(Y, d)$  is complete then both  $\mathcal{C}(X, Y)$  and  $\mathcal{B}(X, Y)$  are complete with respect to  $\bar{\rho}$ . □

# Theorem 43.7

**Theorem 43.7.** Let  $(X, d)$  be a metric space. There is an isometric embedding of  $X$  into a complete space.

**Proof.** Let  $\mathcal{B}(X, \mathbb{R})$  be the set of all bounded functions mapping  $X$  into  $\mathbb{R}$ . Let  $x_0 \in X$  be fixed. Given  $a \in X$ , define  $\varphi_a : X \rightarrow \mathbb{R}$  as  $\varphi_a(x) = d(x, a) - d(x, x_0)$ .

# Theorem 43.7

**Theorem 43.7.** Let  $(X, d)$  be a metric space. There is an isometric embedding of  $X$  into a complete space.

**Proof.** Let  $\mathcal{B}(X, \mathbb{R})$  be the set of all bounded functions mapping  $X$  into  $\mathbb{R}$ . Let  $x_0 \in X$  be fixed. Given  $a \in X$ , define  $\varphi_a : X \rightarrow \mathbb{R}$  as  $\varphi_a(x) = d(x, a) - d(x, x_0)$ . For any  $b \in X$  we have by the Triangle Inequality for  $d$  that for all  $x \in X$  we have

$$d(x, a) \leq d(x, b) + d(a, b) \text{ and } d(x, b) \leq d(x, a) + d(a, b)$$

and combining these inequalities gives  $-d(a, b) \leq d(x, a) - d(x, b) \leq d(a, b)$  or

$$|d(x, a) - d(x, b)| \leq d(a, b). \quad (*)$$



# Theorem 43.7

**Theorem 43.7.** Let  $(X, d)$  be a metric space. There is an isometric embedding of  $X$  into a complete space.

**Proof.** Let  $\mathcal{B}(X, \mathbb{R})$  be the set of all bounded functions mapping  $X$  into  $\mathbb{R}$ . Let  $x_0 \in X$  be fixed. Given  $a \in X$ , define  $\varphi_a : X \rightarrow \mathbb{R}$  as  $\varphi_a(x) = d(x, a) - d(x, x_0)$ . For any  $b \in X$  we have by the Triangle Inequality for  $d$  that for all  $x \in X$  we have

$$d(x, a) \leq d(x, b) + d(a, b) \text{ and } d(x, b) \leq d(x, a) + d(a, b)$$

and combining these inequalities gives  $-d(a, b) \leq d(x, a) - d(x, b) \leq d(a, b)$  or

$$|d(x, a) - d(x, b)| \leq d(a, b). \quad (*)$$

With  $b = x_0$ , we then have  $|\varphi_a(x)| \leq d(a, x_0)$  for all  $x \in X$ . Therefore,  $\varphi_a$  is bounded and  $\varphi_a \in \mathcal{B}(X, \mathbb{R})$ .

# Theorem 43.7

**Theorem 43.7.** Let  $(X, d)$  be a metric space. There is an isometric embedding of  $X$  into a complete space.

**Proof.** Let  $\mathcal{B}(X, \mathbb{R})$  be the set of all bounded functions mapping  $X$  into  $\mathbb{R}$ . Let  $x_0 \in X$  be fixed. Given  $a \in X$ , define  $\varphi_a : X \rightarrow \mathbb{R}$  as  $\varphi_a(x) = d(x, a) - d(x, x_0)$ . For any  $b \in X$  we have by the Triangle Inequality for  $d$  that for all  $x \in X$  we have

$$d(x, a) \leq d(x, b) + d(a, b) \text{ and } d(x, b) \leq d(x, a) + d(a, b)$$

and combining these inequalities gives  $-d(a, b) \leq d(x, a) - d(x, b) \leq d(a, b)$  or

$$|d(x, a) - d(x, b)| \leq d(a, b). \quad (*)$$

With  $b = x_0$ , we then have  $|\varphi_a(x)| \leq d(a, x_0)$  for all  $x \in X$ . Therefore,  $\varphi_a$  is bounded and  $\varphi_a \in \mathcal{B}(X, \mathbb{R})$ .

## Theorem 43.7 (continued 1)

**Proof (continued).** Define  $\Phi : X \rightarrow \mathcal{B}(X, \mathbb{R})$  as  $\Phi(a) = \varphi_a$ . We now show that  $\Phi$  is an isometric embedding of  $(X, d)$  into the complete metric space  $(\mathcal{B}(X, \mathbb{R}), \rho)$ . Notice that since  $(\mathbb{R}, |\cdot|)$  is complete, then  $(\mathcal{B}(X, \mathbb{R}), \bar{\rho})$  is complete by Theorem 43.6. Since  $\bar{\rho}(f, g) = \min\{\rho(f, g), 1\}$  (see the Note before the statement of this theorem) then completeness of  $(\mathcal{B}(X, \mathbb{R}), \rho)$  is equivalent to the completeness of  $(\mathcal{B}(X, \mathbb{R}), \bar{\rho})$  (see the Note before the statement of Lemma 43.1 in these class notes, or see page 264). So  $(\mathcal{B}(X, \mathbb{R}), \rho)$  is in fact a complete metric space.

## Theorem 43.7 (continued 1)

**Proof (continued).** Define  $\Phi : X \rightarrow \mathcal{B}(X, \mathbb{R})$  as  $\Phi(a) = \varphi_a$ . We now show that  $\Phi$  is an isometric embedding of  $(X, d)$  into the complete metric space  $(\mathcal{B}(X, \mathbb{R}), \rho)$ . Notice that since  $(\mathbb{R}, |\cdot|)$  is complete, then  $(\mathcal{B}(X, \mathbb{R}), \bar{\rho})$  is complete by Theorem 43.6. Since  $\bar{\rho}(f, g) = \min\{\rho(f, g), 1\}$  (see the Note before the statement of this theorem) then completeness of  $(\mathcal{B}(X, \mathbb{R}), \rho)$  is equivalent to the completeness of  $(\mathcal{B}(X, \mathbb{R}), \bar{\rho})$  (see the Note before the statement of Lemma 43.1 in these class notes, or see page 264). So  $(\mathcal{B}(X, \mathbb{R}), \rho)$  is in fact a complete metric space. Next, for the isometry claim, let  $a, b \in X$ . Then

$$\begin{aligned} \rho(\varphi_a, \varphi_b) &= \sup\{|\varphi_a(x) - \varphi_b(x)| \mid x \in X\} \text{ since the metric on } \mathbb{R} \text{ is } |\cdot| \\ &= \sup\{|(d(x, a) - d(x, x_0)) - (d(x, b) - d(x, x_0))| \mid x \in X\} \\ &\quad \text{by the definition of } \varphi_a \text{ and } \varphi_b \\ &= \sup\{|d(x, a) - d(x, b)| \mid x \in X\} \\ &\leq d(a, b) \text{ by } (*). \end{aligned}$$

## Theorem 43.7 (continued 1)

**Proof (continued).** Define  $\Phi : X \rightarrow \mathcal{B}(X, \mathbb{R})$  as  $\Phi(a) = \varphi_a$ . We now show that  $\Phi$  is an isometric embedding of  $(X, d)$  into the complete metric space  $(\mathcal{B}(X, \mathbb{R}), \rho)$ . Notice that since  $(\mathbb{R}, |\cdot|)$  is complete, then  $(\mathcal{B}(X, \mathbb{R}), \bar{\rho})$  is complete by Theorem 43.6. Since  $\bar{\rho}(f, g) = \min\{\rho(f, g), 1\}$  (see the Note before the statement of this theorem) then completeness of  $(\mathcal{B}(X, \mathbb{R}), \rho)$  is equivalent to the completeness of  $(\mathcal{B}(X, \mathbb{R}), \bar{\rho})$  (see the Note before the statement of Lemma 43.1 in these class notes, or see page 264). So  $(\mathcal{B}(X, \mathbb{R}), \rho)$  is in fact a complete metric space. Next, for the isometry claim, let  $a, b \in X$ . Then

$$\begin{aligned} \rho(\varphi_a, \varphi_b) &= \sup\{|\varphi_a(x) - \varphi_b(x)| \mid x \in X\} \text{ since the metric on } \mathbb{R} \text{ is } |\cdot| \\ &= \sup\{|(d(x, a) - d(x, x_0)) - (d(x, b) - d(x, x_0))| \mid x \in X\} \\ &\quad \text{by the definition of } \varphi_a \text{ and } \varphi_b \\ &= \sup\{|d(x, a) - d(x, b)| \mid x \in X\} \\ &\leq d(a, b) \text{ by } (*). \end{aligned}$$

## Theorem 43.7 (continued 2)

**Theorem 43.7.** Let  $(X, d)$  be a metric space. There is an isometric embedding of  $X$  into a complete space.

**Proof (continued).** So  $\rho(\varphi_a, \varphi_b) \leq d(a, b)$ . But when  $x = a$   $|d(x, a) - d(x, b)| = d(a, b)$  and so  $\rho(\varphi_a, \varphi_b) = \sup\{|d(x, a) - d(x, b)| \mid x \in X\} \geq d(a, b)$ . Therefore,  $\rho(\varphi_a, \varphi_b) = d(a, b)$  and the mapping  $\Phi : X \rightarrow \mathcal{B}(X, \mathbb{R})$  is an isometry (that is,  $d(a, b) = \rho(\varphi_a, \varphi_b) = \rho(\Phi(a), \Phi(b))$ ). □