


# History of Geometry

## Chapter 2. The Elements of Euclid

### 2.1. Book I—Proofs of Theorems



# Table of contents

- 1 Euclid, Book I Proposition 1
- 2 Euclid, Book I Proposition 2
- 3 Euclid, Book I Proposition 5
- 4 Euclid, Book I Proposition 7
- 5 Euclid, Book I Proposition 8
- 6 Euclid, Book I Proposition 13
- 7 Euclid, Book I Proposition 14
- 8 Euclid, Book I Proposition 15
- 9 Euclid, Book I Proposition 16
- 10 Euclid, Book I Proposition 27
- 11 Euclid, Book I Proposition 29
- 12 Euclid, Book I Proposition 30
- 13 Euclid, Book I Proposition 31

# Euclid, Book I Proposition 1


**Euclid, Book I Proposition 1.** On a given finite straight line  $AB$  to construct an equilateral triangle.

**Proof.** By Postulate 3, we can construct a circle  $\Delta$  centered at point  $A$  and passing through point  $B$ . Also by Postulate 3, construct a circle  $E$  centered at point  $B$  and passing through point  $A$ . Let  $\Gamma$  be a point of intersection of circles  $\Delta$  and  $E$ .

# Euclid, Book I Proposition 1

**Euclid, Book I Proposition 1.** On a given finite straight line  $AB$  to construct an equilateral triangle.


**Proof.** By Postulate 3, we can construct a circle  $\Delta$  centered at point  $A$  and passing through point  $B$ . Also by Postulate 3, construct a circle  $E$  centered at point  $B$  and passing through point  $A$ . Let  $\Gamma$  be a point of intersection of circles  $\Delta$  and  $E$ . Construct a line segment joining points  $\Gamma$  and  $A$ , and construct a line segment joining  $\Gamma$  and  $B$  (using Postulate 1). Then the distance  $A\Gamma$  is equal to  $B\Gamma$  and equal to  $AB$  (by the definition of circle, Book I Definition 15).  $\square$



# Euclid, Book I Proposition 1

**Euclid, Book I Proposition 1.** On a given finite straight line  $AB$  to construct an equilateral triangle.

**Proof.** By Postulate 3, we can construct a circle  $\Delta$  centered at point  $A$  and passing through point  $B$ . Also by Postulate 3, construct a circle  $E$  centered at point  $B$  and passing through point  $A$ . Let  $\Gamma$  be a point of intersection of circles  $\Delta$  and  $E$ . Construct a line segment joining points  $\Gamma$  and  $A$ , and construct a line segment joining  $\Gamma$  and  $B$  (using Postulate 1). Then the distance  $A\Gamma$  is equal to  $B\Gamma$  and equal to  $AB$  (by the definition of circle, Book I Definition 15).  $\square$



## Euclid, Book I Proposition 2


**Euclid, Book I Proposition 2.** To place at a given point  $A$  a straight line  $AE$  equal to a given straight line  $B\Gamma$ .

**Proof.** Use Euclid I.1 to construct an equilateral triangle  $AB\Delta$  on segment  $AB$ . By Postulate 3, construct the circle with center  $B$  and passing through point  $\Gamma$ . Extend line segment  $\Delta B$  from point  $B$  until it intersects this circle at, say, point  $H$  (Postulate 2 allows this).

# Euclid, Book I Proposition 2

**Euclid, Book I Proposition 2.** To place at a given point  $A$  a straight line  $AE$  equal to a given straight line  $B\Gamma$ .


**Proof.** Use Euclid I.1 to construct an equilateral triangle  $AB\Delta$  on segment  $AB$ . By Postulate 3, construct the circle with center  $B$  and passing through point  $\Gamma$ . Extend line segment  $\Delta B$  from point  $B$  until it intersects this circle at, say, point  $H$  (Postulate 2 allows this). Also by Postulate 3, construct a second circle with center  $\Delta$  passing through  $H$ . Extend line segment  $\Delta A$  from point  $A$  until it intersects this second circle at, say, point  $E$  (Postulate 2). The distance  $B\Gamma$  equals the distance  $BH$  (because of properties of the first circle) and distance  $\Delta H$  equals distance  $\Delta E$  (because of properties of the second circle). So  $\Delta E = \Delta A + AE = \Delta H = \Delta B + BH$  or  $AE = BH$ , since  $\Delta B = \Delta A$ .  $\square$



# Euclid, Book I Proposition 2

**Euclid, Book I Proposition 2.** To place at a given point  $A$  a straight line  $AE$  equal to a given straight line  $B\Gamma$ .

**Proof.** Use Euclid I.1 to construct an equilateral triangle  $AB\Delta$  on segment  $AB$ . By Postulate 3, construct the circle with center  $B$  and passing through point  $\Gamma$ . Extend line segment  $\Delta B$  from point  $B$  until it intersects this circle at, say, point  $H$  (Postulate 2 allows this). Also by Postulate 3, construct a second circle with center  $\Delta$  passing through  $H$ . Extend line segment  $\Delta A$  from point  $A$  until it intersects this second circle at, say, point  $E$  (Postulate 2). The distance  $B\Gamma$  equals the distance  $BH$  (because of properties of the first circle) and distance  $\Delta H$  equals distance  $\Delta E$  (because of properties of the second circle). So  $\Delta E = \Delta A + AE = \Delta H = \Delta B + BH$  or  $AE = BH$ , since  $\Delta B = \Delta A$ .  $\square$





# Euclid, Book I Proposition 5

**Euclid, Book I Proposition 5.** If in a triangle,  $a = b$ , then  $\alpha = \beta$ .


**Proof.** Here, we present Euclid's proof and consider Figure 2.2(a). By Postulate 2 we can extend line segments  $CA$  and  $CB$  "continuously" to points  $F$  and  $G$  such that, by Euclid I.2,  $AF = BG$  (without loss of generality,  $CA < CB$  and we can extend  $CA$  and produce a segment  $CF$  longer than  $CB$  and then we can extend  $CB$  to produce a segment  $CG$  equal in length to  $CB$  by Euclid I.2; from this we have  $AF = BG$ ).

## Euclid, Book I Proposition 5

**Euclid, Book I Proposition 5.** If in a triangle,  $a = b$ , then  $\alpha = \beta$ .

**Proof.** Here, we present Euclid's proof and consider Figure 2.2(a). By Postulate 2 we can extend line segments  $CA$  and  $CB$  "continuously" to points  $F$  and  $G$  such that, by Euclid I.2,  $AF = BG$  (without loss of generality,  $CA < CB$  and we can extend  $CA$  and produce a segment  $CF$  longer than  $CB$  and then we can extend  $CB$  to produce a segment  $CG$  equal in length to  $CB$  by Euclid I.2; from this we have  $AF = BG$ ).

Next, by Postulate 2 we can introduce segments  $FB$  and  $AG$ . Now by Euclid I.4 ("side-angle-side"), triangles  $FCB$  and  $GCA$  are equal. So, corresponding angles and sides of the triangles are equal, so that  $\alpha + \delta = \beta + \varepsilon$ ,  $\eta = \zeta$ , and  $FB = GA$ . Again by Euclid I.4, triangle  $AFB$  and  $BGA$  are equal and so  $\delta = \varepsilon$ . Since  $\alpha + \delta = \beta + \varepsilon$  and  $\delta = \varepsilon$ , then  $\alpha = \beta$ , as claimed. □




## Euclid, Book I Proposition 5

**Euclid, Book I Proposition 5.** If in a triangle,  $a = b$ , then  $\alpha = \beta$ .

**Proof.** Here, we present Euclid's proof and consider Figure 2.2(a). By Postulate 2 we can extend line segments  $CA$  and  $CB$  "continuously" to points  $F$  and  $G$  such that, by Euclid I.2,  $AF = BG$  (without loss of generality,  $CA < CB$  and we can extend  $CA$  and produce a segment  $CF$  longer than  $CB$  and then we can extend  $CB$  to produce a segment  $CG$  equal in length to  $CB$  by Euclid I.2; from this we have  $AF = BG$ ).

Next, by Postulate 2 we can introduce segments  $FB$  and  $AG$ . Now by Euclid I.4 ("side-angle-side"), triangles  $FCB$  and  $GCA$  are equal. So, corresponding angles and sides of the triangles are equal, so that  $\alpha + \delta = \beta + \varepsilon$ ,  $\eta = \zeta$ , and  $FB = GA$ . Again by Euclid I.4, triangle  $AFB$  and  $BGA$  are equal and so  $\delta = \varepsilon$ . Since  $\alpha + \delta = \beta + \varepsilon$  and  $\delta = \varepsilon$ , then  $\alpha = \beta$ , as claimed. □



# Euclid, Book I Proposition 7

**Euclid, Book I Proposition 7.** Consider the two triangles of Figure 2.3(a), with the same base  $AB$  and with the third vertex on the same side of the base. If  $a = a'$  and  $b = b'$ , then points  $C$  and  $D$  are the same,  $C = D$ .

**Proof.** We present Euclid's proof. ASSUME  $C \neq D$ . Since triangle  $DAC$  is isosceles by hypothesis, then by Euclid I.5 we have  $\alpha + \beta = \gamma$ . Similarly, triangle  $DBC$  is isosceles and, again by Euclid I.5,  $\beta = \gamma + \delta$ . Since  $\alpha > 0$  then  $\gamma > \beta$ , and since  $\delta > 0$  then  $\gamma < \beta$ , a CONTRADICTION. So the assumption that  $C \neq D$  is false, and hence  $C = D$  as claimed.  $\square$

# Euclid, Book I Proposition 7

**Euclid, Book I Proposition 7.** Consider the two triangles of Figure 2.3(a), with the same base  $AB$  and with the third vertex on the same side of the base. If  $a = a'$  and  $b = b'$ , then points  $C$  and  $D$  are the same,  $C = D$ .

**Proof.** We present Euclid's proof. ASSUME  $C \neq D$ . Since triangle  $DAC$  is isosceles by hypothesis, then by Euclid I.5 we have  $\alpha + \beta = \gamma$ . Similarly, triangle  $DBC$  is isosceles and, again by Euclid I.5,  $\beta = \gamma + \delta$ . Since  $\alpha > 0$  then  $\gamma > \beta$ , and since  $\delta > 0$  then  $\gamma < \beta$ , a CONTRADICTION. So the assumption that  $C \neq D$  is false, and hence  $C = D$  as claimed.  $\square$




Figure 2.3(a)

## Euclid, Book I Proposition 7

**Euclid, Book I Proposition 7.** Consider the two triangles of Figure 2.3(a), with the same base  $AB$  and with the third vertex on the same side of the base. If  $a = a'$  and  $b = b'$ , then points  $C$  and  $D$  are the same,  $C = D$ .

**Proof.** We present Euclid's proof. ASSUME  $C \neq D$ . Since triangle  $DAC$  is isosceles by hypothesis, then by Euclid I.5 we have  $\alpha + \beta = \gamma$ . Similarly, triangle  $DBC$  is isosceles and, again by Euclid I.5,  $\beta = \gamma + \delta$ . Since  $\alpha > 0$  then  $\gamma > \beta$ , and since  $\delta > 0$  then  $\gamma < \beta$ , a CONTRADICTION. So the assumption that  $C \neq D$  is false, and hence  $C = D$  as claimed.  $\square$




Figure 2.3(a)

## Euclid, Book I Proposition 8

**Euclid, Book I Proposition 8.** If two triangles  $ABC$  and  $DEF$  have sides of equal lengths, then they also have equal angles.

**Proof.** Ostermann and Wanner credit the proof they give to Philo of Byzantium (circa 280 BCE–circa 220 BCE), declaring it “more elegant” than Euclid’s proof (though they do not give a specific reference). We move triangle  $ABC$  onto triangle  $DEF$  in such a way that the line segment  $BC$  is placed on segment  $EF$  (this is possible since these are the same length) and the point  $A$  is moved onto point  $G$  on the opposite side of  $EF$  to  $D$ ; see Figures 2.3(b) and (c).

# Euclid, Book I Proposition 8

**Euclid, Book I Proposition 8.** If two triangles  $ABC$  and  $DEF$  have sides of equal lengths, then they also have equal angles.

**Proof.** Ostermann and Wanner credit the proof they give to Philo of Byzantium (circa 280 BCE–circa 220 BCE), declaring it “more elegant” than Euclid’s proof (though they do not give a specific reference). We move triangle  $ABC$  onto triangle  $DEF$  in such a way that the line segment  $BC$  is placed on segment  $EF$  (this is possible since these are the same length) and the point  $A$  is moved onto point  $G$  on the opposite side of  $EF$  to  $D$ ; see Figures 2.3(b) and (c).




Figure 2.3(b) and (c)



# Euclid, Book I Proposition 8

**Euclid, Book I Proposition 8.** If two triangles  $ABC$  and  $DEF$  have sides of equal lengths, then they also have equal angles.

**Proof.** Ostermann and Wanner credit the proof they give to Philo of Byzantium (circa 280 BCE–circa 220 BCE), declaring it “more elegant” than Euclid’s proof (though they do not give a specific reference). We move triangle  $ABC$  onto triangle  $DEF$  in such a way that the line segment  $BC$  is placed on segment  $EF$  (this is possible since these are the same length) and the point  $A$  is moved onto point  $G$  on the opposite side of  $EF$  to  $D$ ; see Figures 2.3(b) and (c).




Figure 2.3(b) and (c)

# Euclid, Book I Proposition 8 (continued)

**Euclid, Book I Proposition 8.** If two triangles  $ABC$  and  $DEF$  have sides of equal lengths, then they also have equal angles.

**Proof (continued).** So  $DE = DG$  and triangle  $DEG$  is isosceles. Hence, by Euclid I.5,  $\alpha = \beta$ . Similarly, triangle  $DFG$  is also isosceles and  $\gamma = \delta$ . So  $\beta + \delta = \alpha + \gamma$ ; that is, the angle at  $A$  (equal to  $\beta + \delta$ ) equals the angle at  $D$  (equal to  $\alpha + \gamma$ ). Similar arguments (with different movements) show the other corresponding angles are also equal.  $\square$




Figure 2.3(b) and (c)

# Euclid, Book I Proposition 13

**Euclid I.13.** Let the line  $AB$  cut the line  $CD$ . With  $\alpha$  and  $\beta$  as the two resulting angles on the same side of line  $CD$ , we have  $\alpha + \beta = 2\perp$ .

**Proof.** Let  $B$  be the point of intersection of the lines. If the lines are perpendicular, then  $\alpha = \beta = \perp$  and we are done.

# Euclid, Book I Proposition 13

**Euclid I.13.** Let the line  $AB$  cut the line  $CD$ . With  $\alpha$  and  $\beta$  as the two resulting angles on the same side of line  $CD$ , we have  $\alpha + \beta = 2\text{ } \perp$ .

**Proof.** Let  $B$  be the point of intersection of the lines. If the lines are perpendicular, then  $\alpha = \beta = \text{ } \perp$  and we are done. So without loss of generality, we may assume that one of  $\alpha$  and  $\beta$  is greater than  $\text{ } \perp$ , say  $\beta > \text{ } \perp$ . Euclid I.11 gives the existence of a perpendicular to  $CD$  through point  $B$ . With  $\eta$  as the angle between this perpendicular and line  $AB$ , we have that  $\beta = \text{ } \perp + \eta$  and  $\alpha + \eta = \text{ } \perp$  (see Figure 2.5). Combining these two equations gives  $\alpha + \eta + \beta = 2\text{ } \perp + \eta$ , or  $\alpha + \beta = 2\text{ } \perp$ , as claimed.  $\square$

## Euclid, Book I Proposition 13

**Euclid I.13.** Let the line  $AB$  cut the line  $CD$ . With  $\alpha$  and  $\beta$  as the two resulting angles on the same side of line  $CD$ , we have  $\alpha + \beta = 2 \perp$ .

**Proof.** Let  $B$  be the point of intersection of the lines. If the lines are perpendicular, then  $\alpha = \beta = \perp$  and we are done. So without loss of generality, we may assume that one of  $\alpha$  and  $\beta$  is greater than  $\perp$ , say  $\beta > \perp$ . Euclid I.11 gives the existence of a perpendicular to  $CD$  through point  $B$ . With  $\eta$  as the angle between this perpendicular and line  $AB$ , we have that  $\beta = \perp + \eta$  and  $\alpha + \eta = \perp$  (see Figure 2.5). Combining these two equations gives  $\alpha + \eta + \beta = 2 \perp + \eta$ , or  $\alpha + \beta = 2 \perp$ , as claimed.  $\square$




Figure 2.5

## Euclid, Book I Proposition 13

**Euclid I.13.** Let the line  $AB$  cut the line  $CD$ . With  $\alpha$  and  $\beta$  as the two resulting angles on the same side of line  $CD$ , we have  $\alpha + \beta = 2 \perp$ .

**Proof.** Let  $B$  be the point of intersection of the lines. If the lines are perpendicular, then  $\alpha = \beta = \perp$  and we are done. So without loss of generality, we may assume that one of  $\alpha$  and  $\beta$  is greater than  $\perp$ , say  $\beta > \perp$ . Euclid I.11 gives the existence of a perpendicular to  $CD$  through point  $B$ . With  $\eta$  as the angle between this perpendicular and line  $AB$ , we have that  $\beta = \perp + \eta$  and  $\alpha + \eta = \perp$  (see Figure 2.5). Combining these two equations gives  $\alpha + \eta + \beta = 2 \perp + \eta$ , or  $\alpha + \beta = 2 \perp$ , as claimed.  $\square$



Figure 2.5

# Euclid, Book I Proposition 14

**Euclid, Book I Proposition 14.** Let line segment  $DB$  and line segment  $BA$  determine an angle  $\beta$ . If segment  $BC$  makes an angle  $\alpha$  with segment  $BA$  where point  $C$  is exterior to the first angle. If  $\alpha + \beta = 2\text{ } \perp$  then  $C$  lies on the line  $DB$ .

**Proof.** Let  $E$  line on the line  $DB$  and let  $\gamma$  be the angle between segments  $BA$  and  $BE$ . Then by Euclid I.13,  $\gamma + \beta = 2\text{ } \perp$ . By hypothesis,  $\alpha + \beta = 2\text{ } \perp$ , so  $\gamma + \beta = \alpha + \beta$  (by Postulate 4) and hence  $\gamma = \alpha$ . So points  $E$  and  $C$  lie on the same line (i.e., lie on the same side of the common angle). Since  $E$  lies on line  $DB$  then we have that  $C$  lies on line  $DB$ , as claimed.  $\square$

# Euclid, Book I Proposition 14

**Euclid, Book I Proposition 14.** Let line segment  $DB$  and line segment  $BA$  determine an angle  $\beta$ . If segment  $BC$  makes an angle  $\alpha$  with segment  $BA$  where point  $C$  is exterior to the first angle. If  $\alpha + \beta = 2\perp$  then  $C$  lies on the line  $DB$ .

**Proof.** Let  $E$  line on the line  $DB$  and let  $\gamma$  be the angle between segments  $BA$  and  $BE$ . Then by Euclid I.13,  $\gamma + \beta = 2\perp$ . By hypothesis,  $\alpha + \beta = 2\perp$ , so  $\gamma + \beta = \alpha + \beta$  (by Postulate 4) and hence  $\gamma = \alpha$ . So points  $E$  and  $C$  lie on the same line (i.e., lie on the same side of the common angle). Since  $E$  lies on line  $DB$  then we have that  $C$  lies on line  $DB$ , as claimed.  $\square$




Figure 2.6



# Euclid, Book I Proposition 14

**Euclid, Book I Proposition 14.** Let line segment  $DB$  and line segment  $BA$  determine an angle  $\beta$ . If segment  $BC$  makes an angle  $\alpha$  with segment  $BA$  where point  $C$  is exterior to the first angle. If  $\alpha + \beta = 2\perp$  then  $C$  lies on the line  $DB$ .

**Proof.** Let  $E$  line on the line  $DB$  and let  $\gamma$  be the angle between segments  $BA$  and  $BE$ . Then by Euclid I.13,  $\gamma + \beta = 2\perp$ . By hypothesis,  $\alpha + \beta = 2\perp$ , so  $\gamma + \beta = \alpha + \beta$  (by Postulate 4) and hence  $\gamma = \alpha$ . So points  $E$  and  $C$  lie on the same line (i.e., lie on the same side of the common angle). Since  $E$  lies on line  $DB$  then we have that  $C$  lies on line  $DB$ , as claimed.  $\square$




Figure 2.6

# Euclid, Book I Proposition 15

**Euclid, Book I Proposition 15.** If two lines cut one another, they make the opposite angles equal to one another.

**Proof.** Consider the opposite angles  $\alpha$  and  $\beta$ , and introduce angle  $\gamma$  as given in Figure 2.7. By Euclid I.13 we have  $\alpha + \gamma = 2 \text{ rt}$  and  $\gamma + \beta = 2 \text{ rt}$ . So  $\alpha + \gamma = \gamma + \beta$  (by Postulate 4). Therefore  $\alpha = \beta$ , as claimed.  $\square$

# Euclid, Book I Proposition 15

**Euclid, Book I Proposition 15.** If two lines cut one another, they make the opposite angles equal to one another.

**Proof.** Consider the opposite angles  $\alpha$  and  $\beta$ , and introduce angle  $\gamma$  as given in Figure 2.7. By Euclid I.13 we have  $\alpha + \gamma = 2 \perp$  and  $\gamma + \beta = 2 \perp$ . So  $\alpha + \gamma = \gamma + \beta$  (by Postulate 4). Therefore  $\alpha = \beta$ , as claimed.  $\square$




Figure 2.7

# Euclid, Book I Proposition 15

**Euclid, Book I Proposition 15.** If two lines cut one another, they make the opposite angles equal to one another.

**Proof.** Consider the opposite angles  $\alpha$  and  $\beta$ , and introduce angle  $\gamma$  as given in Figure 2.7. By Euclid I.13 we have  $\alpha + \gamma = 2 \perp$  and  $\gamma + \beta = 2 \perp$ . So  $\alpha + \gamma = \gamma + \beta$  (by Postulate 4). Therefore  $\alpha = \beta$ , as claimed.  $\square$




Figure 2.7

## Euclid, Book I Proposition 16

**Euclid, Book I Proposition 16.** If one side of a triangle is extended at  $C$ , the exterior angle is greater than both angles in the triangle opposite to  $C$ .

**Proof.** Let  $E$  be the midpoint of  $AC$ , which can be found by Euclid I.10. By Postulate 1, line segment  $BE$  exists. By Postulate 2, line segment  $BE$  can be extended to a point  $F$  such that  $BE = EF$ . The “grey” angles at point  $E$  of Figure 2.8 are opposite angles and so by Euclid I.15 are equal. So by Euclid I.4 (side-angle-side) the two grey triangles are equal.

# Euclid, Book I Proposition 16

**Euclid, Book I Proposition 16.** If one side of a triangle is extended at  $C$ , the exterior angle is greater than both angles in the triangle opposite to  $C$ .

**Proof.** Let  $E$  be the midpoint of  $AC$ , which can be found by Euclid I.10. By Postulate 1, line segment  $BE$  exists. By Postulate 2, line segment  $BE$  can be extended to a point  $F$  such that  $BE = EF$ . The “grey” angles at point  $E$  of Figure 2.8 are opposite angles and so by Euclid I.15 are equal. So by Euclid I.4 (side-angle-side) the two grey triangles are equal. Therefore the grey angle at point  $C$  is  $\alpha$  and is “obviously” smaller than  $\delta$  (we could use Common Notion 5 here); that is,  $\delta > \alpha$ , as claimed. Similarly, by bisecting side  $BC$  and introducing a triangle with a vertex at point  $C$ , we can use the other angle of size  $\delta$  to show  $\delta > \beta$ , as claimed.




Figure 2.8



# Euclid, Book I Proposition 16

**Euclid, Book I Proposition 16.** If one side of a triangle is extended at  $C$ , the exterior angle is greater than both angles in the triangle opposite to  $C$ .

**Proof.** Let  $E$  be the midpoint of  $AC$ , which can be found by Euclid I.10. By Postulate 1, line segment  $BE$  exists. By Postulate 2, line segment  $BE$  can be extended to a point  $F$  such that  $BE = EF$ . The “grey” angles at point  $E$  of Figure 2.8 are opposite angles and so by Euclid I.15 are equal. So by Euclid I.4 (side-angle-side) the two grey triangles are equal. Therefore the grey angle at point  $C$  is  $\alpha$  and is “obviously” smaller than  $\delta$  (we could use Common Notion 5 here); that is,  $\delta > \alpha$ , as claimed. Similarly, by bisecting side  $BC$  and introducing a triangle with a vertex at point  $C$ , we can use the other angle of size  $\delta$  to show  $\delta > \beta$ , as claimed.




Figure 2.8



# Euclid, Book I Proposition 27

**Euclid, Book I Proposition 27.** If some line cuts two line  $a$  and  $b$  such that alternate interior angles  $\alpha$  and  $\beta$  are equal, then lines  $a$  and  $b$  are parallel, denoted  $a \parallel b$ .

**Proof.** Let points  $E$  and  $F$  be the points of intersection of the cutting line with lines  $a$  and  $b$  (see Figure 2.10). ASSUME that lines  $a$  and  $b$  are not parallel; then they meet at some point  $G$ . Without loss of generality, suppose point  $G$  is on side of the cutting line in which angle  $\beta$  lies, as in Figure 2.10. Then  $EGF$  is a triangle with  $\alpha$  as an exterior angle. So by Euclid I.16,  $\alpha > \beta$ . But this CONTRADICTS the assumption that  $\alpha = \beta$ . So the assumption that lines  $a$  and  $b$  are not parallel is false, and hence lines  $a$  and  $b$  are parallel, as claimed.  $\square$



# Euclid, Book I Proposition 27

**Euclid, Book I Proposition 27.** If some line cuts two line  $a$  and  $b$  such that alternate interior angles  $\alpha$  and  $\beta$  are equal, then lines  $a$  and  $b$  are parallel, denoted  $a \parallel b$ .

**Proof.** Let points  $E$  and  $F$  be the points of intersection of the cutting line with lines  $a$  and  $b$  (see Figure 2.10). ASSUME that lines  $a$  and  $b$  are not parallel; then they meet at some point  $G$ . Without loss of generality, suppose point  $G$  is on side of the cutting line in which angle  $\beta$  lies, as in Figure 2.10. Then  $EGF$  is a triangle with  $\alpha$  as an exterior angle. So by Euclid I.16,  $\alpha > \beta$ . But this CONTRADICTS the assumption that  $\alpha = \beta$ . So the assumption that lines  $a$  and  $b$  are not parallel is false, and hence lines  $a$  and  $b$  are parallel, as claimed.  $\square$




Figure 2.10

## Euclid, Book I Proposition 27

**Euclid, Book I Proposition 27.** If some line cuts two line  $a$  and  $b$  such that alternate interior angles  $\alpha$  and  $\beta$  are equal, then lines  $a$  and  $b$  are parallel, denoted  $a \parallel b$ .

**Proof.** Let points  $E$  and  $F$  be the points of intersection of the cutting line with lines  $a$  and  $b$  (see Figure 2.10). ASSUME that lines  $a$  and  $b$  are not parallel; then they meet at some point  $G$ . Without loss of generality, suppose point  $G$  is on side of the cutting line in which angle  $\beta$  lies, as in Figure 2.10. Then  $EGF$  is a triangle with  $\alpha$  as an exterior angle. So by Euclid I.16,  $\alpha > \beta$ . But this CONTRADICTS the assumption that  $\alpha = \beta$ . So the assumption that lines  $a$  and  $b$  are not parallel is false, and hence lines  $a$  and  $b$  are parallel, as claimed.  $\square$




Figure 2.10

# Euclid, Book I Proposition 29

**Euclid, Book I Proposition 29.** Parallel lines cut by some line, have alternate interior angles are equal.

**Proof.** Let  $a$  and  $b$  be parallel lines and let  $\alpha$  and  $\beta$  be alternate interior angles that result from the cutting line (see Figure 2.11). ASSUME  $\alpha > \beta$ . Introduce angle  $\gamma$  as in Figure 2.11. By Euclid I.13 we have  $\alpha + \gamma = 2 \text{ } \perp$ , and hence  $\beta + \gamma < 2 \text{ } \perp$ . By the Parallel Postulate (Postulate 5), lines  $a$  and  $b$  meet (and do so on the side of the cutting line containing angles  $\beta$  and  $\gamma$ ), CONTRADICTING the hypothesis that  $a \parallel b$ . So the assumption that  $\alpha > \beta$  is false and hence we have  $\alpha \leq \beta$ . Similarly, if we assume  $\alpha < \beta$  we can show that  $\alpha \geq \beta$ . Hence  $\alpha = \beta$ , as claimed.  $\square$

# Euclid, Book I Proposition 29

**Euclid, Book I Proposition 29.** Parallel lines cut by some line, have alternate interior angles are equal.

**Proof.** Let  $a$  and  $b$  be parallel lines and let  $\alpha$  and  $\beta$  be alternate interior angles that result from the cutting line (see Figure 2.11). ASSUME  $\alpha > \beta$ . Introduce angle  $\gamma$  as in Figure 2.11. By Euclid I.13 we have  $\alpha + \gamma = 2\text{ } \perp$ , and hence  $\beta + \gamma < 2\text{ } \perp$ . By the Parallel Postulate (Postulate 5), lines  $a$  and  $b$  meet (and do so on the side of the cutting line containing angles  $\beta$  and  $\gamma$ ), CONTRADICTING the hypothesis that  $a \parallel b$ . So the assumption that  $\alpha > \beta$  is false and hence we have  $\alpha \leq \beta$ . Similarly, if we assume  $\alpha < \beta$  we can show that  $\alpha \geq \beta$ . Hence  $\alpha = \beta$ , as claimed.  $\square$




Figure 2.11

# Euclid, Book I Proposition 29

**Euclid, Book I Proposition 29.** Parallel lines cut by some line, have alternate interior angles are equal.

**Proof.** Let  $a$  and  $b$  be parallel lines and let  $\alpha$  and  $\beta$  be alternate interior angles that result from the cutting line (see Figure 2.11). ASSUME  $\alpha > \beta$ . Introduce angle  $\gamma$  as in Figure 2.11. By Euclid I.13 we have  $\alpha + \gamma = 2\text{ } \perp$ , and hence  $\beta + \gamma < 2\text{ } \perp$ . By the Parallel Postulate (Postulate 5), lines  $a$  and  $b$  meet (and do so on the side of the cutting line containing angles  $\beta$  and  $\gamma$ ), CONTRADICTING the hypothesis that  $a \parallel b$ . So the assumption that  $\alpha > \beta$  is false and hence we have  $\alpha \leq \beta$ . Similarly, if we assume  $\alpha < \beta$  we can show that  $\alpha \geq \beta$ . Hence  $\alpha = \beta$ , as claimed.  $\square$




Figure 2.11

# Euclid, Book I Proposition 30

**Euclid, Book I Proposition 30.** For any three (distinct) lines  $a, b, c$ , if  $a \parallel b$  and  $b \parallel a$  then  $a \parallel c$ .

**Proof.** Let the angles between the parallel lines  $a, b, c$  and the cutting line be  $\alpha, \beta, \gamma$ , respectively (see Figure 2.12). If we introduce the angles opposite  $\alpha, \beta, \gamma$  as  $\alpha', \beta', \gamma'$  (not pictured in Figure 2.12) then by Euclid I.15 we have  $\alpha = \alpha'$ ,  $\beta = \beta'$ , and  $\gamma = \gamma'$ . By Euclid I.29,  $\alpha = \beta'$  since  $a \parallel b$ , and  $\beta = \gamma'$  since  $b \parallel c$ . Hence,  $\alpha = \beta' = \beta = \gamma'$ . Finally, by Euclid I.27, we have  $a \parallel c$ , as claimed.  $\square$

# Euclid, Book I Proposition 30

**Euclid, Book I Proposition 30.** For any three (distinct) lines  $a, b, c$ , if  $a \parallel b$  and  $b \parallel c$  then  $a \parallel c$ .

**Proof.** Let the angles between the parallel lines  $a, b, c$  and the cutting line be  $\alpha, \beta, \gamma$ , respectively (see Figure 2.12). If we introduce the angles opposite  $\alpha, \beta, \gamma$  as  $\alpha', \beta', \gamma'$  (not pictured in Figure 2.12) then by Euclid I.15 we have  $\alpha = \alpha'$ ,  $\beta = \beta'$ , and  $\gamma = \gamma'$ . By Euclid I.29,  $\alpha = \beta'$  since  $a \parallel b$ , and  $\beta = \gamma'$  since  $b \parallel c$ . Hence,  $\alpha = \beta' = \beta = \gamma'$ . Finally, by Euclid I.27, we have  $a \parallel c$ , as claimed.  $\square$




Figure 2.12

# Euclid, Book I Proposition 30

**Euclid, Book I Proposition 30.** For any three (distinct) lines  $a, b, c$ , if  $a \parallel b$  and  $b \parallel c$  then  $a \parallel c$ .

**Proof.** Let the angles between the parallel lines  $a, b, c$  and the cutting line be  $\alpha, \beta, \gamma$ , respectively (see Figure 2.12). If we introduce the angles opposite  $\alpha, \beta, \gamma$  as  $\alpha', \beta', \gamma'$  (not pictured in Figure 2.12) then by Euclid I.15 we have  $\alpha = \alpha'$ ,  $\beta = \beta'$ , and  $\gamma = \gamma'$ . By Euclid I.29,  $\alpha = \beta'$  since  $a \parallel b$ , and  $\beta = \gamma'$  since  $b \parallel c$ . Hence,  $\alpha = \beta' = \beta = \gamma'$ . Finally, by Euclid I.27, we have  $a \parallel c$ , as claimed.  $\square$




Figure 2.12



# Euclid, Book I Proposition 31

**Euclid, Book I Proposition 31.** To draw a parallel to a given line  $a$  through a given point  $A$  not on the  $a$ .

**Proof.** By Euclid I.12, we can construct a perpendicular to line  $a$  through point  $A$ . By Euclid I.11, we can construct a perpendicular  $b$  to the perpendicular through point  $A$ . We then have that the alternate interior angles for lines  $a$  and  $b$  are both right angles and hence by Euclid I.27 line  $b$  is parallel to line  $a$  and passes through point  $A$ , as needed.  $\square$

# Euclid, Book I Proposition 31

**Euclid, Book I Proposition 31.** To draw a parallel to a given line  $a$  through a given point  $A$  not on the  $a$ .

**Proof.** By Euclid I.12, we can construct a perpendicular to line  $a$  through point  $A$ . By Euclid I.11, we can construct a perpendicular  $b$  to the perpendicular through point  $A$ . We then have that the alternate interior angles for lines  $a$  and  $b$  are both right angles and hence by Euclid I.27 line  $b$  is parallel to line  $a$  and passes through point  $A$ , as needed.  $\square$




Figure 2.13

# Euclid, Book I Proposition 31

**Euclid, Book I Proposition 31.** To draw a parallel to a given line  $a$  through a given point  $A$  not on the  $a$ .

**Proof.** By Euclid I.12, we can construct a perpendicular to line  $a$  through point  $A$ . By Euclid I.11, we can construct a perpendicular  $b$  to the perpendicular through point  $A$ . We then have that the alternate interior angles for lines  $a$  and  $b$  are both right angles and hence by Euclid I.27 line  $b$  is parallel to line  $a$  and passes through point  $A$ , as needed.  $\square$




Figure 2.13