


## Real Analysis

### Chapter 4. Basic Concepts of Probability

#### 4.8. Independent Random Variables—Proofs of Theorems



()

Real Analysis

February 8, 2019

1 / 13

Theorem 1.3.9. The Monotone Class Theorem

### Theorem 1.3.9. The Monotone Class Theorem

#### Theorem 1.3.9. The Monotone Class Theorem.

Let  $\mathcal{F}_0$  be a field of subsets of  $\Omega$  and  $\mathcal{C}$  a class of subsets of  $\Omega$  that is monotone. If  $\mathcal{C} \supset \mathcal{F}_0$ , then  $\mathcal{C} \supset \sigma(\mathcal{F}_0)$ , the minimal  $\sigma$ -field over  $\mathcal{F}_0$ .

**Proof.** Let  $\mathcal{F} = \sigma(\mathcal{F}_0)$  and let  $\mathcal{M}$  be the smallest monotone class containing all sets of  $\mathcal{F}_0$ . We will show that  $\mathcal{M} = \mathcal{F}$ ; that is, the smallest monotone class containing field  $\mathcal{F}_0$  and the smallest  $\sigma$ -field over field  $\mathcal{F}_0$  are the same.

Let  $A \in \mathcal{M}$  be fixed. Define

$$\mathcal{M}_A = \{B \in \mathcal{M} \mid A \cap B, A \cap B^c, A^c \cap B \in \mathcal{M}\} \subset \mathcal{M}.$$

If  $\{B_n\}_{n=1}^\infty \subset \mathcal{M}_A$  with  $B_1 \subset B_2 \subset \dots$  then  $A \cap B_n, A \cap B_n^c, A^c \cap B_n \in \mathcal{M}$  for all  $n \in \mathbb{N}$ . Then  $\bigcup_{n=1}^\infty B_n \in \mathcal{M}$  since  $\mathcal{M}$  is monotone, and  $A \cap (\bigcup_{n=1}^\infty B_n) = \bigcup_{n=1}^\infty (A \cap B_n) \in \mathcal{M}$  since  $(A \cap B_1) \subset (A \cap B_2) \subset \dots$  and  $\mathcal{M}$  is monotone. Similarly,  $A^c \cap (\bigcup_{n=1}^\infty B_n) = \bigcup_{n=1}^\infty (A^c \cap B_n) \in \mathcal{M}$ .

()

Real Analysis

February 8, 2019

3 / 13

Theorem 1.3.9. The Monotone Class Theorem

### Theorem 1.3.9 (continued 1)

#### Theorem 1.3.9. The Monotone Class Theorem.

Let  $\mathcal{F}_0$  be a field of subsets of  $\Omega$  and  $\mathcal{C}$  a class of subsets of  $\Omega$  that is monotone. If  $\mathcal{C} \supset \mathcal{F}_0$ , then  $\mathcal{C} \supset \sigma(\mathcal{F}_0)$ , the minimal  $\sigma$ -field over  $\mathcal{F}_0$ .

**Proof (continued).** Also,  $(\bigcup_{n=1}^\infty B_n)^c = \bigcap_{n=1}^\infty B_n^c \in \mathcal{M}$  since  $B_1^c \supset B_2^c \supset \dots$  and  $\mathcal{M}$  is monotone. Next,  $A \cap (\bigcup_{n=1}^\infty B_n)^c = A \cap (\bigcap_{n=1}^\infty B_n^c) = \bigcap_{n=1}^\infty (A \cap B_n^c) \in \mathcal{M}$  since  $(A \cap B_1^c) \supset (A \cap B_2^c) \supset \dots$  and  $\mathcal{M}$  is monotone.

If  $\{B_n\}_{n=1}^\infty \subset \mathcal{M}_A$  with  $B_1 \supset B_2 \supset \dots$  then  $A \cap B_n, A \cap B_n^c, A^c \cap B_n \in \mathcal{M}$  for all  $n \in \mathbb{N}$ . Then  $\bigcap_{n=1}^\infty B_n \in \mathcal{M}$  since  $\mathcal{M}$  is monotone, and  $A \cap (\bigcap_{n=1}^\infty B_n) = \bigcap_{n=1}^\infty (A \cap B_n) \in \mathcal{M}$  since  $(A \cap B_1) \supset (A \cap B_2) \supset \dots$  and  $\mathcal{M}$  is monotone. Similarly,  $A^c \cap (\bigcap_{n=1}^\infty B_n) = \bigcap_{n=1}^\infty (A^c \cap B_n) \in \mathcal{M}$ . Also,  $(\bigcap_{n=1}^\infty B_n)^c = \bigcup_{n=1}^\infty B_n^c \in \mathcal{M}$  since  $B_1^c \supset B_2^c \supset \dots$  and  $\mathcal{M}$  is monotone. Next,  $A \cap (\bigcap_{n=1}^\infty B_n)^c = A \cap (\bigcup_{n=1}^\infty B_n^c) = \bigcup_{n=1}^\infty (A \cap B_n^c) \in \mathcal{M}$  since  $(A \cap B_1^c) \supset (A \cap B_2^c) \supset \dots$  and  $\mathcal{M}$  is monotone. Therefore,  $\mathcal{M}_A$  is a monotone class.

()

Real Analysis

February 8, 2019

4 / 13

Theorem 1.3.9. The Monotone Class Theorem

### Theorem 1.3.9 (continued 2)

#### Theorem 1.3.9. The Monotone Class Theorem.

Let  $\mathcal{F}_0$  be a field of subsets of  $\Omega$  and  $\mathcal{C}$  a class of subsets of  $\Omega$  that is monotone. If  $\mathcal{C} \supset \mathcal{F}_0$ , then  $\mathcal{C} \supset \sigma(\mathcal{F}_0)$ , the minimal  $\sigma$ -field over  $\mathcal{F}_0$ .

**Proof (continued).** If  $A \in \mathcal{F}_0$  then  $A^c \in \mathcal{F}_0$  since  $\mathcal{F}_0$  is a field, and for any  $B \in \mathcal{F}_0$  we must also have  $A \cap B, B^c, A \cap B^c, A^c \cap B \in \mathcal{F}_0$  since  $\mathcal{F}_0$  is a field. Since  $B$  satisfies these conditions, then  $B \in \mathcal{M}_A$ ; that is, if  $A \in \mathcal{F}_0$  then  $\mathcal{F}_0 \subset \mathcal{M}_A$ . Since  $\mathcal{M}$  is the smallest monotone monotone class containing all sets in  $\mathcal{F}_0$  and  $\mathcal{M}_A$  is a monotone class containing  $\mathcal{F}_0$  (if  $A \in \mathcal{F}_0$ ) then we must have  $\mathcal{M} \subset \mathcal{M}_A$ . Since  $\mathcal{M}_A \subset \mathcal{M}$  by the definition of  $\mathcal{M}_A$ , then we have  $\mathcal{M}_A = \mathcal{M}$  for any  $A \in \mathcal{F}_0$ . So for any  $B \in \mathcal{M} = \mathcal{M}_A$  we have  $A \cap B, A \cap B^c, A^c \cap B \in \mathcal{M} = \mathcal{M}_A$ , provided  $A \in \mathcal{F}_0$ . That is,  $A \in \mathcal{F}_0$  implies  $A \in \mathcal{M}_B$ , so that  $\mathcal{F}_0 \subset \mathcal{M}_B$ . But as just shown for  $\mathcal{M}_A$ ,  $\mathcal{M}_B$  is also a monotone class (regardless of whether  $B \in \mathcal{F}_0$  or not), and so  $\mathcal{M}_B$  is a monotone class containing  $\mathcal{F}_0$  so that  $\mathcal{M}_B \supset \mathcal{M}$  by the minimality definition of  $\mathcal{M}$ . But  $\mathcal{M}_B \subset \mathcal{M}$  by the definition of  $\mathcal{M}_B$ , so that  $\mathcal{M}_B = \mathcal{M}$  for any  $B \in \mathcal{M}$ .

()

Real Analysis

February 8, 2019

5 / 13

## Theorem 1.3.9 (continued 3)

**Proof (continued).** Next, for any  $A, B \in \mathcal{M} = \mathcal{M}_A$ , we have (by the definition of  $\mathcal{M}_A$ )  $A \cap B, A \cap B^c, A^c \cap B \in \mathcal{M} = \mathcal{M}_A$  so that  $\mathcal{M}$  is closed under complements (since by definition  $\Omega \in \mathcal{F}_0$  so for  $A \in \mathcal{M}$ ,  $A^c \cap \Omega = A^c \in \mathcal{M}$ ) and finite intersections. That is,  $\mathcal{M}$  is a field. We also claim that  $\mathcal{M}$  is a  $\sigma$ -field. Let  $A_1, A_2, \dots \in \mathcal{M}$ . Then  $A_1 \subset (A_1 \cup A_2) \subset (A_1 \cup A_2 \cup A_3) \subset \dots$  and since  $\mathcal{M}$  is a monotone class,  $\bigcup_{n=1}^{\infty} A_n = \bigcup_{k=1}^{\infty} (A_1 \cup A_2 \cup \dots \cup A_k) \in \mathcal{M}$ . Also,  $A_1 \supset (A_1 \cap A_2) \supset (A_1 \cap A_2 \cap A_3) \supset \dots$  and since  $\mathcal{M}$  is a monotone class  $\bigcap_{n=1}^{\infty} A_n = \bigcap_{k=1}^{\infty} (A_1 \cap A_2 \cap \dots \cap A_k) \in \mathcal{M}$ . That is,  $\mathcal{M}$  is a field closed under countable unions (and countable intersections). So  $\mathcal{M}$  is a  $\sigma$ -field which contains  $\mathcal{F}_0$ . Since  $\mathcal{F} = \sigma(\mathcal{F}_0)$  is the minimal  $\sigma$ -field containing  $\mathcal{F}_0$ , then  $\mathcal{F} \subset \mathcal{M}$ . Since  $\mathcal{F}$  is a  $\sigma$ -field, then it is also a monotone class (containing  $\mathcal{F}_0$ ), and by the minimality of monotone class  $\mathcal{M}$ , we have  $\mathcal{M} \subset \mathcal{F}$ . Therefore  $\mathcal{M} = \mathcal{F}$ .

Since  $\mathcal{C}$  is a monotone class of subsets of  $\Omega$  containing  $\mathcal{F}_0$ , then  $\mathcal{C} \supset \mathcal{M} \supset \mathcal{F}_0$ , and so we have  $\mathcal{C} \supset \mathcal{M} = \mathcal{F} = \sigma(\mathcal{F}_0)$ , as claimed.  $\square$

( )

Real Analysis

February 8, 2019

6 / 13

## Theorem 4.8.3

**Theorem 4.8.3.** Let  $X_1, X_2, \dots, X_n$  be random variables on  $(\Omega, \mathcal{F}, P)$ . Let  $F_i$  be the distribution function of  $X_i$ ,  $i = 1, 2, \dots, n$  (so  $F_i(x_i) = P(\{X_i \leq x_i\})$ ) and  $F$  the distribution function of  $X = (X_1, X_2, \dots, X_n)$  (that is,  $F(x_1, x_2, \dots, x_n) = P(\{X_1 \leq x_1, X_2 \leq x_2, \dots, X_n \leq x_n\})$ ). Then  $X_1, X_2, \dots, X_n$  are independent if and only if  $F(x_1, x_2, \dots, x_n) = F_1(x_1)F_2(x_2) \cdots F_n(x_n)$  for all  $x_1, x_2, \dots, x_n \in \mathbb{R}$ .

**Proof.** If  $X_1, X_2, \dots, X_n$  are independent then

$$\begin{aligned} F(x_1, x_2, \dots, x_n) &= P(\{X_1 \leq x_1, X_2 \leq x_2, \dots, X_n \leq x_n\}) \\ &= P(\{X_1 \leq x_1\})P(\{X_2 \leq x_2\}) \cdots P(\{X_n \leq x_n\}) = F_1(x_1)F_2(x_2) \cdots F_n(x_n), \end{aligned}$$

as claimed.

( )

Real Analysis

February 8, 2019

7 / 13

## Theorem 4.8.3 (continued 1)

**Proof (continued).** Now suppose

$F(x_1, x_2, \dots, x_n) = F_1(x_1)F_2(x_2) \cdots F_n(x_n)$  for all  $x_1, x_2, \dots, x_n \in \mathbb{R}$ . Let  $a = (a_1, a_2, \dots, a_n), b = (b_1, b_2, \dots, b_n) \in \mathbb{R}^n$  with  $a \leq b$  (that is,  $a_i \leq b_i$  for  $i = 1, 2, \dots, n$ ). Then

$$(a, b] = \{(x_1, x_2, \dots, x_n) \in \mathbb{R}^n \mid a_i < x_i \leq b_i \text{ for } i = 1, 2, \dots, n\}$$

(see page 26 of the text). So

$$\begin{aligned} P_X((a, b]) &= P(\{X \in (a, b]\}) \\ &= P(\{a_1 < X_1 \leq b_1, a_2 < X_2 \leq b_2, \dots, a_n < X_n \leq b_n\}) \\ &= F((a, b]) = (F_1(b_1) - F_1(a_1))(F_2(b_2) - F_2(a_2)) \cdots \\ &\quad (F_n(b_n) - F_n(a_n)) \text{ by Example 1.4.10(a) of the text} \\ &\quad \text{and the hypothesis on } F \\ &= P_{X_1}(a_1, b_1] P_{X_2}(a_2, b_2] \cdots P_{X_n}(a_n, b_n]. \end{aligned}$$

( )

Real Analysis

February 8, 2019

8 / 13

## Theorem 4.8.3 (continued 2)

**Proof (continued).** So

$$\begin{aligned} &P(\{X_1 \in B_1, X_2 \in B_2, \dots, X_n \in B_n\}) \\ &= P(\{X_1 \in B_1\})P(\{X_2 \in B_2\}) \cdots P(\{X_n \in B_n\}) \quad (1) \end{aligned}$$

holds when each  $B_i$  is of the form  $B_i = (a_i, b_i] \subset \mathbb{R}$ .

Now fix the intervals  $B_2, B_3, \dots, B_n$  and let  $\mathcal{C}$  be the collection of sets  $B_1 \in \mathcal{B}(\mathbb{R})$  for which equation (1) holds (so  $B_1$  need not be of the form  $(a_1, b_1]$  here, but  $\mathcal{C}$  does include all intervals of this form). Suppose  $\{A_n\}_{n=1}^{\infty} \subset \mathcal{C}$  and  $A_1 \subset A_2 \subset \dots$ . Then

$$\{X_1 \in \bigcup_{m=1}^k A_m, X_2 \in B_2, \dots, X_n \in B_n\} = \{X_1 \in A_k, X_2 \in B_2, \dots, X_n \in B_n\}$$

and since  $A_k \in \mathcal{C}$  then

$$\begin{aligned} &P\left(\left\{X_1 \in \bigcup_{m=1}^k A_m, X_2 \in B_2, \dots, X_n \in B_n\right\}\right) \\ &= P(\{X_1 \in A_k, X_2 \in B_2, \dots, X_n \in B_n\}) \end{aligned}$$

( )

Real Analysis

February 8, 2019

9 / 13



## Theorem 4.8.3 (continued 3)

**Proof (continued).** ...

$$= P(\{X_1 \in A_k\})P(\{X_2 \in B_2\}) \cdots P(\{X_n \in B_n\}) \\ = P\left(\left\{X_1 \in \bigcup_{m=1}^k A_m\right\}\right) P(\{X_2 \in B_2\}) \cdots P(\{X_n \in B_n\}).$$

Now

$$\left\{X_1 \in \bigcup_{m=1}^k A_m, X_2 \in B_2, \dots, X_n \in B_n\right\} \\ \subset \left\{X_1 \in \bigcup_{m=1}^{k+1} A_m, X_2 \in B_2, \dots, X_n \in B_n\right\}$$

so this sequence of sets of events indexed by  $k$  is ascending so that

$$\lim_{k \rightarrow \infty} \left\{X_1 \in \bigcup_{m=1}^k A_m, X_2 \in B_2, \dots, X_n \in B_n\right\} \\ = \left\{X_1 \in \bigcup_{m=1}^\infty A_m, X_2 \in B_2, \dots, X_n \in B_n\right\}$$

and since probability measure  $P$  is continuous (Proposition 17.2(i) of Royden and Fitzpatrick)...

## Theorem 4.8.3 (continued 4)

**Proof (continued).** ...

$$P(\{X_1 \in \bigcup_{m=1}^\infty A_m, X_2 \in B_2, \dots, X_n \in B_n\}) \\ = P\left(\lim_{k \rightarrow \infty} \left\{X_1 \in \bigcup_{m=1}^k A_m, X_2 \in B_2, \dots, X_n \in B_n\right\}\right) \\ = P\left(\lim_{k \rightarrow \infty} \{X_1 \in A_k, X_2 \in B_2, \dots, X_n \in B_n\}\right) \\ = \lim_{k \rightarrow \infty} P(\{X_1 \in A_k, X_2 \in B_2, \dots, X_n \in B_n\}) \\ = \lim_{k \rightarrow \infty} P(\{X_1 \in A_k\})P(\{X_2 \in B_2\}) \cdots P(\{X_n \in B_n\}) \text{ since } A_k \in \mathcal{C} \\ = P\left(\lim_{k \rightarrow \infty} \{X_1 \in A_k\}\right) P(\{X_2 \in B_2\}) \cdots P(\{X_n \in B_n\}) \\ = P(\{X_1 \in \bigcup_{m=1}^\infty A_m\}) P(\{X_2 \in B_2\}) \cdots P(\{X_n \in B_n\})$$

and so  $\bigcup_{m=1}^\infty A_m \in \mathcal{C}$ .

## Theorem 4.8.3 (continued 5)

**Proof (continued).** Similarly, if  $\{A_m\}_{m=1}^\infty \subset \mathcal{C}$  with  $A_1 \supset A_2 \supset \dots$ , we have by the continuity of probability measure  $P$  (Proposition 17.2(ii) of Royden and Fitzpatrick) that  $\bigcap_{m=1}^\infty A_m \in \mathcal{C}$ . So collection  $\mathcal{C}$  for which (1) holds is a monotone class. Now if  $B_1$  is a finite union of disjoint "right-semiclosed intervals,"  $B_1 = (a_1^1, b_1^1] \cup (a_1^2, b_1^2] \cup \dots \cup (a_1^\ell, b_1^\ell]$  then

$$\{X_1 \in B_1, X_2 \in B_2, \dots, X_n \in B_n\} = \{X_1 \in (a_1^1, b_1^1], X_2 \in B_2, \dots, X_n \in B_n\} \\ \cup \{X_1 \in (a_1^2, b_1^2], X_2 \in B_2, \dots, X_n \in B_n\} \cup \dots \\ \cup \{X_1 \in (a_1^\ell, b_1^\ell], X_2 \in B_2, \dots, X_n \in B_n\},$$

so by finite additivity of probability measure  $P$ ,

$$P(\{X_1 \in B_1, X_2 \in B_2, \dots, X_n \in B_n\}) \\ = P(\{X_1 \in (a_1^1, b_1^1], X_2 \in B_2, \dots, X_n \in B_n\}) \\ + P(\{X_1 \in (a_1^2, b_1^2], X_2 \in B_2, \dots, X_n \in B_n\}) + \dots \\ + P(\{X_1 \in (a_1^\ell, b_1^\ell], X_2 \in B_2, \dots, X_n \in B_n\})$$

## Theorem 4.8.3 (continued 6)

**Proof (continued).** ...

$$= (P(\{X_1 \in (a_1^1, b_1^1]\}) + P(\{X_1 \in (a_1^2, b_1^2]\}) + \dots \\ + P(\{X_1 \in (a_1^\ell, b_1^\ell]\})) P(\{X_2 \in B_2\}) \cdots P(\{X_n \in B_n\}) \\ = P(\{X_1 \in B_1\})P(\{X_2 \in B_2\}) \cdots P(\{X_n \in B_n\}),$$

so  $\mathcal{C}$  includes the field of all finite unions of right-semiclosed intervals. So, by the Monotone Class Theorem,  $\mathcal{C}$  includes the  $\sigma$ -field generated by the field of all finite unions of right-semiclosed intervals (which, by the Royden and Fitzpatrick Exercise 1.36 is the Borel sets in  $\mathbb{R}$ ). So (1) holds if  $B_1$  is any set. We can now inductively show that (1) holds for any  $B_2, B_3, \dots, B_n$  Borel sets. (As the text says, "Explicitly, we prove by induction that if  $B_1, \dots, B_i$  are arbitrary Borel sets and  $B_{i+1}, \dots, B_n$  are right-semiclosed intervals, then (1) holds for  $B_1, \dots, B_n$ ." )  $\square$