

# Cyclic and Rotational Decompositions of $K_n$ into Stars

Robert B. Gardner

*Institute of Mathematical and Physical Sciences*

*East Tennessee State University*

*Johnson City, Tennessee 37614 - 0296*

**Abstract.** We give necessary and sufficient conditions for the existence of a decomposition of the complete graph into stars which admits either a cyclic or a rotational automorphism.

## 1 Introduction

We denote the complete graph on  $n$  vertices by  $K_n$  and the star with  $m$  edges by  $S_m$ . Let  $m_1 \geq m_2 \geq \dots \geq m_l$  be nonnegative integers. Then a  $S_{m_1}, S_{m_2}, \dots, S_{m_l}$ -decomposition of  $K_n$  (or a *star decomposition* of  $K_n$ , for short) is a collection of stars such that

$$E(S_{m_i}) \cap E(S_{m_j}) = \emptyset \text{ if } i \neq j, \text{ and } \bigcup_{i=1}^l E(S_{m_i}) = E(K_n).$$

It was recently shown in [2] that such a decomposition exists if and only if

$$\sum_{i=1}^k m_i \leq \sum_{i=1}^k (n-i) \text{ for } k = 1, 2, \dots, n-1, \text{ and } \sum_{i=1}^l m_i = \binom{n}{2}.$$

An *automorphism* of a star decomposition is a permutation of  $V(K_n)$  which fixes the set  $\{S_{m_1}, S_{m_2}, \dots, S_{m_l}\}$ . The *orbit* of a star under an automorphism  $\pi$  is the collection of images of the star under the powers of  $\pi$ . A permutation of  $V(K_n)$  which consists of a single cycle of length  $n$  is said to be *cyclic*. A permutation of  $V(K_n)$  consisting of a fixed point and a cycle of length  $n-1$  is said to be *rotational*. Several graph and digraph decompositions have been studied which admit either a cyclic or rotational automorphism. See, for example, [1, 3, 4, 5]. The purpose of this paper is to give necessary and sufficient conditions for the existence of star decompositions of  $K_n$  which admit either a cyclic automorphism or a rotational automorphism.

## 2 Cyclic Star Decompositions of $K_n$

Throughout this section, we assume the vertex set of  $K_n$  is  $\{0, 1, \dots, n-1\}$  and we will construct star decompositions of  $K_n$  admitting  $\pi = (0, 1, \dots, n-1)$

1) as an automorphism.

**Lemma 2.1** *If there exists a  $S_{m_1}, S_{m_2}, \dots, S_{m_l}$ -decomposition of  $K_n$  which admits a cyclic automorphism and if  $n$  is even, then  $|\{i \mid m_i = 1\}| \equiv n/2 \pmod{n}$ .*

**Proof.** The edge  $(0, n/2)$  must lie in some star, say  $S_{m_s}$ . Then  $\pi^{n/2}((0, n/2)) = (0, n/2)$  and since each edge occurs in exactly one star of the decomposition, it must be that  $\pi^{n/2}(S_{m_s}) = S_{m_s}$ . Therefore  $m_s = 1$ . Let  $A = \{\pi^i(S_{m_s}) \mid i \in \mathbb{Z}\}$ . Then  $|A| = n/2$  and if  $S_{m_t} \notin A$  then the length of the orbit of  $S_{m_t}$  is  $n$ . Therefore  $|\{i \mid m_i = 1\}| \equiv n/2 \pmod{n}$ . ■

As argued in Lemma 2.1, the length of the orbit of every star in a cyclic star decomposition of  $K_n$  is  $n$  except for the special "short orbit" stars in set  $A$ . We therefore have:

**Lemma 2.2** *If there exists a  $S_{m_1}, S_{m_2}, \dots, S_{m_l}$ -decomposition of  $K_n$  which admits a cyclic automorphism, then for  $k = 1, 2, \dots, n-1$ ,  $|\{i \mid m_i = k\}| \equiv 0 \pmod{n}$ , except for the case  $k = 1$  when  $n$  is even.*

We show the necessary conditions of Lemmas 2.1 and 2.2, along with the necessary conditions for the existence of a star decomposition of  $K_n$ , are sufficient for the existence of a cyclic star decomposition of  $K_n$ .

**Theorem 2.1** *Let  $m_1 \geq m_2 \geq \dots \geq m_l$  be nonnegative integers. Then there is a cyclic  $S_{m_1}, S_{m_2}, \dots, S_{m_l}$ -decomposition of  $K_n$  if and only if*

$$\sum_{i=1}^k m_i \leq \sum_{i=1}^k (n-i) \text{ for } k = 1, 2, \dots, n-1, \quad \sum_{i=1}^l m_i = \binom{n}{2}$$

and

- (a)  $|\{i \mid m_i = k\}| \equiv 0 \pmod{n}$  for all  $k = 1, 2, \dots, n-1$  if  $n$  is odd, or
- (b)  $|\{i \mid m_i = 1\}| \equiv n/2 \pmod{n}$  and  $|\{i \mid m_i = k\}| \equiv 0 \pmod{n}$  for all  $k = 2, 3, \dots, n-1$  if  $n$  is even.

**Proof.** We need only establish sufficiency. Without loss of generality, we may assume  $m_l \geq 1$ . If  $n$  is odd, consider the collection of stars with edge sets

$$E(S_{m_l - kn - i}) = \{(i, i + r + \sum_{j=1}^k m_{l-(j-1)n}) \mid r = 1, 2, \dots, m_l - kn\}$$

for  $i = 0, 1, \dots, n-1$  and  $k = 0, 1, \dots, l/n - 1$ . If  $n$  is even, consider the collection of stars with edge sets

$$E(S_{m_{l-n/2-i}}) = \{(i, i + n/2)\}$$

for  $i = 0, 1, \dots, n/2 - 1$ , and

$$E(S_{m_{l-n/2-kn-i}}) = \{(i, i+r+\sum_{j=1}^k m_{l-n/2-(j-1)n}) \mid r = 1, 2, \dots, m_{l-n/2-kn}\}$$

for  $i = 0, 1, \dots, n-1$  and  $k = 0, 1, \dots, (l - n/2)/n - 1$ . In each case, the given collection of stars forms a cyclic star decomposition of  $K_n$ . ■

### 3 Rotational Star Decompositions of $K_n$

Throughout this section, we assume the vertex set of  $K_n$  is  $\{\infty, 0, 1, \dots, n-2\}$  and we will construct star decompositions of  $K_n$  admitting  $\pi = (\infty)(0, 1, \dots, n-2)$  as an automorphism.

As in Lemma 2.1, if  $n-1$  is even, then the edge  $(0, (n-1)/2)$  must occur in some  $S_{m_s}$  where  $m_s = 1$ . We analogously have:

**Lemma 3.1** *If there exists a  $S_{m_1}, S_{m_2}, \dots, S_{m_l}$ -decomposition of  $K_n$  which admits a rotational automorphism and if  $n$  is odd, then  $|\{i \mid m_i = 1\}| \equiv (n-1)/2 \pmod{n-1}$ .*

The orbit of each star of a rotational star decomposition of  $K_n$  is of length  $n-1$ , with two possible types of exceptions: (1) if  $n$  is odd, then the stars  $S_1$  with edge sets  $\{(i, i + (n-1)/2)\}$  for some  $i$  have orbits of length  $(n-1)/2$ , and (2) if  $m \mid (n-1)$ ,  $m \neq 1$ , say  $(n-1)/m = p$  then the stars  $S_m$  with edge sets  $\{(\infty, i), (\infty, i+p), \dots, (\infty, i+n-1-p)\}$  for some  $i$  have orbits of length  $p$ .

**Theorem 3.2** *Let  $m_1 \geq m_2 \geq \dots \geq m_l$  be nonnegative integers. Then there is a rotational  $S_{m_1}, S_{m_2}, \dots, S_{m_l}$ -decomposition of  $K_n$  if and only if*

$$\sum_{i=1}^k m_i \leq \sum_{i=1}^k (n-i) \text{ for } k = 1, 2, \dots, n-1, \quad \sum_{i=1}^l m_i = \binom{n}{2}$$

and

(a)  $|\{i \mid m_i = k\}| \equiv 0 \pmod{n-1}$  for all  $k = 1, 2, \dots, n-1$  if  $n$  is even,  
or

- (b)  $|\{i \mid m_i = 1\}| \equiv (n-1)/2 \pmod{n-1}$  and  $|\{i \mid m_i = k\}| \equiv 0 \pmod{n-1}$  for all  $k = 2, 3, \dots, n-1$  if  $n$  is odd, or
- (c) if  $m \mid (n-1)$ , say  $(n-1)/m = p$ , for some  $m \in \{m_1, m_2, \dots, m_l\}$ ,  $m \neq 1$ , then  $|\{i \mid m_i = m\}| \equiv p \pmod{n-1}$  and  $|\{i \mid m_i = k\}| \equiv 0 \pmod{n-1}$  for all  $k = 1, 2, \dots, m-1, m+1, \dots, n-1$  if  $n$  is even, or
- (d) if  $m \mid (n-1)$ , say  $(n-1)/m = p$ , for some  $m \in \{m_1, m_2, \dots, m_l\}$ ,  $m \neq 1$ , then  $|\{i \mid m_i = m\}| \equiv p \pmod{n-1}$ ,  $|\{i \mid m_i = 1\}| \equiv (n-1)/2 \pmod{n-1}$  and  $|\{i \mid m_i = k\}| \equiv 0 \pmod{n-1}$  for all  $k = 2, 3, \dots, m-1, m+1, \dots, n-1$  if  $n$  is odd.

**Proof.** We need only establish sufficiency. Without Loss of generality, we may assume  $m_l \geq 1$ . We consider the four cases separately.

(a) Consider the collection of stars with edge sets

$$E(S_{m_l-i}) = \{(\infty, i)\} \cup \{(i, i+r) \mid r = 1, 2, \dots, m_l-1\}$$

for  $i = 0, 1, \dots, n-2$  and

$$E(S_{m_l-k(n-1)-i}) = \{(i, i+r-1 + \sum_{j=1}^k m_{l-(j-1)(n-1)}) \mid r = 1, 2, \dots, m_{l-k(n-1)}\}$$

for  $i = 0, 1, \dots, n-2$  and  $k = 1, 2, \dots, l/(n-1) - 1$ .

(b) Consider the collection of stars with edge sets

$$E(S_{m_l-i}) = \{(i, i + (n-1)/2)\}$$

for  $i = 0, 1, \dots, (n-1)/2 - 1$ ,

$$E(S_{m_l-(n-1)/2-i}) = \{(\infty, i)\} \cup \{(i, i+r) \mid r = 1, 2, \dots, m_{l-(n-1)/2-k(n-1)} - 1\}$$

for  $i = 0, 1, \dots, n-2$ , and

$$E(S_{m_l-(n-1)/2-k(n-1)-i}) = \{(i, i+r-1 + \sum_{j=1}^k m_{l-(n-1)/2-(j-1)(n-1)}) \mid r = 1, 2, \dots, m_{l-(n-1)/2-k(n-1)}\}$$

for  $i = 0, 1, \dots, n-2$  and  $k = 1, 2, \dots, (l - (n-1)/2)/(n-1) - 1$ .

- (c) Let  $t$  be the largest index such that  $m_t = m$ . Consider the collection of stars with edge sets

$$E(S_{m_{l-k(n-1)-i}}) = \{(i, i + r + \sum_{j=1}^k m_{l-(j-1)(n-1)}) \mid r = 1, 2, \dots, m_{l-k(n-1)}\}$$

for  $i = 0, 1, \dots, n-2$  and  $k = 0, 1, \dots, (l-t)/(n-1) - 1$ ,

$$E(S_{m_{t-i}}) = \{(\infty, i + rp) \mid r = 0, 1, \dots, m_t - 1\}$$

for  $i = 0, 1, \dots, p-1$ ,

$$E(S_{m_{t-p-k(n-1)-i}}) = \{(i, i + r + \sum_{j=1}^{(l-t)/(n-1)} m_{l-(j-1)(n-1)} + \sum_{j=1}^k m_{t-p-(j-1)(n-1)}) \mid r = 1, 2, \dots, m_{t-p-k(n-1)}\}$$

for  $i = 0, 1, \dots, n-2$  and  $k = 0, 1, \dots, (t-p)/(n-1) - 1$ .

- (d) Let  $t$  be the largest index such that  $m_t = m$ . Consider the collection of stars with edge sets

$$E(S_{m_{l-i}}) = \{(i, i + (n-1)/2)\}$$

for  $i = 0, 1, \dots, (n-1)/2 - 1$ ,

$$E(S_{m_{l-(n-1)/2-k(n-1)-i}}) = \{(i, i + r + \sum_{j=1}^k m_{l-(n-1)/2-(j-1)(n-1)}) \mid r = 1, 2, \dots, m_{l-(n-1)/2-k(n-1)}\}$$

for  $i = 0, 1, \dots, n-2$  and  $k = 0, 1, \dots, (l-t)/(n-1) - 1$ ,

$$E(S_{m_{t-i}}) = \{(\infty, i + rp) \mid r = 0, 1, \dots, m_t - 1\}$$

for  $i = 0, 1, \dots, p-1$ ,

$$E(S_{m_{t-p-(n-1)/2-k(n-1)-i}}) = \{(i, i + r + \sum_{j=1}^{(l-t)/(n-1)} m_{l-(n-1)/2-(j-1)(n-1)} + \sum_{j=1}^k m_{t-p-(n-1)/2-(j-1)(n-1)}) \mid r = 1, 2, \dots, m_{t-p-(n-1)/2-k(n-1)}\}$$

for  $i = 0, 1, \dots, n-2$  and  $k = 0, 1, \dots, (t-p-(n-1)/2)/(n-1) - 1$ .

In each case, the given stars form a rotational decomposition of  $K_n$ . ■

## References

- [1] C. J. Cho, Rotational Mendelsohn Triple Systems, *Kyungpook Math J.* **20** (1986), 5-9.
- [2] C. Lin and T. Shyu, A Necessary and Sufficient Condition for the Star Decomposition of Complete Graphs, *J. Graph Theory* **23**(4) (1996), 361-364.
- [3] B. Micale and M. Pennisi, Cyclic Mendelsohn Quadruple Systems, *Ars Combinatoria* **35** (1993), 225-236.
- [4] R. Peltesohn, A Solution to both of Heffter's Difference Problems (in German), *Compositio Math.* **6** (1939), 251-257.
- [5] M. Pennisi, On the Rotational Mendelsohn Designs, *J. Combinatorics, Information and Systems Sciences*, to appear.